

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
5 December 2002 (05.12.2002)

PCT

(10) International Publication Number  
**WO 02/097127 A2**

(51) International Patent Classification<sup>7</sup>: **C12Q 1/68**

(21) International Application Number: PCT/EP02/05835

(22) International Filing Date: 28 May 2002 (28.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0113266.1 31 May 2001 (31.05.2001) GB

(71) Applicant (for all designated States except US): **BAYER AKTIENGESELLSCHAFT** [DE/DE]; 51368 Leverkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **OELLERS, Nadja** [DE/DE]; Steinfeldergasse 9, 50670 Köln (DE). **GEHRMANN, Mathias** [DE/DE]; Rosenstr. 68, 47918 Tönisvorst (DE). **KALLABIS, Harald** [DE/DE]; Wolfskaul 8, 51061 Köln (DE). **HALL, Roderick** [GB/GB]; 10 Acacia Close, Woodham, Surrey KT 15 3SJ (GB). **SCHULZE, Thomas** [DE/DE]; Kaiserwaldstr. 2c, 91083 Baiersdorf (DE). **KROEGEL, Claus** [DE/DE]; Falkenweg 18, 07751 Cospeda (DE).

(74) Common Representative: **BAYER AKTIENGESELLSCHAFT**; 51368 Leverkusen (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,

LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GENES AND PROTEINS FOR PREVENTION, PREDICTION, DIAGNOSIS, PROGNOSIS AND TREATMENT OF CHRONIC LUNG DISEASE

(57) Abstract: Genes that are differentially expressed in lung tissue of COPD patients versus lungs of normal people are disclosed. The genes provide novel methods for the prevention, prediction, diagnosis, prognosis and treatment of chronic lung disease.



WO 02/097127 A2

**GENES AND PROTEINS FOR PREVENTION, PREDICTION, DIAGNOSIS,  
PROGNOSIS AND TREATMENT OF CHRONIC LUNG DISEASE**

5      **TECHNICAL FIELD OF THE INVENTION**

The invention relates to polynucleotides and, more particularly to genes that are differentially expressed in diseased lungs as compared to normal lung. In particular genes that are up- or downregulated in lungs of patients with Chronic Obstructive  
10      Pulmonary Disease (COPD) are disclosed. Also disclosed are methods for utilizing such genes, polynucleotides or polypeptides derived from the genes as predictive, diagnostic or prognostic markers for chronic lung disease, particularly COPD. Also disclosed are antibodies specific to the markers to be used to identify the genes or their protein products for predictive, diagnostic or prognostic purposes or to  
15      modulate their activity in order to prevent or treat lung disease. Methods of screening for modulators (activators or inhibitors) which can be used for the regulation of the genes or the polypeptides derived therefrom and preventive or therapeutic uses of these modulators are also disclosed.

20      **BACKGROUND OF THE INVENTION**

Chronic obstructive pulmonary (or airways) disease (COPD) is a condition defined physiologically as airflow obstruction that generally results from a mixture of emphysema and peripheral airway obstruction due to chronic bronchitis (1).  
25      Emphysema is characterised by destruction of alveolar walls leading to abnormal enlargement of the air spaces of the lung. Chronic bronchitis is defined clinically as the presence of chronic productive cough for three months in each of two successive years. In COPD, airflow obstruction is usually progressive and is only partially reversible. By far the most important risk factor for development of COPD is  
30      cigarette smoking, although the disease does occur in non-smokers.

Chronic inflammation of the airways is a key pathological feature of COPD (1). The inflammatory cell population comprises increased numbers of macrophages, neutrophils and CD8<sup>+</sup> lymphocytes. Inhaled irritants such as cigarette smoke activate macrophages resident in the respiratory tract as well as epithelial cells leading to release of chemokines (e.g. interleukin-8) and other chemotactic factors which act to increase the neutrophil/monocyte trafficking from the blood into the lung tissue and airways. Neutrophils and monocytes recruited into the airways can release a variety of potentially damaging mediators such as proteolytic enzymes and reactive oxygen species. Matrix degradation and emphysema, along with airway wall thickening, surfactant dysfunction and mucus hypersecretion are all potential sequelae of this inflammatory response that lead to impaired airflow and gas exchange.

Since there is a lack of the understanding of the underlying mechanisms of COPD, the therapeutic standard is not very high. COPD is the 5<sup>th</sup> leading cause of death in the world and the need for drugs is extremely high. The main risk factor for development of COPD is cigarette smoking. But only a portion (up to 20%) of all smokers develop COPD, independently of their smoking habits. A genetic predisposition of a portion of smokers to develop COPD is very likely. COPD is a subgroup of the chronic lung diseases which includes also asthma and which are characterized by a chronic inflammation and/or fibrosis of the airway tissue. Many pathophysiological features are shared among these diseases. There is a need for a better understanding of the pathophysiology of chronic lung diseases on the molecular level in order to identify those genes that could serve as novel predictive, diagnostic and/ or prognostic markers and/ or as targets for preventive and/ or therapeutic drugs for chronic lung disease. Certain genes involved in the pathophysiology of chronic lung disease have been shown to be (a) upregulated in animal models of disease (108) and human patients suffering from chronic lung disease (109) or (b) downregulated in human patients suffering from chronic lung disease (110). Consequently there is a demand to identify more genes that are implicated in the disease process since novel preventive, predictive, diagnostic, prognostic and therapeutic methods can be based on them.

### SUMMARY OF THE INVENTION

5 The present invention is based on the identification of genes that are differentially expressed in lung tissue of patients with a clear clinical evidence of COPD compared to lung tissue of individuals without any evidence of COPD. Accordingly disclosed herein are 28 genes that are differentially expressed in COPD, as well as derivatives, fragments, analogs and homologues thereof. Any of these genes is named hereinafter “COPD GENE”.

10 It is an objective of the invention to provide methods and reagents for the prevention, prediction, diagnosis, prognosis and treatment of chronic lung disease and COPD in particular.

15 In one embodiment of the invention a “COPD GENE” or a gene product of a “COPD GENE” can be used to identify cells or tissue in individuals which exhibit a phenotype predisposed to disease or a diseased phenotype, thereby (a) predicting whether an individual is at risk for the development, or (b) diagnosing whether an individual is having, or (c) prognosing the progression or the outcome of the treatment of chronic lung disease and COPD in particular.

20 In another embodiment the invention provides methods of screening for agents which regulate the activity of a polypeptide encoded by a “COPD GENE”. A test compound is contacted with a polypeptide encoded by a “COPD GENE”. Binding of the test compound to the polypeptide is detected. A test compound which binds to the polypeptide is thereby identified as a potential therapeutic agent for the treatment of chronic lung disease and more particularly COPD.

25 In one embodiment the invention provides another method of screening for agents which regulate the activity of a polypeptide encoded by a “COPD GENE”. A test compound is contacted with a polypeptide encoded by a “COPD GENE”. A



biological activity mediated by the polypeptide is detected. A test compound which decreases the biological activity is thereby identified as a potential therapeutic agent for decreasing the activity of the polypeptide encoded by a "COPD GENE" in chronic lung disease and COPD in particular. A test compound which increases the biological activity is thereby identified as a potential therapeutic agent for increasing the activity of the polypeptide encoded by a "COPD GENE" in chronic lung disease and COPD in particular.

In another embodiment the invention provides a method of screening for agents which regulate the activity of a "COPD GENE". A test compound is contacted with a "COPD GENE" polynucleotide. Binding of the test compound to the "COPD GENE" polynucleotide is detected. A test compound which binds to the polynucleotide is thereby identified as a potential therapeutic agent for regulating the activity of the "COPD GENE" in chronic lung disease and COPD in particular.

The invention thus provides "COPD GENES" which can be used to identify compounds which may act, for example, as regulators or modulators such as agonists and antagonists, partial agonist, inverse agonist, activators, co-activators and inhibitors of the polypeptide encoded by a "COPD GENE". Accordingly, the invention provides reagents and methods for regulating a "COPD GENE" polynucleotide or a polypeptide encoded by a "COPD GENE" in chronic lung disease and more particularly COPD. The regulation can be up- or downregulation. Reagents that modulate the expression, stability or amount of a "COPD GENE" polynucleotide or the activity of the polypeptide encoded by a "COPD GENE" can be a protein, a peptide, a peptidomimetic, a nucleic acid, a nucleic acid analogue (e.g. peptide nucleic acid, locked nucleic acid) or a small molecule. Methods that the expression, stability or amount of a "COPD GENE" polynucleotide or the activity of the polypeptide encoded by a "COPD GENE" can be gene replacement therapies, antisense, ribozyme and triplex nucleic acid approaches.

One embodiment of the invention provides antibodies which specifically bind to a full-length or partial "COPD GENE" for use in prevention, diagnosis, prognosis and treatment of chronic lung disease.

5 Yet another embodiment of the invention is the use of a reagent which specifically binds to a "COPD GENE" polynucleotide or a polypeptide encoded by a "COPD GENE" in the preparation of a medicament for the treatment of chronic lung disease and COPD in particular.

10 Still another embodiment of the invention is a pharmaceutical composition which includes a reagent which specifically binds to a "COPD GENE" polynucleotide or a polypeptide encoded by a "COPD GENE", and a pharmaceutically acceptable carrier.

#### DETAILED DESCRIPTION OF THE INVENTION

15

The present invention relates to 28 genes that are differentially regulated in the lungs of patients with clinical evidence of COPD. "Gene or Genes" as used herein refers to the polynucleotides of SEQ ID NO. 1 to 28, as well as derivatives, fragments, analogs and homologues thereof, the polypeptides encoded thereby, the polypeptides  
20 of SEQ ID NO. 29 to 56 and the corresponding genomic transcription units which can be derived or identified with standard techniques well known in the art using the information disclosed in Tables 1 to 5. The accession numbers of the polynucleotide sequences of the SEQ IDs NO. 1 to 28 are shown in Table 3. The accession numbers of the polypeptide sequences encoded by the polynucleotide sequences are also  
25 shown in Table 3.

The invention further relates to the use of:

- 30 a) a polynucleotide comprising at least one of the sequences of SEQ ID NO. 1 to 28;

- b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) and encodes a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;
- 5 c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;
- d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);
- 10 e) an antisense molecule targeting one of the polynucleotide sequences specified in (a) to (d);
- f) a purified polypeptide encoded by a polynucleotide sequence specified in (a) to (d)
- 15 g) a purified polypeptide comprising at least one of the sequences of SEQ ID NO. 29 to 56;
- 20 h) an antibody capable of binding to one of the polynucleotide specified in (a) to (d) or a polypeptide specified in (f) and (g)
- i) a reagent identified by any of the methods as specified below that modulates the amount or activity of a polynucleotide sequence specified in (a) to (d) or a polypeptide specified in (f) and (g)
- 25

for the prevention, prediction, diagnosis, prognosis and treatment of chronic lung disease.

### Polynucleotides

A „COPD GENE“ polynucleotide can be single- or double-stranded and comprises a coding sequence or the complement of a coding sequence for a „COPD GENE“ polypeptide. Degenerate nucleotide sequences encoding human „COPD GENE“ polypeptides, as well as homologous nucleotide sequences which are at least about 50, 55, 60, 65, 70, preferably about 75, 90, 96, or 98% identical to the nucleotide sequences of SEQ ID NO. 1 to 28 also are „COPD GENE“ polynucleotides. Percent sequence identity between the sequences of two polynucleotides is determined using computer programs such as ALIGN which employ the FASTA algorithm, using an affine gap search with a gap open penalty of -12 and a gap extension penalty of -2. Complementary DNA (cDNA) molecules, species homologues, and variants of „COPD GENE“ polynucleotides which encode biologically active „COPD GENE“ polypeptides also are „COPD GENE“ polynucleotides.

### Identification of Polynucleotide Variants and Homologues

Variants and homologues of the „COPD GENE“ polynucleotides described above also are „COPD GENE“ polynucleotides. Typically, homologous „COPD GENE“ polynucleotide sequences can be identified by hybridization of candidate polynucleotides to known „COPD GENE“ polynucleotides under stringent conditions, as is known in the art. For example, using the following wash conditions: 2X SSC (0.3 M NaCl, 0.03 M sodium citrate, pH 7.0), 0.1% SDS, room temperature twice, 30 minutes each; then 2X SSC, 0.1% SDS, 50 EC once, 30 minutes; then 2X SSC, room temperature twice, 10 minutes each homologous sequences can be identified which contain at most about 25-30% basepair mismatches. More preferably, homologous nucleic acid strands contain 15-25% basepair mismatches, even more preferably 5-15% basepair mismatches.

Species homologues of the „COPD GENE“ polynucleotides disclosed herein also can be identified by making suitable probes or primers and screening cDNA expression

libraries from other species, such as mice, monkeys, or yeast. Human variants of „COPD GENE“ polynucleotides can be identified, for example, by screening human cDNA expression libraries. It is well known that the  $T_m$  of a double-stranded DNA decreases by 1-1.5°C with every 1% decrease in homology (2). Variants of human „COPD GENE“ polynucleotides or „COPD GENE“ polynucleotides of other species can therefore be identified by hybridizing a putative homologous „COPD GENE“ polynucleotide with a polynucleotide comprising any of the nucleotide sequences of SEQ ID NO. 1 to 28 or the complement thereof to form a test hybrid. The melting temperature of the test hybrid is compared with the melting temperature of a hybrid comprising polynucleotides having perfectly complementary nucleotide sequences, and the number or percent of basepair mismatches within the test hybrid is calculated.

Nucleotide sequences which hybridize to „COPD GENE“ polynucleotides or their complements following stringent hybridization and/or wash conditions also are „COPD GENE“ polynucleotides. Stringent wash conditions are well known and understood in the art and are disclosed, for example in Ref. 3 at pages 9.50-9.51.

Typically, for stringent hybridization conditions a combination of temperature and salt concentration should be chosen that is approximately 12-20°C below the calculated  $T_m$  of the hybrid under study. The  $T_m$  of a hybrid between a „COPD GENE“ polynucleotide comprising a nucleotide sequence selected from the polynucleotides of SEQ ID NO. 1 to 28 or the complement thereof and a polynucleotide sequence which is at least about 50, preferably about 75, 90, 96, or 98% identical to one of those nucleotide sequences can be calculated, for example, using the equation of Bolton and McCarthy (4):

$$T_m = 81.5^\circ\text{C} - 16.6(\log_{10}[\text{Na}^+]) + 0.41(\%G + C) - 0.63(\%\text{formamide}) - 600/l,$$

where  $l$  = the length of the hybrid in basepairs.

Stringent wash conditions include, for example, 4X SSC at 65°C, or 50% formamide, 4X SSC at 42°C, or 0.5X SSC, 0.1% SDS at 65°C. Highly stringent wash conditions include, for example, 0.2X SSC at 65°C.

5     Preparation of Polynucleotides

A naturally occurring „COPD GENE“ polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the  
10     polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique for obtaining a polynucleotide can be used to obtain isolated „COPD GENE“ polynucleotides. For example, restriction enzymes and probes can be used to isolate  
15     polynucleotide fragments which comprise „COPD GENE“ nucleotide sequences. Isolated polynucleotides are in preparations which are free or at least 70, 80, or 90% free of other molecules.

„COPD GENE“ cDNA molecules can be made with standard molecular biology techniques, using „COPD GENE“ mRNA as a template. „COPD GENE“ cDNA  
20     molecules can thereafter be replicated using molecular biology techniques known in the art and disclosed in manuals such as Sambrook et al. (3). An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

25     Alternatively, synthetic chemistry techniques can be used to synthesize „COPD GENE“ polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode a „COPD GENE“ polypeptide or a biologically active variant thereof.

30

Extending Polynucleotides

Various PCR-based methods can be used to extend the nucleic acid sequences disclosed herein to detect upstream sequences such as promoters and regulatory elements. For example, restriction site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus (5). Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region (6). Primers can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 2230 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA (7). In this method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

Another method which can be used to retrieve unknown sequences is that of Parker et al (8). Additionally, PCR, nested primers, and PROMOTERFINDER libraries (CLONTECH, Palo Alto, Calif.) can be used to walk genomic DNA (CLONTECH,

Palo Alto, Calif.). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' nontranscribed regulatory regions.

Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate software (e.g. GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic data display can be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

#### Polypeptides

“COPD GENE” polypeptides according to the invention comprise an amino acid selected from SEQ ID NO. 29 to 56 or which are encoded by the polynucleotide sequences of SEQ ID NO. 1 to 28 or derivatives, fragments, analogs and homologues thereof. A “COPD GENE” polypeptide of the invention therefore can be a portion, a full-length, or a fusion protein comprising all or a portion of a “COPD GENE” polypeptide.



The invention additionally, encompasses "COPD GENE" polypeptides which are differentially modified during or after translation, e.g. by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc.

5 Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH<sub>4</sub>; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

10 Additional post-translational modifications encompassed by the invention include, for example, e.g., N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of procaryotic  
15 host cell expression. The "COPD GENE" polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the protein.

Also provided by the invention are chemically modified derivatives of the "COPD  
20 GENE" polypeptides which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Patent No. 4,179,337). The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose,  
25 dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

Biologically Active Variants

„COPD GENE“ polypeptide variants which are biologically active, i.e., retain an „COPD GENE“ activity, also are „COPD GENE“ polypeptides. Preferably, naturally or non-naturally occurring „COPD GENE“ polypeptide variants have amino acid sequences which are at least about 60, 65, or 70, preferably about 75, 80, 85, 90, 92, 94, 96, or 98% identical to the amino acid sequence which is encoded by a polynucleotide comprising at least one of the polynucleotide sequences of SEQ ID NO. 1 to 28 or which comprises at least one polypeptide of SEQ ID NO. 29 to 56 or a fragment thereof. Percent identity between a putative „COPD GENE“ polypeptide variant and an amino acid sequence which is encoded by a polynucleotide comprising at least one of the polynucleotide sequences of SEQ ID NO. 1 to 28 or which comprises at least one polypeptide of SEQ ID NO. 29 to 56 is determined using the Needleman/Wunsch algorithm (111) with the substitutions-matrix BLOSUM62 (112) and a gap creation penalty of 8 and a gap extension penalty of 2.

Variations in percent identity can be due, for example, to amino acid substitutions, insertions, or deletions. Amino acid substitutions are defined as one for one amino acid replacements. They are conservative in nature when the substituted amino acid has similar structural and/or chemical properties. Examples of conservative replacements are substitution of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

Amino acid insertions or deletions are changes to or within an amino acid sequence. They typically fall in the range of about 1 to 5 amino acids. Guidance in determining which amino acid residues can be substituted, inserted, or deleted without abolishing biological or immunological activity of a „COPD GENE“ polypeptide can be found using computer programs well known in the art, such as DNASTAR software. Whether an amino acid change results in a biologically active „COPD GENE“ polypeptide can readily be determined by assaying for „COPD GENE“ activity, as described for example, in the specific examples, below. Larger insertions or deletions

can also be caused by alternative splicing. Protein domains can be inserted or deleted without altering the main activity of the protein.

### Fusion Proteins

5

Fusion proteins are useful for generating antibodies against „COPD GENE“ polypeptide amino acid sequences and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of a „COPD GENE“ polypeptide. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

10

A „COPD GENE“ polypeptide fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment comprises at least 25, 50, 75, 100, 150, 200, 300, 400, 500, 600, 700 or 750 contiguous amino acids of an amino acid sequence encoded by a polynucleotide comprising at least one of the polynucleotide sequences of SEQ ID NO. 1 to 28 or of a biologically active variant, such as those described above. The first polypeptide segment also can comprise a full-length „COPD GENE“ polypeptide.

15

20

The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include  $\beta$ -galactosidase,  $\beta$ -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. A

25

30

fusion protein also can be engineered to contain a cleavage site located between the „COPD GENE“ polypeptide-encoding sequence and the heterologous protein sequence, so that the „COPD GENE“ polypeptide can be cleaved and purified away from the heterologous moiety.

5

A fusion protein can be synthesized chemically, as is known in the art. Preferably, a fusion protein is produced by covalently linking two polypeptide segments or by standard procedures in the art of molecular biology. Recombinant DNA methods can be used to prepare fusion proteins, for example, by making a DNA construct which comprises coding sequences selected from polynucleotides comprising at least one of the polynucleotide sequences of SEQ ID NO. 1 to 28 in proper reading frame with nucleotides encoding the second polypeptide segment and expressing the DNA construct in a host cell, as is known in the art. Many kits for constructing fusion proteins are available from companies such as Promega Corporation (Madison, WI), Stratagene (La Jolla, CA), CLONTECH (Mountain View, CA), Santa Cruz Biotechnology (Santa Cruz, CA), MBL International Corporation (MIC; Watertown, MA), and Quantum Biotechnologies (Montreal, Canada; 1-888-DNA-KITS).

10

15

#### Obtaining Polypeptides

20

„COPD GENE“ polypeptides can be obtained, for example, by purification from human cells, by expression of „COPD GENE“ polynucleotides, or by direct chemical synthesis.

25

#### Protein Purification

30

„COPD GENE“ polypeptides can be purified from any cell which expresses the enzyme, including host cells which have been transfected with „COPD GENE“ expression constructs. Lung is an especially useful source of „COPD GENE“ polypeptides. A purified „COPD GENE“ polypeptide is separated from other compounds which normally associate with the „COPD GENE“ polypeptide in the cell, such as

certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis. A preparation of purified „COPD GENE“ polypeptides is at least 80% pure; preferably, the preparations are 90%, 95%, or 99% pure. Purity of the preparations can be assessed by any means known in the art, such as SDS-polyacrylamide gel electrophoresis.

#### Expression of Polynucleotides

10

To express a „COPD GENE“ polynucleotide, the polynucleotide can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding „COPD GENE“ polypeptides and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described, for example, in Sambrook et al. (3) and in Ausubel et al. (9). A variety of expression vector/host systems can be utilized to contain and express sequences encoding a „COPD GENE“ polypeptide. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (e.g., baculovirus), plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids), or animal cell systems.

20

25

30

The control elements or regulatory sequences are those nontranslated regions of the vector enhancers, promoters, 5' and 3' untranslated regions which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any

number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and  
5 the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO, and storage protein genes) or from plant viruses (e.g., viral promoters or leader sequences) can be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is  
10 necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding a „COPD GENE“ polypeptide, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

#### Bacterial and Yeast Expression Systems

15 In bacterial systems, a number of expression vectors can be selected depending upon the use intended for the „COPD GENE“ polypeptide. For example, when a large quantity of the „COPD GENE“ polypeptide is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified  
20 can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding the „COPD GENE“ polypeptide can be ligated into the vector in frame with sequences for the amino terminal Met and the subsequent 7 residues of  $\beta$ -galactosidase so that a hybrid protein is produced. pIN  
25 vectors (Van Heeke & Schuster, J. Biol. Chem. 264, 55035509, 1989) or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione agarose beads followed by elution in the presence of free glutathione.  
30 Proteins made in such systems can be designed to include heparin, thrombin, or

factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

5 In the yeast *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH can be used. For reviews, see Ausubel et al. (9) and Grant et al. (10).

#### Plant and Insect Expression Systems

10 If plant expression vectors are used, the expression of sequences encoding „COPD GENE“ polypeptides can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV (11). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be  
15 used (12,13,14). These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection. Such techniques are described in a number of generally available reviews e.g., Hobbs or Murray (15).

20 An insect system also can be used to express a „COPD GENE“ polypeptide. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. Sequences encoding „COPD GENE“ polypeptides can be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of „COPD  
25 GENE“ polypeptides will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which „COPD GENE“ polypeptides can be expressed (16).

### Mammalian Expression Systems

A number of viral-based expression systems can be used to express „COPD GENE“ polypeptides in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding „COPD GENE“ polypeptides can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a nonessential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing a „COPD GENE“ polypeptide in infected host cells (17). If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (e.g., liposomes, polycationic amino polymers, or vesicles).

Specific initiation signals also can be used to achieve more efficient translation of sequences encoding „COPD GENE“ polypeptides. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding a „COPD GENE“ polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used (see Ref. 18).



### Host Cells

A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed „COPD GENE“ polypeptide in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Posttranslational processing which cleaves a "prepro" form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for Post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen to ensure the correct modification and processing of the foreign protein.

Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express „COPD GENE“ polypeptides can be transformed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 12 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced „COPD GENE“ sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. See, for example, ANIMAL CELL CULTURE, R.I. Freshney, ed., 1986.

Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (19) and adenine phosphoribosyltransferase (20) genes which can be employed in tk<sup>-</sup> or aprt<sup>-</sup> cells, respectively. Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, dhfr confers resistance to metho-

trexate (21), npt confers resistance to the aminoglycosides, neomycin and G418 (22), and als and pat confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murray, 1992, supra). Additional selectable genes have been described. For example, trpB allows cells to utilize indole in place of  
5 tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (23). Visible markers such as anthocyanins,  $\beta$ -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system (24).

10

#### Detecting Polynucleotide and Polypeptide Expression in Host Cells

Although the presence of marker gene expression suggests that the „COPD GENE“ polynucleotide is also present, its presence and expression may need to be confirmed.  
15 For example, if a sequence encoding a „COPD GENE“ polypeptide is inserted within a marker gene sequence, transformed cells containing sequences which encode a „COPD GENE“ polypeptide can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding a „COPD GENE“ polypeptide under the control of a single promoter.  
20 Expression of the marker gene in response to induction or selection usually indicates expression of the „COPD GENE“ polynucleotide.

Alternatively, host cells which contain a „COPD GENE“ polynucleotide and which express a „COPD GENE“ polypeptide can be identified by a variety of procedures  
25 known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of nucleic acid or protein. For example, the presence of a polynucleotide sequence encoding a „COPD GENE“ polypeptide can be detected  
30 by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding a „COPD GENE“ polypeptide.

Nucleic acid amplification-based assays involve the use of oligonucleotides selected from sequences encoding a „COPD GENE“ polypeptide to detect transformants which contain a „COPD GENE“ polynucleotide.

5 A variety of protocols for detecting and measuring the expression of a „COPD GENE“ polypeptide, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immuno-sorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal  
10 antibodies reactive to two non-interfering epitopes on a „COPD GENE“ polypeptide can be used, or a competitive binding assay can be employed. These and other assays are described in Hampton et al. (25) and (26).

A wide variety of labels and conjugation techniques are known by those skilled in the  
15 art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding „COPD GENE“ polypeptides include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, sequences encoding a „COPD GENE“ polypeptide can be cloned into  
20 a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes in vitro by addition of labeled nucleotides and an appropriate RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable  
25 reporter molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

### Expression and Purification of Polypeptides

Host cells transformed with nucleotide sequences encoding a „COPD GENE“ polypeptide can be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode „COPD GENE“ polypeptides can be designed to contain signal sequences which direct secretion of soluble „COPD GENE“ polypeptides through a prokaryotic or eukaryotic cell membrane or which direct the membrane insertion of membrane-bound „COPD GENE“ polypeptide.

As discussed above, other constructions can be used to join a sequence encoding a „COPD GENE“ polypeptide to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor Xa or enterokinase (Invitrogen, San Diego, CA) between the purification domain and the „COPD GENE“ polypeptide also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a „COPD GENE“ polypeptide and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography, as described in (27) while the enterokinase cleavage site provides a means for purifying the „COPD GENE“ polypeptide from the fusion protein. Vectors which contain fusion proteins are disclosed in (28).

### Chemical Synthesis

Sequences encoding a „COPD GENE“ polypeptide can be synthesized, in whole or in part, using chemical methods well known in the art (see Ref. 29, 30). Alternatively, a  
5 „COPD GENE“ polypeptide itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques (31, 32). Protein synthesis can be performed using manual techniques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Optionally,  
10 fragments of „COPD GENE“ polypeptides can be separately synthesized and combined using chemical methods to produce a full-length molecule.

The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography (e.g. Ref. 33). The composition of a synthetic  
15 „COPD GENE“ polypeptide can be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; see Creighton, supra). Additionally, any portion of the amino acid sequence of the „COPD GENE“ polypeptide can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

20

### Production of Altered Polypeptides

As will be understood by those of skill in the art, it may be advantageous to produce „COPD GENE“ polypeptide-encoding nucleotide sequences possessing non-natural  
25 occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

30 The nucleotide sequences disclosed herein can be engineered using methods generally known in the art to alter „COPD GENE“ polypeptide-encoding sequences

for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

#### Predictive, Diagnostic and Prognostic Methods and Reagents

10

The present invention provides methods and reagents for determining whether a subject is at risk for developing chronic lung disease or COPD in particular by detecting the disclosed biomarkers, i.e., the disclosed polynucleotides specified in Table 3 and/or polypeptides encoded thereby. One embodiment is a method for the prediction, diagnosis or prognosis of chronic lung disease by the detection:

15

- a) a polynucleotide comprising at least one of the sequences of SEQ ID NO.1 to 28;
- 20 b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;
- c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;
- 25 d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

in a biological sample comprising the following steps: hybridizing at least one polynucleotide specified in (a) to (d) to a nucleic acid material of a biological sample, thereby forming a hybridization complex; and detecting said hybridization complex.

- 5 Another embodiment is a method as described above, wherein before hybridization, the nucleic acid material of the biological sample is amplified.

Another embodiment is a method for the prediction, diagnosis or prognosis of chronic lung disease by the detection of:

10

a) a polynucleotide comprising at least one of the sequences of SEQ ID NO. 1 to 28;

15

b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;

20

c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;

d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

25

e) a purified polypeptide encoded by a polynucleotide sequence specified in (a) to (d);

f) a purified polypeptide comprising at least one of the sequences of SEQ ID NO. 29 to 56;

comprising the steps of contacting a biological sample with a reagent which specifically interacts with the polynucleotide specified in (a) to (d) or the polypeptide specified in (e) and (f).

- 5 Another embodiment is a diagnostic kit for conducting any of the methods described in the embodiments above.

In clinical applications, human tissue samples can be screened for the presence and/or absence of the biomarkers („COPD GENES“) identified herein. Such samples  
10 could consist of needle biopsy cores, surgical resection samples, lymph node tissue, whole blood, plasma or serum. For example, these methods include obtaining a biopsy, which is optionally fractionated by cryostat sectioning to enrich diseased cells to about 80% of the total cell population. In certain embodiments, nucleic acids extracted from these samples may be amplified using techniques well known in the  
15 art. The levels of selected markers can be detected in diseased lung and healthy lung tissue samples and compared within statistically valid reference groups. An abnormal level of the „COPD GENE“ polypeptide or mRNA levels is likely to be indicative of chronic lung disease such as COPD.

20 In even another embodiment the diagnostic method comprises determining whether a subject has an abnormal mRNA and/or protein level of the disclosed markers, such as by Northern blot analysis, reverse transcription-polymerase chain reaction (RT-PCR), *in situ* hybridization, immunoprecipitation, Western blot analysis, or immuno-histochemistry. According to the method, cells are obtained from a subject and the  
25 levels of the disclosed „COPD GENES“, protein or mRNA level, is determined and compared.

Accordingly, in one aspect, the invention provides probes and primers that are specific to the unique polynucleotide „COPD GENES“ disclosed herein. Accord-  
30 ingly, the polynucleotide probes comprise a nucleotide sequence at least 12 nucleotides in length, preferably at least 15 nucleotides, more preferably, 25



nucleotides, and most preferably at least 40 nucleotides, and up to all or nearly all of the coding sequence which is complementary to a portion of the coding sequence of a „COPD GENE“ polynucleotide sequence comprising at least one of the sequences of SEQ ID NO. 1 to 28 or a sequence complementary thereto.

5

In one embodiment, the method comprises using a polynucleotide probe to determine the presence of chronic lung disease cells in a tissue from a patient. Specifically, the method comprises:

- 10 1. providing a polynucleotide probe comprising a nucleotide sequence at least 12 nucleotides in length, preferably at least 15 nucleotides, more preferably, 25 nucleotides, and most preferably at least 40 nucleotides, and up to all or nearly all of the coding sequence which is complementary to a portion of the coding sequence of a polynucleotide sequence comprising at least one of the  
15 sequences of SEQ ID NO. 1 to 28 or a sequence complementary thereto and is differentially expressed in chronic lung disease, such as COPD ;
2. obtaining a tissue sample from a patient with chronic lung disease and COPD in particular;
- 20 3. providing reference tissue sample from patients with no lung disease;
4. contacting the polynucleotide probe under stringent conditions with RNA of each of said first and reference tissue samples (e.g., in a Northern blot or in  
25 situ hybridization assay); and
5. comparing (a) the amount of hybridization of the probe with RNA of the first tissue sample, with (b) the amount of hybridization of the probe with RNA of the reference tissue samples;

30

wherein a statistically significant difference in the amount of hybridization with the RNA of the first tissue sample as compared to the amount of hybridization of the RNA with the reference tissue samples is indicative of chronic lung disease and COPD in particular in the first tissue sample.

5

In one aspect, the method comprises in situ hybridization with a probe derived from a given „COPD GENE“ polynucleotide, which polynucleotide sequence comprises at least one of the polynucleotide sequences of SEQ ID NO. 1 to 28 or a sequence complementary thereto. The method comprises contacting the labeled hybridization probe with a sample of a given type of tissue from a patient potentially having chronic lung disease and COPD in particular as well as normal tissue from a person with no lung disease, and determining whether the probe labels tissue of the patient to a degree significantly different (e.g., by at least a factor of two, or at least a factor of five, or at least a factor of twenty, or at least a factor of fifty) than the degree to which normal tissue is labeled.

10  
15

Another such method includes the step of providing an antibody specific for the polypeptide encoded by a polynucleotide comprising at least one of the polynucleotide sequences of SEQ ID NO. 1 to 28.

20

The subject invention further provides a method of determining whether a cell sample obtained from a subject possesses an abnormal amount of „COPD GENE“ polypeptide which comprises (a) obtaining a cell sample from the subject, (b) quantitatively determining the amount of the „COPD GENE“ polypeptide in the sample so obtained, and (c) comparing the amount of the „COPD GENE“ polypeptide so determined with a known standard, so as to thereby determine whether the cell sample obtained from the subject possesses an abnormal amount of the „COPD GENE“ polypeptide. Such „COPD GENE“ polypeptide may be detected by immuno-histochemical assays, dot-blot assays, ELISA and the like.

25  
30

Immunoassays are commonly used to quantify the levels of proteins in cell samples, and many other immunoassay techniques are known in the art. The invention is not limited to a particular assay procedure, and therefore is intended to include both homogeneous and heterogeneous procedures. Exemplary immunoassays which can be conducted according to the invention include fluorescence polarization immunoassay (FPIA), fluorescence immunoassay (FIA), enzyme immunoassay (EIA), nephelometric inhibition immunoassay (NIA), enzyme linked immunosorbent assay (ELISA), and radioimmunoassay (RIA). An indicator moiety, or label group, can be attached to the subject antibodies and is selected so as to meet the needs of various uses of the method which are often dictated by the availability of assay equipment and compatible immunoassay procedures. General techniques to be used in performing the various immunoassays noted above are known to those of ordinary skill in the art.

In another embodiment, the level of the encoded product, i.e., the product encoded by a polynucleotide comprising at least one of the polynucleotides sequences of SEQ ID NO. 1 to 28 or a sequence complementary thereto, in a biological fluid (e.g., blood or urine) of a patient may be determined as a way of monitoring the level of expression of the „COPD GENE“ polynucleotide sequence in cells of that patient. Such a method would include the steps of obtaining a sample of a biological fluid from the patient, contacting the sample (or proteins from the sample) with an antibody specific for a encoded „COPD GENE“ polypeptide, and determining the amount of immune complex formation by the antibody, with the amount of immune complex formation being indicative of the level of the marker encoded product in the sample. This determination is particularly instructive when compared to the amount of immune complex formation by the same antibody in a control sample taken from a normal individual or in one or more samples previously or subsequently obtained from the same person.

Of particular importance to the subject invention is the ability to quantify the level of „COPD GENE“ polypeptide as determined by the number of cells associated with a

normal or abnormal „COPD GENE“ polypeptide level. The number of cells with a particular „COPD GENE“ polypeptide phenotype may then be correlated with patient prognosis. In one embodiment of the invention, the „COPD GENE“ polypeptide phenotype of the lesion is determined as a percentage of cells in a biopsy which are found to have abnormally high/low levels of the „COPD GENE“ polypeptide. Such expression may be detected by immunohistochemical assays, dot-blot assays, ELISA and the like.

Where tissue samples are employed, immunohistochemical staining may be used to determine the number of cells having the „COPD GENE“ polypeptide phenotype. For such staining, a multiblock of tissue is taken from the biopsy or other tissue sample and subjected to proteolytic hydrolysis, employing such agents as protease K or pepsin. In certain embodiments, it may be desirable to isolate a nuclear fraction from the sample cells and detect the level of the „COPD GENE“ polypeptide in the nuclear fraction.

The tissues samples are fixed by treatment with a reagent such as formalin, glutaraldehyde, methanol, or the like. The samples are then incubated with an antibody, preferably a monoclonal antibody, with binding specificity for the „COPD GENE“ polypeptides. This antibody may be conjugated to a label for subsequent detection of binding. Samples are incubated for a time sufficient for formation of the immunocomplexes. Binding of the antibody is then detected by virtue of a label conjugated to this antibody. Where the antibody is unlabeled, a second labeled antibody may be employed, e.g., which is specific for the isotype of the anti-„COPD GENE“ polypeptide antibody. Examples of labels which may be employed include radionuclides, fluorescers, chemilumescers, enzymes and the like.

Where enzymes are employed, the substrate for the enzyme may be added to the samples to provide a colored or fluorescent product. Examples of suitable enzymes for use in conjugates include horseradish peroxidase, alkaline phosphatase, malate dehydrogenase and the like. Where not commercially available, such antibody-

enzyme conjugates are readily produced by techniques known to those skilled in the art.

5 In one embodiment, the assay is performed as a dot blot assay. The dot blot assay finds particular application where tissue samples are employed as it allows determination of the average amount of the „COPD GENE“ polypeptide associated with a single cell by correlating the amount of „COPD GENE“ polypeptide in a cell-free extract produced from a predetermined number of cells.

10 In one embodiment, the present invention also provides a method wherein polynucleotide probes are immobilized on a DNA chip in an organized array. Oligonucleotides can be bound to a solid support by a variety of processes, including lithography. For example a chip can hold up to 250,000 oligonucleotides (GeneChip, Affymetrix). These polynucleotide probes comprise a nucleotide sequence at least  
15 about 12 nucleotides in length, preferably at least about 15 nucleotides, more preferably at least about 25 nucleotides, and most preferably at least about 40 nucleotides, and up to all or nearly all of a sequence which is complementary to a portion of the coding sequence of a „COPD GENE“ polynucleotide sequence selected from at least one of the polynucleotides of SEQ ID NO. 1 to 28. The present  
20 invention provides significant advantages over the available tests for various chronic lung diseases, such as COPD, because it increases the reliability of the test by providing an array of polynucleotide markers on a single chip.

The method includes obtaining tissue or cells from airways such as alveolar  
25 macrophages, e.g. by bronchoalveolar lavage (BAL). The DNA or RNA is extracted, amplified, and analyzed with a DNA chip to determine the presence or absence of the „COPD GENE“ polynucleotide sequences. In one embodiment, the polynucleotide probes are spotted onto a substrate in a two-dimensional matrix or array. Samples of polynucleotides can be labeled and then hybridized to the probes. Double-stranded  
30 polynucleotides, comprising the labeled sample polynucleotides bound to probe

polynucleotides, can be detected once the unbound portion of the sample is washed away.

5 The probe polynucleotides can be spotted on substrates including glass, nitro-cellulose, etc. The probes can be bound to the substrate by either covalent bonds or by non-specific interactions, such as hydrophobic interactions. The sample polynucleotides can be labeled using radioactive labels, fluorophores, chromophores, etc. Techniques for constructing arrays and methods of using these arrays are described in EP No. 0 799 897; PCT No. WO 97/29212; PCT No. WO 97/27317; 10 EP No. 0 785 280; PCT No. WO 97/02357; U.S. Pat. No. 5,593,839; U.S. Pat. No. 5,578,832; EP No. 0 728 520; U.S. Pat. No. 5,599,695; EP No. 0 721 016; U.S. Pat. No. 5,556,752; PCT No. WO 95/22058; and U.S. Pat. No. 5,631,734. Further, arrays can be used to examine differential expression of genes and can be used to determine gene function. For example, arrays of the instant polynucleotide sequences can be 15 used to determine if any of the polynucleotide sequences are differentially expressed between normal cells and cells in chronic lung disease, for example. High expression of a particular message in a cell from chronically diseased lung, which is not observed in a corresponding normal cell, can indicate a protein specific for chronic lung disease such as COPD.

20 In yet another embodiment, the invention contemplates using a panel of antibodies which are generated against the "COPD GENE" polypeptides of this invention, which polypeptides comprise a polynucleotide selected from SEQ ID NO. 29 to 56 or fragments thereof or are encoded by a polynucleotide which comprises a sequence 25 selected from at least one of the polynucleotides of SEQ ID NO. 1 to 28. Such a panel of antibodies may be used as a reliable diagnostic probe for chronic lung disease and COPD in particular. The assay of the present invention comprises contacting a biopsy sample containing cells, e.g., macrophages obtained by bronchoalveolar lavage, with a panel of antibodies to one or more of the encoded 30 products to determine the presence or absence of the "COPD GENE" polypeptides.

The diagnostic methods of the subject invention may also be employed as follow-up to treatment, e.g., quantification of the level of „COPD GENE“ polypeptide may be indicative of the effectiveness of current or previously employed chronic lung disease therapies as well as the effect of these therapies upon patient prognosis.

5

Accordingly, the present invention makes available diagnostic assays and reagents for detecting gain and/or loss of „COPD GENE“ polypeptides from a cell in order to aid in the diagnosis and phenotyping of chronic lung disorders, and COPD in particular.

10

The diagnostic assays described above can be adapted to be used as prognostic assays, as well. Such an application takes advantage of the sensitivity of the assays of the invention to events which take place at characteristic stages in the progression of chronic lung disease. For example, a given „COPD GENE“ may be up- or down-regulated at a very early stage, perhaps before the lung is irreversibly damaged, while another „COPD GENE“ may be characteristically up or down regulated only at a much later stage. Such a method could involve the steps of contacting the mRNA of a test cell with a polynucleotide probe derived from a given „COPD GENE“ polynucleotide which is expressed at different characteristic levels in chronically diseased lung tissue cells and COPD tissue cells in particular at different stages of progression, and determining the approximate amount of hybridization of the probe to the mRNA of the cell, such amount being an indication of the level of expression of the gene in the cell, and thus an indication of the stage of progression of chronic lung disease and COPD; alternatively, the assay can be carried out with an antibody specific for the gene product of the given „COPD GENE“ polynucleotide, contacted with the proteins of the test cell. The methods of the invention can also be used to follow the clinical course of chronic lung disease. For example, the assay of the invention can be applied to a tissue sample from a patient (e.g. macrophages from BAL); following treatment of the patient for chronic lung disease, another tissue sample is taken and the test repeated. Successful treatment will result in removal of all cells which demonstrate differential expression characteristic of the chronically

30

diseased lung tissue cells and COPD tissue cells in particular, or a substantial increase in expression of the gene in those cells, perhaps approaching or even surpassing normal levels.

5 In yet another embodiment, the invention provides methods for determining whether a subject is at risk for developing a disease, such as a predisposition to develop chronic lung disease, for example COPD, associated with an aberrant activity of at least one of the polypeptides of SEQ ID NO. 29 to 56, wherein the aberrant activity of the polypeptide is characterized by detecting the presence or absence of a genetic  
10 lesion characterized by at least one of (i) an alteration affecting the integrity of a gene encoding a „COPD GENE“ polypeptides, or (ii) the mis-expression of the encoding polynucleotide. To illustrate, such genetic lesions can be detected by ascertaining the existence of at least one of (i) a deletion of one or more nucleotides from the polynucleotide sequence, (ii) an addition of one or more nucleotides to the  
15 polynucleotide sequence, (iii) a substitution of one or more nucleotides of the polynucleotide sequence, (iv) a gross chromosomal rearrangement of the polynucleotide sequence, (v) a gross alteration in the level of a messenger RNA transcript of the polynucleotide sequence, (vi) aberrant modification of the polynucleotide sequence, such as of the methylation pattern of the genomic DNA, (vii) the presence  
20 of a non-wild type splicing pattern of a messenger RNA transcript of the gene, (viii) a non-wild type level of the „COPD GENE“ polypeptide, (ix) allelic loss of the gene, and/or (x) inappropriate post-translational modification of the „COPD GENE“ polypeptide

25 The present invention provides assay techniques for detecting lesions in the encoding polynucleotide sequence. These methods include, but are not limited to, methods involving sequence analysis, Southern blot hybridization, restriction enzyme site mapping, and methods involving detection of absence of nucleotide pairing between the polynucleotide to be analyzed and a probe.

30



Specific diseases or disorders, e.g., genetic diseases or disorders, are associated with specific allelic variants of polymorphic regions of certain genes, which do not necessarily encode a mutated protein. Thus, the presence of a specific allelic variant of a polymorphic region of a gene in a subject can render the subject susceptible to developing a specific disease or disorder. Polymorphic regions in genes, can be identified, by determining the nucleotide sequence of genes in populations of individuals. If a polymorphic region is identified, then the link with a specific disease can be determined by studying specific populations of individuals, e.g, individuals which developed a specific disease, such as chronic lung disease like COPD. A polymorphic region can be located in any region of a gene, e.g., exons, in coding or non coding regions of exons, introns, and promoter region.

In an exemplary embodiment, there is provided a polynucleotide composition comprising a polynucleotide probe including a region of nucleotide sequence which is capable of hybridizing to a sense or antisense sequence of a gene or naturally occurring mutants thereof, or 5' or 3' flanking sequences or intronic sequences naturally associated with the subject genes or naturally occurring mutants thereof. The polynucleotide of a cell is rendered accessible for hybridization, the probe is contacted with the polynucleotide of the sample, and the hybridization of the probe to the sample polynucleotide is detected. Such techniques can be used to detect lesions or allelic variants at either the genomic or mRNA level, including deletions, substitutions, etc., as well as to determine mRNA transcript levels.

A preferred detection method is allele specific hybridization using probes overlapping the mutation or polymorphic site and having about 5, 10, 20, 25, or 30 nucleotides around the mutation or polymorphic region. In a preferred embodiment of the invention, several probes capable of hybridizing specifically to allelic variants are attached to a solid phase support, e.g., a "chip". Mutation detection analysis using these chips comprising oligonucleotides, also termed "DNA probe arrays" is described e.g., in (34). In one embodiment, a chip comprises all the allelic variants of at least one polymorphic region of a gene. The solid phase support is then contacted

with a test polynucleotide and hybridization to the specific probes is detected. Accordingly, the identity of numerous allelic variants of one or more genes can be identified in a simple hybridization experiment.

5 In certain embodiments, detection of the lesion comprises utilizing the probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligase chain reaction (LCR) (see, e.g. Ref. 35 and 36), the latter of which can be particularly  
10 useful for detecting point mutations in the gene (see Ref. 37). In a merely illustrative embodiment, the method includes the steps of (i) collecting a sample of cells from a patient, (ii) isolating polynucleotide (e.g., genomic, mRNA or both) from the cells of the sample, (iii) contacting the polynucleotide sample with one or more primers which specifically hybridize to a polynucleotide sequence under conditions such that hybridization and amplification of the polynucleotide (if present) occurs, and (iv)  
15 detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

20 Alternative amplification methods include: self sustained sequence (38), transcriptional amplification system (39), Q-Beta Replicase (40), or any other polynucleotide amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are  
25 especially useful for the detection of polynucleotide molecules if such molecules are present in very low numbers.

In a preferred embodiment of the subject assay, mutations in, or allelic variants, of a gene from a sample cell are identified by alterations in restriction enzyme cleavage  
30 patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are

determined by gel electrophoresis. Moreover; the use of sequence specific ribozymes (see, for example, U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

## 5     Antibodies

Any type of antibody known in the art can be generated to bind specifically to an epitope of a „COPD GENE“ polypeptide. An antibody as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab)<sub>2</sub>, and  
10     Fv, which are capable of binding an epitope of a „COPD GENE“ polypeptide. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, e.g., at least 15, 25, or 50 amino acids.

15     An antibody which specifically binds to an epitope of a „COPD GENE“ polypeptide can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmunoassays, immunohistochemical assays, immuno-precipitations, or other immunochemical assays known in the art. Various immuno-  
20     assays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the immunogen.

Typically, an antibody which specifically binds to a „COPD GENE“ polypeptide  
25     provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to „COPD GENE“ polypeptides do not detect other proteins in immunochemical assays and can immunoprecipitate a „COPD  
30     GENE“ polypeptide from solution.

„COPD GENE“ polypeptides can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, a „COPD GENE“ polypeptide can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on  
5 the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (e.g., aluminum hydroxide), and surface active substances (e.g. lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially useful.  
10

Monoclonal antibodies which specifically bind to a „COPD GENE“ polypeptide can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not  
15 limited to, the hybridoma technique, the human B cell hybridoma technique, and the EBV hybridoma technique (41, 42, 43, 44). In addition, techniques developed for the production of chimeric antibodies, the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used (45, 46, 47). Monoclonal and other antibodies also  
20 can be humanized to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in treatment or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which  
25 differ from those in the human sequences by site directed mutagenesis of individual residues or by grafting of entire complementarity determining regions. Alternatively, humanized antibodies can be produced using recombinant methods, as described in GB2188638B. Antibodies which specifically bind to a „COPD GENE“ polypeptide can contain antigen binding sites which are either partially or fully humanized, as  
30 disclosed in U.S. Patent 5,565,332.

Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to „COPD GENE“ polypeptides. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries (48).

Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template (49). Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught, for example, in (50). Construction of bivalent, bispecific single-chain antibodies is taught in (51). A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology (52, 53).

Antibodies which specifically bind to „COPD GENE“ polypeptides also can be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature (54, 55). Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the antibodies described in WO 94/13804, also can be prepared.

Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which a „COPD GENE“ polypeptide is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

Antisense Oligonucleotides

Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 6 nucleotides in length, but can be at least 7, 8, 10, 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of „COPD GENE“ gene products in the cell.

Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, peptide nucleic acids (PNAs; described in US 5,714,331), locked nucleic acids (LNAs; described in WO 99/14226), or a combination of them. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamides, carboxymethyl esters, carbonates, and phosphate triesters (see Ref. 56, 57, 58). Modifications of „COPD GENE“ polynucleotide expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the „COPD GENE“. Oligonucleotides derived from the transcription initiation site, e.g., between positions 10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature (e.g., Ref. 59). An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a „COPD GENE“ polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a „COPD GENE“ polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent „COPD GENE“ nucleotides, can provide sufficient targeting specificity for „COPD GENE“ mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular „COPD GENE“ polynucleotide sequence.

Antisense oligonucleotides can be modified without affecting their ability to hybridize to a „COPD GENE“ polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5' substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art (see, e.g., 60, 61, 62).

### Ribozymes

Ribozymes are RNA molecules with catalytic activity (see, e.g., Ref. 63, 64, 65, 66).

Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art (e.g., Ref. 67). The mechanism of ribozyme action involves

sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammer-head motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences.

5

The transcribed sequence of a „COPD GENE“ can be used to generate ribozymes which will specifically bind to mRNA transcribed from a „COPD GENE“ genomic locus. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art (see Ref. 68). For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target (see, for example, Ref. 69).

15

Specific ribozyme cleavage sites within a „COPD GENE“ RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate „COPD GENE“ RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing

30



DNA construct into cells in which it is desired to decrease „COPD GENE“ expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A  
5 ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells.

As taught in (67), ribozymes can be engineered so that ribozyme expression will  
10 occur in response to factors which induce expression of a target gene. Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

#### 15 Screening Methods

The invention provides assays for screening test compounds which bind to or modulate the activity of a „COPD GENE“ polypeptide or a „COPD GENE“ polynucleotide. A test compound preferably binds to a „COPD GENE“ polypeptide  
20 or polynucleotide. More preferably, a test compound decreases „COPD GENE“ activity by at least about 10, preferably about 50, more preferably 90, or 100% relative to the absence of the test compound; or a test compound increases „COPD GENE“ activity by at least 0.3, preferably by 0.5, more preferably by 1, 2 or 5 or more fold relative to the absence of the test compound.

25

#### Test Compounds

Test compounds can be pharmacologic agents already known in the art or can be compounds previously unknown to have any pharmacological activity. The com-  
30 pounds can be naturally occurring or designed in the laboratory. They can be isolated from microorganisms, animals, or plants, and can be produced recombinantly, or

synthesized by chemical methods known in the art. If desired, test compounds can be obtained using any of the numerous combinatorial library methods known in the art, including but not limited to, biological libraries, spatially addressable parallel solid phase or solution phase libraries, synthetic library methods requiring deconvolution, the one-bead one-compound library method, and synthetic library methods using affinity chromatography selection. The biological library approach is limited to polypeptide libraries, while the other four approaches are applicable to polypeptide, non-peptide oligomer, or small molecule libraries of compounds (see Ref.70).

Methods for the synthesis of molecular libraries are well known in the art (see, for example, Ref. 71, 72, 73, 74, 75, 76, 77). Libraries of compounds can be presented in solution (see, e.g., Ref. 78), or on beads (79), chips (80), bacteria or spores (81), plasmids (82), or phage (83, 84, 85, 86 and 81).

#### High Throughput Screening

Test compounds can be screened for the ability to bind to „COPD GENE“ polypeptides or polynucleotides or to affect „COPD GENE“ activity or „COPD GENE“ gene expression using high throughput screening. Using high throughput screening, many discrete compounds can be tested in parallel so that large numbers of test compounds can be quickly screened. The most widely established techniques utilize 96-well, 384-well or 1536-well microtiter plates. The wells of the microtiter plates typically require assay volumes that range from 5 to 500 µl. In addition to the plates, many instruments, materials, pipettors, robotics, plate washers, and plate readers are commercially available to fit the microwell formats.

Alternatively, Free format assays, or assays that have no physical barrier between samples, can be used. For example, an assay using pigment cells (melanocytes) in a simple homogeneous assay for combinatorial peptide libraries is described by (87). The cells are placed under agarose in petri dishes, then beads that carry combinatorial compounds are placed on the surface of the agarose. The combinatorial compounds

are partially released the compounds from the beads. Active compounds can be visualized as dark pigment areas because, as the compounds diffuse locally into the gel matrix, the active compounds cause the cells to change colors.

5 Another example of a free format assay is described by Chelsky, "Strategies for Screening Combinatorial Libraries: Novel and Traditional Approaches," reported at the First Annual Conference of The Society for Biomolecular Screening in Philadelphia, Pa., Nov. 710, 1995. Chelsky placed a simple homogenous enzyme  
10 assay for carbonic anhydrase inside an agarose gel such that the enzyme in the gel would cause a color change throughout the gel. Thereafter, beads carrying combinatorial compounds via a photolinker were placed inside the gel and the compounds were partially released by UV light. Compounds that inhibited the enzyme were observed as local zones of inhibition having less color change.

15 Yet another example is described in (88). In this example, combinatorial libraries were screened for compounds that had cytotoxic effects on cancer cells growing in agar.

20 Another high throughput screening method is described in (89). In this method, test samples are placed in a porous matrix. One or more assay components are then placed within, on top of, or at the bottom of a matrix such as a gel, a plastic sheet, a filter, or other form of easily manipulated solid support. When samples are introduced to the porous matrix they diffuse sufficiently slowly, such that the assays can be performed without the test samples running together.

25

#### Binding Assays

For binding assays, the test compound is preferably a small molecule which binds to and occupies, for example, the ATP/GTP binding site of the enzyme or the active site  
30 of a „COPD GENE“ polypeptide, such that normal biological activity is prevented.

Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules.

5 In binding assays, either the test compound or a „COPD GENE“ polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to a „COPD GENE“ polypeptide can then be accomplished, for example, by direct counting of radioemission, by scintillation counting, or by determining conversion of an appropriate  
10 substrate to a detectable product.

Alternatively, binding of a test compound to a „COPD GENE“ polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a „COPD GENE“  
15 polypeptide. A microphysiometer (e.g., CytosensorJ) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and a „COPD GENE“ polypeptide (90).

20 Determining the ability of a test compound to bind to a „COPD GENE“ polypeptide also can be accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) (91 and 92). BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore™).  
25 Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

In yet another aspect of the invention, a „COPD GENE“ polypeptide can be used as a "bait protein" in a two-hybrid assay or three-hybrid assay (see, e.g., Ref. 93, 94, 95,  
30 96, 97 and 98), to identify other proteins which bind to or interact with the „COPD GENE“ polypeptide and modulate its activity.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide  
5 encoding a „COPD GENE“ polypeptide can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (e.g., GAL4). In the other construct a DNA sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact in vivo to  
10 form an protein- dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can  
15 be isolated and used to obtain the DNA sequence encoding the protein which interacts with the „COPD GENE“ polypeptide.

It may be desirable to immobilize either a „COPD GENE“ polypeptide (or polynucleotide) or the test compound to facilitate separation of bound from unbound  
20 forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either a „COPD GENE“ polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex,  
25 polystyrene, or glass beads). Any method known in the art can be used to attach a „COPD GENE“ polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid  
30 support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to a „COPD GENE“ polypeptide (or polynucleotide) can

be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

5 In one embodiment, a „COPD GENE“ polypeptide is a fusion protein comprising a domain that allows the „COPD GENE“ polypeptide to be bound to a solid support. For example, glutathione S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the nonadsorbed „COPD GENE“ polypeptide; the mixture is then  
10 incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

15 Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either a „COPD GENE“ polypeptide (or polynucleotide) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated „COPD GENE“  
20 polypeptides (or polynucleotides) or test compounds can be prepared from biotin NHS (N-hydroxysuccinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies which specifically bind to a „COPD GENE“ polypeptide, polynucleotide, or a test com-  
25 pound, but which do not interfere with a desired binding site, such as the ATP/GTP binding site or the active site of the „COPD GENE“ polypeptide, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

30 Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using anti-

bodies which specifically bind to a „COPD GENE“ polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of a „COPD GENE“ polypeptide, and SDS gel electrophoresis under non-reducing conditions.

- 5 Screening for test compounds which bind to a „COPD GENE“ polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a „COPD GENE“ polypeptide or polynucleotide can be used in a cell-based assay system. A „COPD GENE“ polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test  
10 compound to a „COPD GENE“ polypeptide or polynucleotide is determined as described above.

#### Enzyme Assays

- 15 As given in Table 3 some of the „COPD GENE“ polypeptides have enzymatic activity. Test compounds can be tested for the ability to increase or decrease the activity of a „COPD GENE“ polypeptide that has enzymatic activity. Enzymatic activity of a „COPD GENE“ polypeptide can be measured, for these polypeptides, as known in the art. Enzyme assays can be carried out after contacting either a purified  
20 „COPD GENE“ polypeptide, a cell membrane preparation, or an intact cell with a test compound.

#### Modulation of Gene Expression

- 25 In another embodiment, test compounds which increase or decrease „COPD GENE“ gene expression are identified. A „COPD GENE“ polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of the „COPD GENE“ polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of  
30 expression of mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a modulator of expression based on this

comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

The level of „COPD GENE“ mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of a „COPD GENE“ polynucleotide can be determined, for example, using a variety of techniques known in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined in vivo, in a cell culture, or in an in vitro translation system by detecting incorporation of labeled amino acids into a „COPD GENE“ polypeptide.

Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses a „COPD GENE“ polynucleotide can be used in a cell-based assay system. A „COPD GENE“ polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Either a primary culture or an established cell line, such as CHO or human embryonic kidney 293 cells, can be used.

## Pharmaceutical Compositions

The invention also provides pharmaceutical compositions which can be administered to a patient to achieve a therapeutic effect. Pharmaceutical compositions of the invention can comprise, for example, a „COPD GENE“ polypeptide, „COPD GENE“ polynucleotide, ribozymes or antisense oligonucleotides, antibodies which specifically bind to a „COPD GENE“ polypeptide, or mimetics, agonists, antagonists,



partial agonists, inverse agonists, activators, co-activators or inhibitors of a „COPD GENE“ polypeptide activity. The compositions can be administered alone or in combination with at least one other agent, such as stabilizing compound, which can be administered in any sterile, biocompatible pharmaceutical carrier, including, but  
5 not limited to, saline, buffered saline, dextrose, and water. The compositions can be administered to a patient alone, or in combination with other agents, drugs or hormones.

In addition to the active ingredients, these pharmaceutical compositions can contain  
10 suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Pharmaceutical compositions of the invention can be administered by any number of routes including, but not limited to, oral, inhalation, intravenous, intramuscular, intraarterial, intramedullary, intrathecal, intraventricular,  
15 transdermal, subcutaneous, intraperitoneal, intranasal, parenteral, topical, sublingual, or rectal means. Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels,  
20 syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to  
25 obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethylcellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or  
30 solubilizing agents can be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores can be used in conjunction with suitable coatings, such as concentrated sugar solutions, which also can contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds can be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration can be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions can contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers also can be used for delivery. Optionally, the suspension also can contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention can be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. The pharmaceutical composition can be provided as a salt and  
5 can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation can be a lyophilized powder which can contain any or all of the following: 150 mM histidine, 0.1%2% sucrose, and 27% mannitol, at a pH range  
10 of 4.5 to 5.5, that is combined with buffer prior to use.

Further details on techniques for formulation and administration can be found in the latest edition of REMINGTON'S PHARMACEUTICAL SCIENCES (Maack Publishing Co., Easton, Pa.). After pharmaceutical compositions have been prepared, they can be  
15 placed in an appropriate container and labeled for treatment of an indicated condition. Such labeling would include amount, frequency, and method of administration.

#### Therapeutic Indications and Methods

20 Therapies for treatment of COPD primarily relied upon medications to open the airways and decrease inflammation, oxygen supplementation and pulmonary rehabilitation. The advent of genomics-driven molecular target identification has opened up the possibility of identifying new lung disease-specific targets for  
25 therapeutic intervention that will provide safer, more effective treatments for chronic lung disease patients. Thus, newly discovered COPD-associated genes and their products can be tested for their role(s) in chronic lung disease and used as tools to discover and develop innovative therapies.

30 Genes playing important roles in any of the physiological processes outlined above can be characterized as chronic lung disease targets. Genes or gene fragments

identified through genomics can readily be expressed in one or more heterologous expression systems to produce functional recombinant proteins. These proteins are characterized *in vitro* for their biochemical properties and then used as tools in high-throughput molecular screening programs to identify chemical modulators of their biochemical activities. Modulators of target protein activity can be identified in this manner and subsequently tested in cellular and *in vivo* disease models for therapeutic activity. Optimization of lead compounds with iterative testing in biological models and detailed pharmacokinetic and toxicological analyses form the basis for drug development and subsequent testing in humans.

„COPD GENES“ can be utilized as therapeutic targets for chronic lung disease and COPD in particular. Inhibition of the activity of the upregulated „COPD GENE“ polynucleotides or polypeptides leads to a decreased activity of the „COPD GENE“ polynucleotides or polypeptides, overexpressed in COPD. Activation of the downregulated „COPD GENE“ polynucleotides or polypeptides leads to an enhanced activity of the „COPD GENE“ polynucleotides or polypeptides, expressed at a lower level in COPD patients.

This invention further pertains to the use of novel agents identified by the screening assays described above. Accordingly, it is within the scope of this invention to use a test compound identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a modulating agent, an antisense nucleic acid molecule, a specific antibody, ribozyme, or a human „COPD GENE“ polypeptide binding molecule) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above described screening assays for treatments as described herein.

A reagent which affects human „COPD GENE“ activity can be administered to a human cell, either in vitro or in vivo, to reduce or increase human „COPD GENE“ activity. The reagent preferably binds to an expression product of a human „COPD GENE“ gene. If the expression product is a protein, the reagent is preferably an antibody. For treatment of human cells *ex vivo*, an antibody can be added to a preparation of stem cells which have been removed from the body. The cells can then be replaced in the same or another human body, with or without clonal propagation, as is known in the art.

In one embodiment, the reagent is delivered using a liposome. Preferably, the liposome is stable in the animal into which it has been administered for at least about 30 minutes, more preferably for at least about 1 hour, and even more preferably for at least about 24 hours. A liposome comprises a lipid composition that is capable of targeting a reagent, particularly a polynucleotide, to a particular site in an animal, such as a human. Preferably, the lipid composition of the liposome is capable of targeting to a specific organ of an animal, such as the lung, liver, spleen, heart brain, lymph nodes, and skin.

A liposome useful in the present invention comprises a lipid composition that is capable of fusing with the plasma membrane of the targeted cell to deliver its contents to the cell. Preferably, the transfection efficiency of a liposome is about 0.5 µg of DNA per 16 nmole of liposome delivered to about  $10^6$  cells, more preferably about 1.0 µg of DNA per 16 nmole of liposome delivered to about  $10^6$  cells, and even more preferably about 2.0 µg of DNA per 16 nmol of liposome delivered to about  $10^6$  cells. Preferably, a liposome is between about 100 and 500 nm, more preferably between about 150 and 450 nm, and even more preferably between about 200 and 400 nm in diameter.

Suitable liposomes for use in the present invention include those liposomes standardly used in, for example, gene delivery methods known to those of skill in the art. More preferred liposomes include liposomes having a polycationic lipid

composition and/or liposomes having a cholesterol backbone conjugated to polyethylene glycol. Optionally, a liposome comprises a compound capable of targeting the liposome to a particular cell type, such as a cell-specific ligand exposed on the outer surface of the liposome.

5

Complexing a liposome with a reagent such as an antisense oligonucleotide or ribozyme can be achieved using methods which are standard in the art (see, for example, Ref. 99). Preferably, from about 0.1  $\mu$ g to about 10  $\mu$ g of polynucleotide is combined with about 8 nmol of liposomes, more preferably from about 0.5  $\mu$ g to about 5  $\mu$ g of polynucleotides are combined with about 8 nmol liposomes, and even more preferably about 1.0  $\mu$ g of polynucleotides is combined with about 8 nmol liposomes.

10

15

In another embodiment, antibodies can be delivered to specific tissues in vivo using receptor-mediated targeted delivery. Receptor-mediated DNA delivery techniques are taught in, for example, in (Ref. 100, 101, 102, 103, 104, 105).

#### Determination of a Therapeutically Effective Dose

20

The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases human „COPD GENE“ activity relative to the human „COPD GENE“ activity which occurs in the absence of the therapeutically effective dose.

25

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

30

Therapeutic efficacy and toxicity, e.g.,  $ED_{50}$  (the dose therapeutically effective in 50% of the population) and  $LD_{50}$  (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it  
5 can be expressed as the ratio,  $LD_{50}/ED_{50}$ .

Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is  
10 preferably within a range of circulating concentrations that include the  $ED_{50}$  with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to  
15 the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response  
20 to treatment. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts can vary from 0.1 to 100,000 micrograms, up to a total dose  
25 of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions,  
30 locations, etc.

If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either ex vivo or in vivo using well-established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, 5 protoplast fusion, viral infection, electroporation, a gene gun, and DEAE- or calcium phosphate-mediated transfection.

Effective in vivo dosages of an antibody are in the range of about 5 µg to about 10 50 µg/kg, about 50 µg to about 5 mg/kg, about 100 µg to about 500 µg/kg of patient body weight, and about 200 to about 250 µg/kg of patient body weight. For administration of polynucleotides encoding single-chain antibodies, effective in vivo dosages are in the range of about 100 ng to about 200 ng, 500 ng to about 50 mg, about 1 µg to about 2 mg, about 5 µg to about 500 µg, and about 20 µg to about 15 100 µg of DNA.

If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described 20 above.

Preferably, a reagent reduces expression of a „COPD GENE“ polynucleotide or the activity of a „COPD GENE“ polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of a 25 „COPD GENE“ polynucleotide or the activity of a „COPD GENE“ polypeptide can be assessed using methods well known in the art, such as hybridization of nucleotide probes to „COPD GENE“-specific mRNA, quantitative RT-PCR, immunologic detection of a „COPD GENE“ polypeptide, or measurement of the „COPD GENE“ 30 polypeptide activity.



In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination treatment can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

Any of the therapeutic methods described above can be applied to any subject in need of such treatment, including, livestock, for example, cows, pigs, sheep, goats, horses and domestic animals, for example birds, dogs, cats, rabbits and other animals such as monkeys, and most preferably, humans.

All patents and patent applications cited in this disclosure are expressly incorporated herein by reference. The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples which are provided for purposes of illustration only and are not intended to limit the scope of the invention.

20

EXAMPLESEXAMPLE 15     Material and methods1.a) Probe selection for the differential gene expression analysis

10     Three patients with clear clinical evidence of COPD and a cigarette consumption equivalent of less than 20 pack years compared to five patients with a cigarette consumption equivalent of more than 40 pack years and no evidence of COPD provide the material for the study. These two groups of smokers represent the end-points of the COPD disease spectrum.

15     The clinical study is based on a patient-stratification scheme which is exactly followed.

In detail, the two groups of patients are classified as follows:

20     Group 1:     smokers without COPD

- cigarette smoking equivalent > 40 pack years (1py= 1 pack of cigarettes per day for one year)
- sex: male
- 25     - no airway obstruction (FEV1> 70%)
- normal diffusing capacity (> 85% of normal)
- thorax computed tomography (CT): no signs of emphysema

30     Group 2:     smokers with severe COPD plus emphysema

- cigarette smoking equivalent < 20 pack years

- sex: male
- airway obstruction (FEV1 < 70%) and no reversibility (< 12%)
- reduced diffusion capacity (< 60%)
- thorax CT: clear emphysema

5

The following groups are excluded from the study:

- patients with alpha1 anti-trypsin deficiency
- patients with serious diseases such as diabetes, cardiovascular diseases, liver or kidney diseases
- patients who have been treated with systemic or oral steroids, antibiotics or NSAIDs in the last two months.

10

15

The two different populations of smokers underwent bronchoalveolar lavages (BALs) in two lung segments of the same side. The BAL fluid was characterized and the presence of > 90% macrophages and at least  $1.5 \times 10^7$  cells were criteria for including the samples into the study. Macrophages were isolated.

20 1.b) Differential DNA expression profiling

25

30

Expression profiling was carried out using the Affymetrix Array Technology. Total RNA was prepared from these macrophages according to a standard protocol as described in: Expression Analysis Technical Manual, Affymetrix Inc., CA, USA, cDNA was generated, labeled and hybridized to the Affymetrix Hg-U95A DNA array containing oligonucleotides of 12626 human annotated genes (according to the manufacturer's instructions). The data have been analyzed by bioinformatics tools, using the Affymetrix software and additional filter criteria. The bioinformatics analysis is described in detail below.

28 Genes were identified to be at least 1.5 fold, differentially expressed in patients with COPD in comparison to patients without COPD. 18 genes which are up-regulated (Table 1) and 10 genes which are down-regulated (Table 2) in COPD patients have been identified.

5

To confirm the results obtained by the array analysis with a second independent experimental approach, these 28 genes were analyzed by real-time quantitative PCR (TaqMan), using the PRISM 7700 Sequence Detection System of PE Applied Biosystems (Perkin Elmer, Foster City, CA, USA. Here a fluorogenic probe, consisting of an oligonucleotide labeled with both a fluorescent reporter dye and a quencher dye, is included in a typical PCR. Amplification of the probe-specific product causes cleavage of the probe, generating an increase in reporter fluorescence. To standardize the amount of sample RNA, GAPDH was selected as a reference, since it was not differentially regulated in IC versus IVA patients. Primers and probes were selected using the Primer Express software and localized mostly in the 3' region of the coding sequence or in the 3' untranslated region (see Tables 4 and 5 for primer- and probe- sequences). All primer pairs were checked for specificity by conventional PCR reactions. TaqMan validation experiments were performed showing that the efficiencies of the target and the control amplifications are approximately equal which is a prerequisite for the relative quantitation of gene expression by the comparative  $C_T$  method.

10  
15  
20

To further verify the disease relevance of a selection of the upregulated genes identified in Table 1, real-time PCR (Taqman) using the PRISM 7700 Sequence Detection System was carried out as described above but with  $\beta_2$ -microglobulin instead of GAPDH for standardization using mRNA prepared from BAL macrophages from a second independent donor set (COPD patients, and smoking matched controls) -see Table 6 for primers. Up regulated genes were confirmed for the second donor set (Table 6) in 3 to 6 donors and by greater than 2-fold increase in expression.

25  
30

1.c) Data analysis

According to Affymetrix' measurement technique a single gene expression measurement on one chip yields the *average difference* value and the *absolute call*. The *average difference* is a real number supposed to represent the expression value of that gene. The *absolute call* can take the values 'A' (absent), 'M' (marginal), or 'P' (present) and denotes the quality of a single hybridization. We used both the quantitative information given by the *average difference* and the qualitative information given by the *absolute call* to identify the genes which are differentially expressed in the COPD versus the normal population.

We calculated the differential expression E in the COPD versus the normal population as follows. Given D *average difference* values  $c_1, c_2, \dots, c_D$  in the COPD population and N *average difference* values  $m_1, m_2, \dots, m_N$  in the population of normals, we compute

$$E \equiv \exp\left(\frac{1}{D} \sum_{i=1}^D \ln(c_i) - \frac{1}{N} \sum_{i=1}^N \ln(m_i)\right).$$

20

If  $c_i < L$  or  $d_j < L$  for one or more values of i and j, we set these particular values equal to the positive number L. L represents the *average difference* level below which the measurements are not meaningful. Typically we set  $L=50$ .

25

Note that the particular computation of E allows for a correct comparison to TaqMan results as obtained by the ' $\Delta\Delta C_t$  - method'.

A gene is called *up-regulated* in COPD versus normal if  $E \geq 1.5$  and if the number of *absolute calls* equal to 'P' in the COPD population is greater than  $D/2$ .

5 A gene is called *down-regulated* in COPD versus normal if  $E \leq 1.5$  and if the number of *absolute calls* equal to 'P' in the normal population is greater than  $N/2$ .

10 Based on the *average difference* values  $c_i$  and  $d_j$  we also calculated p-values by means of Wilcoxon's rank test. In this way we were able to test for the significance of the expression value difference of the genes in the COPD versus the normal population.

15 The final list of differentially regulated genes consists of all *up-regulated* and all *down-regulated* genes in the COPD versus the normal population. Those genes on this list which are interesting for a pharmaceutical application were finally validated by TaqMan.

## 20 2. Physiological and biochemical significance of the results

20 Tables 1 to 3 show the summary of the genes, including the differential expression values of array analysis and TaqMan analysis, which show an excellent correlation of gene expression levels.

25 All 28 sequences were previously described but have not been previously recognized as being differentially expressed in COPD patients versus non-COPD patients.

30 Of these 28 genes, three genes (SEQ ID NO: 7, 8, 21) belong to the family of chemokines, which appear to be involved in a variety of pro-inflammatory diseases and this makes them and their receptors very attractive therapeutic

targets. Inflammation of the lung is a key feature of COPD. In COPD, interleukin-8 has already been identified as a major up-regulated chemokine. Three G-protein coupled receptors (SEQ ID NO: 11, 12, 16) have been shown to be differentially regulated. GPCRs are involved in a variety of signal transduction pathways and have proven to be among the most successful drug targets for many indications. They have been shown to be expressed in lung tissue but were not implicated in COPD. Therefore novel GPCRs will be good targets for therapeutic intervention of respiratory diseases.

Two proteases (SEQ ID NO: 10, 24) have been recognized as differentially expressed. Different classes of proteases have been identified that have the potential to contribute to lung matrix destruction. A selective inhibitor of PDE4 isoforms has demonstrated already therapeutic utility in clinical trials of subjects with COPD (106).

Transcription factors ( SEQ ID NO: 1, 3, 4, 5, 14), two signaling molecules (SEQ ID NO: 17, 20) as well as the two phosphatases (SEQ ID NO: 6, 25) were identified, all of them directly involved in signal transduction processes. Additionally two transporter (SEQ ID NO: 15, 26) and a transferase (SEQ ID NO: 18) have been identified.

## EXAMPLE 2

### Identification of test compounds that bind to „COPD GENE“ polypeptides

A purified „COPD GENE“ polypeptide comprising a glutathione-S-transferase protein and absorbed onto glutathione-derivatized wells of 96-well microtiter plates is contacted with test compounds from a small molecule library at pH 7.0 in a physiological buffer solution. The test compounds comprise a fluorescent tag. The samples are incubated for 5 minutes to one hour. Control samples are incubated in the absence of a test compound.

The buffer solution containing the test compounds is washed from the wells. Binding of a test compound to a „COPD GENE“ polypeptide is detected by fluorescence measurements of the contents of the wells. A test compound which increases the fluorescence in a well by at least 15% relative to fluorescence of a well in which a test compound is not incubated is identified as a compound which binds to a „COPD GENE“ polypeptide.

### EXAMPLE 3

10

#### Identification of a test compound which decreases „COPD GENE“ gene expression

A test compound is administered to a culture of human cells transfected with a „COPD GENE“ expression construct and incubated at 37°C for 10 to 45 minutes. A culture of the same type of cells which have not been transfected is incubated for the same time without the test compound to provide a negative control.

RNA is isolated from the two cultures as described in (107). Northern blots are prepared using 20 to 30 µg total RNA and hybridized with a <sup>32</sup>P-labeled „COPD GENE“ -specific probe at 65°C in Express-hyb (CLONTECH). The probe comprises at least 11 contiguous nucleotides selected from the complement of SEQ ID NO. 19. A test compound which decreases the „COPD GENE“ -specific signal relative to the signal obtained in the absence of the test compound is identified as an inhibitor of „COPD GENE“ gene expression.

25

### EXAMPLE 4

#### Treatment of COPD in an animal model

Guinea pigs are exposed on a single occasion to tobacco smoke for 50 minutes. Animals are sacrificed between 10 minutes and 24 hour following the end of the

30



exposure and their lungs placed in RNAlater™. The lung tissue is homogenized and total RNA is extracted using a Qiagen RNeasy™ Maxi kit. Molecular Probes RiboGreen™ RNA quantitation method is used to quantify the amount of RNA in each sample. Total RNA is reverse transcribed and the resultant cDNA was used in a  
5 real-time polymerase chain reaction (PCR). The cDNA is added to a solution containing the sense and anti-sense primers and the 6-carboxy-tetramethyl-rhodamine labeled probe of the "COPD GENE". Cyclophilin is used as the housekeeping gene. The expression of the "COPD GENE" is measured using the TaqMan real-time PCR system that generates an amplification curve for each sample. From this curve a  
10 threshold cycle value is calculated: the fractional cycle number at which the amount of amplified target reaches a fixed threshold. A sample containing many copies of the "COPD GENE" will reach this threshold earlier than a sample containing fewer copies. The threshold is set at 0.2 and the threshold cycle  $C_T$  is calculated from the amplification curve. The  $C_T$  value for the "COPD GENE" is normalized using the  
15  $C_T$  value for the housekeeping gene.

Expression of the "COPD GENE" is increased by at least 1,5-fold between 10 minutes and 3 hours post tobacco smoke exposure compared to air exposed control  
20 animals.

Test compounds are evaluated as follows. Animals are pre-treated with a test compound between 5 minutes and 1 hour prior to the tobacco smoke exposure and they are then sacrificed up to 3 hours after the tobacco smoke exposure has been completed. Control animals are pre-treated with the vehicle of the test compound via  
25 the route of administration chosen for the test compound. A test compound that reduces the tobacco smoke induced upregulation of the "COPD GENE" relative to the expression seen in vehicle treated tobacco smoke exposed animals is identified as an inhibitor of "COPD GENE" expression.

References

1. Barnes, P.J. Mechanisms in COPD. Differences from asthma. Chest 2000, 117:10S14S.
- 5 2. Bonner et al., J. Mol. Biol. 81, 123 (1973).
3. Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2d ed., 1989.
4. Bolton and McCarthy, Proc. Natl. Acad. Sci. U.S.A. 48, 1390 (1962).
5. Sarkar, PCR Methods Applic. 2, 318322, 1993.
- 10 6. Triglia et al., Nucleic Acids Res. 16, 8186, 1988.
7. Lagerstrom et al., PCR Methods Applic. 1, 111119, 1991.
8. Parker et al., Nucleic Acids Res. 19, 30553060, 1991.
9. Ausubel et al., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, N.Y., 1989.
- 15 10. Grant et al., Methods Enzymol. 153, 516544, 1987.
11. Takamatsu, EMBO J. 6, 307311, 1987.
12. Coruzzi et al., EMBO J. 3, 16711680, 1984.
13. Broglie et al., Science 224, 838843, 1984.
14. Winter et al., Results Probl. Cell Differ. 17, 85105, 1991.
- 20 15. MCGRAW HILL YEARBOOK OF SCIENCE AND TECHNOLOGY, McGraw Hill, New York, N.Y., pp. 191196, 1992.
16. Engelhard et al., Proc. Nat. Acad. Sci. 91, 32243227, 1994.
17. Logan & Shenk, Proc. Natl. Acad. Sci. 81, 36553659, 1984.
18. Scharf et al., Results Probl. Cell Differ. 20, 125162, 1994.
- 25 19. Wigler et al., Cell 11, 22332, 1977.
20. Lowy et al., Cell 22, 81723, 1980.
21. Wigler et al., Proc. Natl. Acad. Sci. 77, 356770, 1980.
22. Colbere-Garapin et al., J. Mol. Biol. 150, 114, 1981.
23. Hartman & Mulligan, Proc. Natl. Acad. Sci. 85, 804751, 1988.
- 30 24. Rhodes et al., Methods Mol. Biol. 55, 121131, 1995.

25. Hampton et al., SEROLOGICAL METHODS: A LABORATORY MANUAL, APS Press, St. Paul, Minn., 1990.
26. Maddox et al., J. Exp. Med. 158, 12111216, 1983.
27. Porath et al., Prot. Exp. Purif. 3, 263281, 1992.
- 5 28. Kroll et al., DNA Cell Biol. 12, 441453, 1993.
29. Caruthers et al., Nucl. Acids Res. Symp. Ser. 215223, 1980.
30. Horn et al. Nucl. Acids Res. Symp. Ser. 225232, 1980.
31. Merrifield, J. Am. Chem. Soc. 85, 21492154, 1963.
32. Roberge et al., Science 269, 202204, 1995.
- 10 33. Creighton, PROTEINS: STRUCTURES AND MOLECULAR PRINCIPLES, WH and Co., New York, N.Y., 1983.
34. Cronin et al. (1996) Human Mutation 7:244.
35. Landegran *et al.* (1988) *Science* 241:1077-1080.
36. Nakazawa *et al.* (1994) PNAS 91:360-364.
- 15 37. Abravaya et al. (1995) *Nuc Acid Res* 23:675-682.
38. Guatelli, J.C. *et al.*, 1990, Proc. Natl. Acad. Sci. USA 87:1874-1878.
39. Kwoh, D.Y. *et al.*, 1989, Proc. Natl. Acad. Sci. USA 86:1173 1177.
40. Lizardi, P.M. *et al.*, 1988, Bio/Technology 6:1197.
41. Kohler et al., Nature 256, 495497, 1985.
- 20 42. Kozbor et al., J. Immunol. Methods 81, 3142, 1985.
43. Cote et al., Proc. Natl. Acad. Sci. 80, 20262030, 1983.
44. Cole et al., Mol. Cell Biol. 62, 109120, 1984.
45. Morrison et al., Proc. Natl. Acad. Sci. 81, 68516855, 1984.
46. Neuberger et al., Nature 312, 604608, 1984.
- 25 47. Takeda et al., Nature 314, 452454, 1985.
48. Burton, Proc. Natl. Acad. Sci. 88, 1112023, 1991.
49. Thirion et al., 1996, Eur. J. Cancer Prev. 5, 507-11.
50. Coloma & Morrison, 1997, Nat. Biotechnol. 15, 159-63.
51. Mallender & Voss, 1994, J. Biol. Chem. 269, 199-206.
- 30 52. Verhaar et al., 1995, Int. J. Cancer 61, 497-501.
53. Nicholls et al., 1993, J. Immunol. Meth. 165, 81-91.

54. Orlandi et al., Proc. Natl. Acad. Sci. 86, 3833 3837, 1989.
55. Winter et al., Nature 349, 293299, 1991.
56. Brown, Meth. Mol. Biol. 20, 18, 1994.
57. Sonveaux, Meth. Mol. Biol. 26, 1-72, 1994.
- 5 58. Uhlmann et al., Chem. Rev. 90, 543583, 1990.
59. Gee et al., in Huber & Carr, MOLECULAR AND IMMUNOLOGIC APPROACHES, Publishing Co., Mt. Kisco, N.Y., 1994.
60. Agrawal et al., Trends Biotechnol. 10, 152158, 1992.
61. Uhlmann et al., Chem. Rev. 90, 543584, 1990.
- 10 62. Uhlmann et al., Tetrahedron. Lett. 215, 35393542, 1987.
63. Cech, Science 236, 15321539; 1987.
64. Cech, Ann. Rev. Biochem. 59, 543568; 1990.
65. Cech, Curr. Opin. Struct. Biol. 2, 605609; 1992.
66. Couture & Stinchcomb, Trends Genet. 12, 510515, 1996.
- 15 67. Haseloff et al., U.S. Patent 5,641,673.
68. Haseloff et al. Nature 334, 585591, 1988.
69. Gerlach et al., EP 321,201.
70. Lam, Anticancer Drug Des. 12, 145, 1997.
71. DeWitt et al., Proc. Natl. Acad. Sci. U.S.A. 90, 6909, 1993.
- 20 72. Erb et al. Proc. Natl. Acad. Sci. U.S.A. 91, 11422, 1994.
73. Zuckermann et al., J. Med. Chem. 37, 2678, 1994.
74. Cho et al., Science 261, 1303, 1993.
75. Carell et al., Angew. Chem. Int. Ed. Engl. 33, 2059, 1994.
76. Carell et al., Angew. Chem. Int. Ed. Engl. 33, 2061
- 25 77. Gallop et al., J. Med. Chem. 37, 1233, 1994.
78. Houghten, BioTechniques 13, 412421, 1992.
79. Lam, Nature 354, 8284, 1991.
80. Fodor, Nature 364, 555556, 1993.
81. Ladner, U.S. Patent 5,223,409.
- 30 82. Cull et al., Proc. Natl. Acad. Sci. U.S.A. 89, 18651869, 1992.
83. Scott & Smith, Science 249, 386390, 1990.

84. Devlin, Science 249, 404406, 1990.
85. Cwirla et al., Proc. Natl. Acad. Sci. 97, 63786382, 1990.
86. Felici, J. Mol. Biol. 222, 301310, 1991.
87. Jayawickreme et al., Proc. Natl. Acad. Sci. U.S.A. 19, 161418, 1994.
- 5 88. Salmon et al., Molecular Diversity 2, 5763, 1996.
89. Beutel et al., U.S. Patent 5,976,813.
90. McConnell et al., Science 257, 19061912, 1992.
91. Sjolander & Urbaniczky, Anal. Chem. 63, 23382345, 1991.
92. Szabo et al., Curr. Opin. Struct. Biol. 5, 699705, 1995.
- 10 93. U.S. Patent 5,283,317.
94. Zervos et al., Cell 72, 223232, 1993.
95. Madura et al., J. Biol. Chem. 268, 1204612054, 1993.
96. Bartel et al., BioTechniques 14, 920924, 1993.
97. Iwabuchi et al., Oncogene 8, 16931696, 1993.
- 15 98. Brent W094/10300.
99. U.S. Patent 5,705,151.
100. Findeis et al. Trends in Biotechnol. 11, 202-05, 1993.
101. Chiou et al., GENE THERAPEUTICS: METHODS AND APPLICATIONS OF DIRECT GENE TRANSFER (J.A. Wolff, ed.), 1994.
- 20 102. Wu & Wu, J. Biol. Chem. 263, 621-24, 1988.
103. Wu et al., J. Biol. Chem. 269, 542-46, 1994.
104. Zenke et al., Proc. Natl. Acad. Sci. U.S.A. 87, 3655-59, 1990.
105. Wu et al., J. Biol. Chem. 266, 338-42, 1991.
106. Barnette & Underwood, Curr Opin Pulm Med. 6,164-9, 2000.
- 25 107. Chirgwin et al., Biochem. 18, 5294-99, 1979.
108. Walter et al., J. Exp. Med. 193, 339-52, 2001
109. Cowburn et al., J. Clin. Invest. 101, 834-46, 1998
110. Pilette et al., Am. J. Crit. Care Med. 163, 185-94, 2001
111. Needleman SB, Wunsch, J Mol Biol. 48, 443-53, 1970
- 30 112. Henikoff S, Henikoff JG, Proc. Natl. Acad. Sci. USA 89,10915-10919, 1992

## SEQUENCE LISTING

<110> Bayer AG

<120> GENES AND PROTEINS FOR PREVENTION, PREDICTION, DIAGNOSIS,  
PROGNOSIS AND TREATMENT OF CHRONIC LUNG DISEASE

<130> LeA 35183 Foreign Countries

<150> GB 0113266.1

<151> 2001-05-31

<160> 182

<170> PatentIn version 3.1

<210> 1

<211> 1345

<212> DNA

<213> Homo sapiens

<400> 1  
aaaacagccg gggctccagc gggagaacga taatgcaaag tgctatgttc ttggctgttc 60  
aacacgactg cagacccatg gacaagagcg caggcagtgg ccacaagagc gaggagaagc 120  
gagaaaagat gaaacggacc cttttaaaag attggaagac ccgtttgagc tacttcttac 180  
aaaattcctc tactcctggg aagcccaaaa ccggcaaaaa aagcaaacag caagctttca 240  
tcaagccttc tcctgaggaa gcacagctgt ggtcagaagc atttgacgag ctgctagcca 300  
gcaaatatgg tcttgctgca ttcagggctt ttttaaagtc ggaattctgt gaagaaaata 360  
ttgaattctg gctggcctgt gaagacttca aaaaaaccaa atcaccccaa aagctgtcct 420  
caaaagcaag gaaaatatat actgacttca tagaaaagga agctccaaaa gagataaaca 480  
tagattttca aaccaaaaact ctgattgccc agaatataca agaagctaca agtggctgct 540

ttacaactgc	ccagaaaagg	gtatacagct	tgatggagaa	caactcttat	cctcgtttct	600
tgagagtcaga	attctaccag	gacttgtgta	aaaagccaca	aatcaccaca	gagcctcatg	660
ctacatgaaa	tgtaaaagg	agcccagaaa	tgaggagacat	ttcattcttt	ttcctgaggg	720
gaaggactgt	gacctgccat	aaagactgac	cttgaattca	gcctgggtgt	tcaggaaaca	780
tcactcagaa	ctattgattc	aaagttgggt	agtgaatcag	gaagccagta	actgactagg	840
agaagctgg	atcagaacag	cttcctcac	tgtgtacaga	acgcaagaag	ggaatagggtg	900
gtctgaacgt	ggtgtctcac	tctgaaaagc	aggaatgtaa	gatgatgaaa	gagacaatgt	960
aatactgttg	gtccaaaagc	attttaaatac	aatagatctg	ggattatgtg	gccttaggta	1020
gctgggttgta	catctttccc	taaatcgatc	catgttacca	catagtagtt	ttagtttagg	1080
attcagtaac	agtgaagtgt	ttactatgtg	caagggtatt	gaagttctta	tgaccacaga	1140
tcatcagtac	tgttgtctca	tgtaatgcta	aaactgaaat	ggtccgtgtt	tgcattgtta	1200
aaaatgatgt	gtgaaataga	atgagtgcta	tggtgttgaa	aactgcagtg	tccgttatga	1260
gtgccaaaaa	tctgtcttga	aggcagctac	actttgaagt	ggtctttgaa	tacttttaat	1320
aaatttat	tgataaataa	tattg				1345

&lt;210&gt; 2

&lt;211&gt; 5722

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

ggaagcacag	gcattccccg	cgccccctcca	gccctcgccg	ccctcgccac	cgctcccggc	60
cgccgcgctc	cggtagacac	aggatccctg	ctgggcacca	acagctccac	catggggctg	120
gcctggggac	taggcgtcct	gttcctgatg	catgtgtgtg	gcaccaaccg	cattccagag	180
tctggcggag	acaacagcgt	gtttgacatc	tttgaactca	ccggggccgc	ccgcaagggg	240
tctgggcgcc	gactgggtgaa	gggccccgac	ccttcacagc	cagctttccg	catcgaggat	300
gccaacctga	tccccctgt	gcctgatgac	aagttccaag	acctgggtgga	tgctgtgcgg	360
gcagaaaagg	gtttcctcct	tctggcatcc	ctgaggcaga	tgaagaagac	ccggggcacg	420
ctgctggccc	tgagcgggaa	agaccactct	ggccaggtct	tcagcgtggg	gtccaatggc	480
aaggcgggca	ccctggacct	cagcctgacc	gtccaaggaa	agcagcacgt	ggtgtctgtg	540

gaagaagctc	tcctggcaac	cggccagtgg	aagagcatca	ccctgtttgt	gcaggaagac	600
agggcccagc	tgtacatcga	ctgtgaaaag	atggagaatg	ctgagttgga	cgtccccatc	660
caaagcgtct	tcaccagaga	cctggccagc	atcgccagac	tccgcacgcg	aaaggggggc	720
gtcaatgaca	atttccaggg	ggtgctgcag	aatgtgaggt	ttgtcttttg	aaccacacca	780
gaagacatcc	tcaggaacaa	aggctgctcc	agctctacca	gtgtcctcct	cacccttgac	840
aacaacgtgg	tgaatggttc	cagccctgcc	atccgcacta	actacattgg	ccacaagaca	900
aaggacttgc	aagccatctg	cggcatctcc	tgtgatgagc	tgtccagcat	ggtcctggaa	960
ctcagggggc	tgcgcaccat	tgtgaccacg	ctgcaggaca	gcacccgcaa	agtgactgaa	1020
gagaacaaaag	agttggccaa	tgagctgagg	cggcctcccc	tatgctatca	caacggagtt	1080
cagtacagaa	ataacgagga	atggactgtt	gatagctgca	ctgagtgtca	ctgtcagaac	1140
tcagttacca	tctgcaaaaa	ggtgtcctgc	cccatcatgc	cctgctccaa	tgccacagtt	1200
cctgatggag	aatgctgtcc	tcgctgttgg	cccagcgact	ctgcggacga	tggctggtct	1260
ccatggtccg	agtggacctc	ctgttctacg	agctgtggca	atggaattca	gcagcgcggc	1320
cgtccttgcg	atagcctcaa	caaccgatgt	gagggctcct	cggctccagac	acggacctgc	1380
cacattcagg	agtgtgacaa	aagatttaaa	caggatggtg	gctggagcca	ctggtccccg	1440
tggtcacatt	gttctgtgac	atgtggtgat	ggtgtgatca	caaggatccg	gctctgcaac	1500
tctcccagcc	cccagatgaa	tgggaaaccc	tgtgaaggcg	aagcgcggga	gaccaaagcc	1560
tgcaagaaaag	acgcctgccc	catcaatgga	ggctggggtc	cttggtcacc	atgggacatc	1620
tgttctgtca	cctgtggagg	aggggtacag	aaacgtagtc	gtctctgcaa	caaccccgca	1680
ccccagtttg	gaggcaagga	ctgcgttggt	gatgtaacag	aaaaccagat	ctgcaacaag	1740
caggactgtc	caattgatgg	atgcctgtcc	aatccctgct	ttgccggcgt	gaagtgtact	1800
agctaccctg	atggcagctg	gaaatgtggt	gcttgtcccc	ctggttacag	tggaaatggc	1860
atccagtgca	cagatgttga	tgagtgcaaa	gaagtgcctg	atgcctgctt	caaccacaat	1920
ggagagcacc	ggtgtgagaa	cacggacccc	ggctacaact	gcctgccctg	ccccccacgc	1980
ttcacccggt	cacagccctt	cggccagggg	gtcgaacatg	ccacggccaa	caaacaggtg	2040
tgcaagcccc	gtaacccttg	cacggatggg	accacagact	gcaacaagaa	cgccaagtgc	2100
aactacctgg	gccactatag	cgaccccatg	taccgctgcg	agtgcgaagc	tggctacgct	2160
ggcaatggca	tcattctgcg	ggaggacaca	gacctggatg	gctggcccaa	tgagaacctg	2220
gtgtgcgtgg	ccaatgcgac	ttaccactgc	aaaaaggata	attgccccaa	ccttcccaac	2280
tcagggcagg	aagactatga	caaggatgga	attggtgatg	cctgtgatga	tgacgatgac	2340



aatgataaaa	ttccagatga	cagggacaac	tgtccattcc	attacaaccc	agctcagtat	2400
gactatgaca	gagatgatgt	gggagaccgc	tgtgacaact	gtccctacaa	ccacaaccca	2460
gatcaggcag	acacagacaa	caatggggaa	ggagacgcct	gtgctgcaga	cattgatgga	2520
gacggtatcc	tcaatgaacg	ggacaactgc	cagtacgtct	acaatgtgga	ccagagagac	2580
actgatatgg	atgggggttg	agatcagtgt	gacaattgcc	ccttggaaca	caatccggat	2640
cagctggact	ctgactcaga	ccgcattgga	gatacctgtg	acaacaatca	ggatattgat	2700
gaagatggcc	accagaacaa	tctggacaac	tgtccctatg	tgcccaatgc	caaccaggct	2760
gaccatgaca	aagatggcaa	gggagatgcc	tgtgaccacg	atgatgacaa	cgatggcatt	2820
cctgatgaca	aggacaactg	cagactcgtg	cccaatcccg	accagaagga	ctctgacggc	2880
gatggtcgag	gtgatgcctg	caaagatgat	tttgaccatg	acagtgtgcc	agacatcgat	2940
gacatctgtc	ctgagaatgt	tgacatcagt	gagaccgatt	tccgccgatt	ccagatgatt	3000
cctctggacc	ccaaagggac	atccccaaat	gaccctaact	gggttgtagc	ccatcagggt	3060
aaagaactcg	tccagactgt	caactgtgat	cctggactcg	ctgtaggtta	tgatgagttt	3120
aatgctgtgg	acttcagtgg	caccttcttc	atcaacaccg	aaagggacga	tgactatgct	3180
ggatattgtct	ttggctacca	gtccagcagc	cgcttttatg	ttgtgatgtg	gaagcaagtc	3240
accagtcct	actgggacac	caacccacg	agggtcagg	gatactcggg	cctttctgtg	3300
aaagtgttaa	actccaccac	agggcctggc	gagcacctgc	ggaacgccct	gtggcacaca	3360
ggaaacaccc	ctggccaggt	gcgcaccctg	tggcatgacc	ctcgtcacat	aggctggaaa	3420
gatttcaccg	cctacagatg	gcgtctcagc	cacaggccaa	agacgggttt	cattagagtg	3480
gtgatgtatg	aagggaagaa	aatcatggct	gactcaggac	ccatctatga	taaaacctat	3540
gctggtggta	gactagggtt	gtttgtcttc	tctcaagaaa	tggtgttctt	ctctgacctg	3600
aaatacgaat	gtagagatcc	ctaatacatca	aattgttgat	tgaaagactg	atcataaacc	3660
aatgctggta	ttgcaccttc	tggaactatg	ggcttgagaa	aacccccagg	atcacttctc	3720
cttggtcttc	ttcttttctg	tgcttgcatc	agtgtggact	cctagaacgt	gcgacctgcc	3780
tcaagaaaat	gcagttttca	aaaacagact	catcagcatt	cagcctccaa	tgaataagac	3840
atcttccaag	catataaaca	attgcttttg	tttccctttg	aaaaagcatc	tacttgcttc	3900
agttgggaag	gtgcccattc	cactctgcct	ttgtcacaga	gcagggtgct	attgtgaggc	3960
catctctgag	cagtggactc	aaaagcattt	tcaggcatgt	cagagaaggg	aggactcact	4020
agaattagca	aacaaaacca	ccctgacatc	ctccttcagg	aacacgggga	gcagaggcca	4080

aagcactaag gggagggcgc atacccgaga cgattgtatg aagaaaatat ggaggaactg	4140
ttacatgttc ggtactaagt ctttttcagg ggattgaaag actattgctg gatttcatga	4200
tgctgactgg cgtttagctga ttaacccatg taaataggca cttaaataga agcaggaaag	4260
ggagacaaag actggcttct ggacttcctc cctgatcccc acccttactc atcaccttgc	4320
agtggccaga attaggggaat cagaatcaaa ccagtgtgaa gcagtgtctg ctgccattgc	4380
ctggtcacat tgaaattggg ggcttcattc tagatgtagc ttgtgcagat gtagcaggaa	4440
aataggaaaa cctaccatct cagtgcacac cagctgcctc ccaaaggagg ggcagccgtg	4500
cttatatttt tatgggttaca atggcacaaa attattatca acctaaactaa aacattcctt	4560
ttctcttttt tccgtaatta ctaggtagtt ttctaattct ctcttttgga agtatgattt	4620
ttttaaagtc tttacgatgt aaaatattta ttttttactt attctggaag atctggctga	4680
aggattattc atggaacagg aagaagcgtg aagactatcc atgtcatctt tgttgagagt	4740
cttcgtgact gtaagattgt aaatacagat tatttattaa ctctgttctg cctggaaatt	4800
taggcttcat acggaaagtg tttgagagca agtagttgac atttatcagc aaatctcttg	4860
caagaacagc acaaggaaaa tcagtctaata agctgtctct gcccttgtg ctgagagtgg	4920
atgttatggg attccttttt tctctgtttt atcttttcaa gtggaattag ttggttatcc	4980
at ttgcaaat gtttttaaatt gcaaagaaag ccatgaggtc ttcaatactg ttttacccca	5040
tcccttgtgc atatttccag ggagaaggaa agcatataca cttttttctt tcatttttcc	5100
aaaagagaaa aaatgacaa aaggtgaaac ttacatacaa atattacctc atttgttgtg	5160
tgactgagta aagaattttt ggatcaagcg gaaagagttt aagtgtctaa caaacttaaa	5220
gctactgtag tacctaaaaa gtcagtgttg tacatagcat aaaaactctg cagagaagta	5280
ttcccaataa ggaaatagca ttgaaatgtt aaatacaatt tctgaaagtt atgttttttt	5340
tctatcatct ggtataccat tgctttattt ttataaatta ttttctcatt gccattggaa	5400
tagaatattc agattgtgta gatatgctat ttaaataatt tatcaggaaa tactgcctgt	5460
agagttagta tttctatttt tatataatgt ttgcacactg aattgaagaa ttgttggttt	5520
tttctttttt ttgttttttt tttttttttt tttttttttg cttttgacct cccattttta	5580
ctatttgcca ataccttttt ctaggaatgt gctttttttt gtacacattt ttatccattt	5640
tacattctaa agcagtgtaa gttgtatatt actgtttctt atgtacaagg aacaacaata	5700
aatcatatgg aaatttatat tt	5722

&lt;210&gt; 3

&lt;211&gt; 6210

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 3

```

gcaggaacag tgctagtatt gctcgagccc gagggctgga ggtagggga tgaaggctctg      60
cttccacgct ttgcactgaa ttagggctag aattggggat gggggtaggg gcgcattcct      120
tcgggagccg aggcettaagt cctcgggggtc ctgtactcga tgccgtttct cctatctctg      180
agcctcagaa ctgtcttcag ttccgtaca agggtaaaaa ggcgctctct gccccatccc      240
ccccgacctc gggaacaagg gtccgcattg aaccagggtgc gaatgtttct tctcattctg      300
cgccgttccc gcctccccctc ccccagccgc ggcccccgcc tcccccgca ctgcacctc      360
ggtgttggtc gcagcccgcg agcagttccc gtcaatccct ccccccttac acaggatgtc      420
catattagga catctgcgtc agcaggtttc cacggccttt cctgttagcc ctgggggggag      480
ccatccccga aaccctcat cttggggggc ccacgagacc tctgagacag gaactgcgaa      540
atgctcacga gattaggaca cgcgccaaag cgggggcagg gagctgcgag cgctggggac      600
gcagccgggc ggccgcagaa gcgcccaggc ccgcgcgcca cccctctggc gccaccgtgg      660
ttgagcccgt gacgtttaca ctcatccta aaacgcttgt tataaaagca gtggctgcgg      720
cgcctcgtac tccaaccgca tctgcagcga gcaactgaga agccaagact gagccggcgg      780
ccgcggcgca gcgaacgagc agtgaccgtg ctctaccga gctctgcttc acagcgccca      840
cctgtctccg cccctcggcc cctcgcccgg ctttgcttaa ccgccacgat gatgttctcg      900
ggcttcaacg cagactacga ggcgtcatcc tcccgctgca gcagcgcgtc cccggccggg      960
gatagcctct ctactacca ctacccgca gactccttct ccagcatggg ctgcctgtc      1020
aacgcgcagg taaggctggc ttcccgctgc cgcggggccc ggggcttggg gtcgcggagg      1080
aggagacacc gggcgggacg ctccagtaga tgagtagggg gctcccttgt gcctggaggg      1140
aggctgccgt ggccggagcg gtgccggctc gggggctcgg gacttgctct gagcgcacgc      1200
acgcttgcca tagtaagaat tggttcccc ttccggaggc aggttcgttc tgagcaacct      1260
ctggtctgca ctccaggacg gatctctgac attagctgga gcagacgtgt cccaagcaca      1320
aactcgctaa ctagagcctg gcttcttcgg ggagggtggc gaaagcggca atccccctc      1380
ccccggcagc ctggagcacg gaggagggat gaggaggag ggtgcagcgg gcgggtgtgt      1440

```

aaggcagttt cattgataaa aagcgagttc attctggaga ctccggagcg gcgcctgcgt	1500
cagcgcagac gtcagggata ttataacaa accccctttc aagcaagtga tgctgaaggg	1560
ataacgggaa cgcagcggca ggatggaaga gacaggcact gcgctgcgga atgcctggga	1620
ggaaaagggg gagacctttc atccaggatg agggacattt aagatgaaat gtccgtggca	1680
ggatcgtttc tcttcaactgc tgcattgcggc actgggaact cgcctcacct gtgtccggaa	1740
cctgctcgct cacgtcggct tcccccttct gttttgttct aggacttctg cacggacctg	1800
gccgtctcca gtgccaaact cattcccacg gtcactgcca tctcgaccag tccggacctg	1860
cagtggctgg tgcagccgc cctcgtctcc tctgtggccc catcgcagac cagagccct	1920
caccctttcg gagtccccgc cccctccgct ggggcttact ccagggttg cgttgtgaag	1980
accatgacag gaggccgagc gcagagcatt ggcaggaggg gcaagggtga acagggtgagg	2040
aactctagcg tactcttctt gggaatgtgg gggctgggtg ggaagcagcc cgggagatgc	2100
aggagcccag tacagaggat gaagccactg atggggcttg ctgcacatcc gtaactggga	2160
gccctggctc caagccatt ccatcccaac tcagactctg agtctcacc taagaagtac	2220
tctcatagtt tcttcctaa gtttcttacc gcatgctttc agactgggct cttctttgtt	2280
ctcttgctga ggatcttatt ttaaattgca gtcacacctt ttctgcaact gcaggctcaga	2340
aatggtttca cagtggggtg ccaggaagca gggaaagtgc aggagccagt tctactgggg	2400
tgggtgaatg gaggtgatgg cagacacttt tactgaatgt cggctttttt ttgtgattat	2460
tctagttatc tccagaagaa gaagagaaaa ggagaatccg aagggaagg aataagatgg	2520
ctgcagccaa atgccgcaac cggaggaggg agctgactga tacactcaa gcggtaggta	2580
ctctgtgggt tgctcctttt taaaacttaa gggaaagttg gagattgagc ataagggccc	2640
ttgagtaaga ctgtgtctta tgctttcctt tatccctctg tatacaggag acagaccaac	2700
tagaagatga gaagtctgct ttgcagaccg agattgccaa cctgctgaag gagaaggaaa	2760
aactagagtt catcctggca gctcacccgac ctgcctgcaa gatccctgat gacctgggct	2820
tcccagaaga gatgtctgtg gcttcccttg atctgactgg gggcctgcc gaggttgcca	2880
ccccggagtc tgaggaggcc ttcacctgc ctctcctcaa tgacctgag cccaagccct	2940
cagtggaaacc tgtcaagagc atcagcagca tggagctgaa gaccgagccc tttgatgact	3000
tctgtttccc agcatcatcc aggccagtg gctctgagac agcccgctcc gtgocagaca	3060
tggacctatc tgggtccttc tatgcagcag actgggagcc tctgcacagt ggctccctgg	3120
ggatggggcc catggccaca gagctggagc cctgtgcac tccggtggtc acctgtactc	3180
ccagctgcac tgcttacacg tcttccctcg tcttcacctc ccccgaggct gactccttcc	3240

ccagctgtgc	agctgcccac	cgcaagggca	gcagcagcaa	tgagccttcc	tctgactcgc	3300
tcagctcacc	cacgctgctg	gccctgtgag	ggggcagggg	aggggaggca	gccggcaccc	3360
acaagtgcc	ctgcccagac	tggtgcatta	cagagaggag	aaacacatct	tccctagagg	3420
gttctctgtg	acctagggag	gaccttatct	gtgcgtgaaa	cacaccaggc	tgtgggcctc	3480
aaggacttga	aagcatccat	gtgtggactc	aagtccttac	ctcttccgga	gatgtagcaa	3540
aacgcatgga	gtgtgtattg	ttcccagtga	cacttcagag	agctggtagt	tagtagcatg	3600
ttgagccagg	cctgggtctg	tgtctctttt	ctctttctcc	ttagtcttct	catagcatta	3660
actaatctat	tgggttcatt	attggaatta	acctgggtgct	ggatatatttc	aaattgtatc	3720
tagtgcagct	gattttaaca	ataactactg	tgttcctggc	aatagtgtgt	tctgattaga	3780
aatgaccaat	attatactaa	gaaaagatac	gactttatatt	tctggtagat	agaaataaat	3840
agctatatcc	atgtactgta	gtttttcttc	aacatcaatg	ttcattgtaa	tgttactgat	3900
catgcattgt	tgaggtggtc	tgaatgttct	gacattaaca	gttttccatg	aaaacgtttt	3960
attgtgtttt	taatttatatt	attaagatgg	attctcagat	atttatattt	ttattttatt	4020
tttttctacc	ttgaggtcct	ttgacatgtg	gaaagtgaat	ttgaatgaaa	aatttaagca	4080
ttgtttgctt	attgttccaa	gacattgtca	ataaaagcat	ttaagttgaa	tgcgaccaac	4140
cttgtgctct	tttcattctg	gaagtcttgt	aagtttctga	aaggatttat	tggagaccag	4200
tttgtcaaga	agggtagctg	ctggaggggg	acacaccctc	tgtctgatcc	cttatcaaag	4260
aggacaagga	aactatagag	ctgatttttag	aatatatttac	aaatacatgc	cttccattgg	4320
aatgctaaga	ttttctactg	cttctggggg	cgggaaaccg	ctgtgtaaca	gcttttgtgg	4380
gaatacattt	tttctgtttc	agtactcgca	gggggaaata	tttaaatttt	gttgtgctaa	4440
tattaaattc	agatgttttg	atcttaaagg	aaccctttaa	gcaaacagaa	cctagctttg	4500
tacagactat	tttaactttt	tattctcaca	aatcacgtg	gagggttatt	ctacttcaaa	4560
gatgagcaaa	ttgaagaatg	gttagaataa	acaactttct	tgatattccg	ttatcggcat	4620
tagaatcttc	ctgctcgtaa	tcgtatccag	caggctgaac	tgctcttga	tacttggtta	4680
aaaaaaattt	tcaggccggg	cgcggtggcc	catgcctgta	atcctagcac	tttgggaggc	4740
cgaggcaggc	ggatcacctg	aggctggggg	ttcgagacca	gcctgaccaa	catggagaaa	4800
ccccgtcttt	actaaaaata	caaaattagc	ctgggtgtgg	ggtgcatgcc	tgtaatccta	4860
gctacttgag	aggctgagac	aggaaaatca	cttgaactcg	ggaggcggat	gttgcagcga	4920
actgagattg	cgccattgca	ctccagcctg	ggcaacaaga	ttgaaactct	gtttaaaaaa	4980

```

aaaagttttc actaatgtgt acatTTTTTT gtactctttt attctogaaa gggaaggagg 5040
gctattgccc tatcccttat taataaatgc attgtgggtt ctggttttct taataccata 5100
tgcccttcat tcagttttata gtgggcggaa gtgggggaga aaaagttgct cagaaatcaa 5160
aagatatctc aaacagcaca aataatggct gatcgttctg caaacaaaaa gttacataat 5220
agctcaagaa ggagaagtca acatgactct gaacaagctt taacttagaa actttatcat 5280
cttaaggaag aacgtgacct ttgtccagga cgtctctggg aatggggcac ttacacacac 5340
atgcacacgt acaaaccaca gggaaaggag accgcccttc tgccctctgct cgcgagtatc 5400
acgcaggcac catgcactat gttttcacac acactgggtg gaagaagagc ttcagcgcca 5460
gtcttctaata gctttgggtga taatgaaaat cactgggtgc ttatggggtg tcatattcaa 5520
tcgagttaaa agtttttaatt caaatgaca gttttactga ggttgatgtt ctcgctctatg 5580
atatctctgc ccctcccata aaaatggaca tttaaaagca acttaccgct ctttagatca 5640
ctcctatata acacaccact tgggggtgctg tttctgctag acttgatgac acagtggcct 5700
taggatccct gtttgctgtt caaagggcaa atattttata gcctttaaat atacctaaac 5760
taaatacaga attaataata ctaacaaaca cctggtctga aataacaagg tgatctaccc 5820
tggaaggaac ccagctgggtg ggccaggagc ggtggctcac acctgtaatt ccagcacttt 5880
gggaggctga gacaggagga tcactggagt ccaggagttt gagaccagcc tgggcaacat 5940
ggcaaaaccc agtgtgtctc tgttgctcca gctacactac tcaggaggct gaggcaggag 6000
tatgacttga gcctgggagg gggagggtgc agagaactga tattgcacca ccactgcact 6060
ccagcctggg tgacacagca aaaccctatc tcaaaaaaaaa aaaaaaaaaa aaggaacca 6120
gctggttcct gtaggtgtgc aataataaca accagaggaa gaaaaggaag acgatttccc 6180
agatgaagaa gggcagctgg accttcggac 6210

```

&lt;210&gt; 4

&lt;211&gt; 2136

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

```

cgcgagccgc ctctccccc tccccacgct cgaggagggg ggcgcggggg cccggctccg 60
gcgaaggcca atcggagcgc acttccgtgg ctgactagcg cgggtataaag gcgtgtggct 120
caggctgagc ggctgggacc ttgagagcgg ccaggccagc ctgggagcca gcaggagct 180

```

gggagctggg	ggaaacgacg	ccaggaaagc	tatcgcgcca	gagagggcga	cgggggctcg	240
ggaagcctga	cagggctttt	gcgcacagct	gccggctggc	tgctaccgcg	ccgcgccagc	300
ccccgagaac	gcgcgaccag	gcacccagtc	cggtcaccgc	agcggagagc	tcgccgctcg	360
ctgcagcgag	gcccgggagc	gccccgcagg	gacctcccc	agaccgcctg	ggccgcccgg	420
atgtgcacta	aaatggaaca	gcccttctac	cacgacgact	catacacagc	tacgggatac	480
ggccggggccc	ctggtggcct	ctctctacac	gactacaaac	tcctgaaacc	gagcctggcg	540
gtcaacctgg	ccgaccccta	ccggagtctc	aaagcgcttg	gggctcgcgg	acccggccca	600
gagggcgggc	gtggcggcag	ctacttttct	ggtcagggct	cggacaccgg	cgcgtctctc	660
aagctcgctt	cttcggagct	ggaacgcctg	attgtcccca	acagcaacgg	cgtgatcacg	720
acgacgccta	cacccccggg	acagtacttt	tacccccgcg	ggggtggcag	cgggtggaggt	780
gcagggggcg	cagggggcg	cgtcaccgag	gagcaggagg	gcttcgccga	cggctttgtc	840
aaagccctgg	acgatctgca	caagatgaac	cacgtgacac	cccccaacgt	gtccctgggc	900
gctaccgggg	ggcccccggc	tgggccccgg	ggcgtctacg	ccggccccga	gccacctccc	960
gtttacacca	acctcagcag	ctactcccca	gcctctgcgt	cctcgggagg	cgcgggggct	1020
gccgtcggga	ccgggagctc	gtacccgacg	accaccatca	gctacctccc	acacgcgcgc	1080
cccttcgccg	gtggccaccc	ggcgagctg	ggcttggggc	gcggcgctc	caccttcaag	1140
gaggaaccgc	agaccgtgcc	ggaggcgcg	agccgggacg	ccacgccgcc	ggtgtcccc	1200
atcaacatgg	aagaccaaga	gcgcatcaaa	gtggagcgca	agcggctgcg	gaaccggctg	1260
gcggccacca	agtgccggaa	gcggaagctg	gagcgcctcg	cgcgcctgga	ggacaagggt	1320
aagacgctca	aggccgagaa	cgcggggctg	tcgagtaccg	ccggcctcct	ccgggagcag	1380
gtggcccagc	tcaaacagaa	ggtcatgacc	cacgtcagca	acggctgtca	gctgctgctt	1440
ggggtcaagg	gacacgcctt	ctgaacgtcc	cctgccccct	tacggacacc	ccctcgcttg	1500
gacggctggg	cacacgcctc	ccactggggg	ccagggagca	ggcgggtggg	acccaccctg	1560
ggacctaggg	gcgcgcgaaa	ccacactgga	ctccggccct	cctacctgc	gcccagtcct	1620
tcacacctga	cgtttacaag	cccccccttc	cacttttttt	tgtatgtttt	ttttctgctg	1680
gaaacagact	cgattcatat	tgaatataat	atatttgtgt	atttaacagg	gaggggaaga	1740
ggggggcgatc	gcggcgagc	tggccccgcc	gcctgggtact	caagcccgcg	gggacattgg	1800
gaaggggacc	cccgccccct	gccctcccc	ctctgcaccg	tactgtggaa	aagaaacacg	1860
cacttagtct	ctaaagagtt	tattttaaga	cgtgtttgtg	tttgtgtgtg	tttgttcttt	1920

ttattgaatc tatttaagta aaaaaaaaaat tggttcttta ttaatttctg ttgtcttttt	1980
ttccaagctg ggagggcggg gggaaaaaaaa aagcactggg ttgccccag ctcagtgtg	2040
ttggtggctc ggtcctgtat gtgtccccc cgtcggttcg ggcaggcat cttgtgggcc	2100
cagcccagga gtcccaccct tcccgcgtcc ccagat	2136

&lt;210&gt; 5

&lt;211&gt; 1398

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

<400> 5	
tagatggcaa cctccctatc tgcccgagg tcatagaggc gacacgtagc gtcactgtac	60
cctgaagcaa aggcattctc actccaaagt tagacaaaat gccaggaatg ttcttctctg	120
ctaaccctaaa ggaattgaaa ggaaccactc attcacttct agacgacaaa atgcaaaaaa	180
ggaggccaaa gacttttggg atggatatga aagcatacct gagatctatg atcccacatc	240
tggaaatctgg aatgaaatct tccaagtcca aggatgtact ttctgctgct gaagtaatgc	300
aatggtctca atctctggaa aaacttcttg ccaaccaaac tgggtcaaat gtctttggaa	360
gtttcctaaa gtctgaattc agtgaggaga atattgagtt ctggctggct tgtgaagact	420
ataagaaaac agagtctgat cttttgccct gtaaagcaga agagatatat aaagcatttg	480
tgcattcaga tgctgctaaa caaatcaata ttgacttccg cactcgagaa tctacagcca	540
agaagattaa agcaccaacc cccacgtgtt ttgatgaagc acaaaaagtc atatatactc	600
ttatggaaaa ggactcttat ccaggttcc tcaaatcaga tatttactta aatcttctaa	660
atgacctgca ggctaatagc cttaaagtgac tgggtccctgg ctgaaggga ttaacagata	720
gtatcaaggc acgaaggaat gtgccagtat gggtccctgg gtgaacagct tggccttttt	780
tgggtgtctt gacaggccaa gaagaacaaa tgactcagaa tggattaaca tgaaagtat	840
ccaggcgcag agttgaagaa gcataagcaa gacaaaaaca gagagaccgc agaaggagga	900
agatactgtg gtactgtcat aaaaaacagt ggagctctgt attagaaagc ccctcagaac	960
tgggaaggcc aggtaactct agttacacag aaactgtgac taaagtctat gaaactgatt	1020
acaacaggct gtaagaatca aagtcaactg acatctatgc tacatattat tatatagttt	1080
gtactgagct attgaagtcc cattaactta aagtatatgt tttcaaattg ccattgctac	1140
tattgcttgt cgggtgtattt ttttttattg tttttgactt tggaagagat gaactgtgta	1200



tttaacttaa	gctattgctc	ttaaaaccag	ggatcagaat	atatttgtaa	gttaaatacat	1260
tggtgcta	aatataatgtg	gattttgtat	taaaatatat	agaagcaatt	tctgtttaca	1320
tgctcctt	gctactttaaa	acttgcattt	attcctcaga	ttttaaaaaat	aaataaataa	1380
ttcattttaa	aaaaaaaaa					1398

&lt;210&gt; 6

&lt;211&gt; 2000

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 6

tttgggctgt	gtgtgctgacg	cgggtcggag	gggcagtcgg	gggaaccgcg	aagaagccga	60
ggagcccga	gccccgcgtg	acgtcctct	ctcagtcctaa	aagcggcttt	tggttcggcg	120
cagagagacc	cgggggtcta	gcttttctc	gaaaagcgcc	gccctgccct	tggccccgag	180
aacagacaaa	gagcaccgca	gggcccgatca	cgtcgggggc	gctgaggccg	gccatgggtca	240
tggaagtggg	caccctggac	gctggaggcc	tgccggcgct	gctgggggag	cgagcggcgc	300
aatgcctgct	gctggactgc	cgtccttct	tcgctttcaa	cgcggccac	atcgccggct	360
ctgtcaacgt	gcgcttcagc	accatcgtgc	ggcgccgggc	caagggcgcc	atgggcctgg	420
agcacatcgt	gccccacgcc	gagctccgcg	gccgcctgct	ggccggcgcc	taccacgccg	480
tggtgttgct	ggacgagcgc	agcgccgccc	tggacggcgc	caagcgcgac	ggcaccctgg	540
ccctggcggc	cggcgcgctc	tgcgcgagg	cgcgcgcgc	gcaagtcttc	ttcctcaaag	600
gaggatacga	agcgttttcg	gcttctctgc	cggagctgtg	cagcaaacag	tcgaccccca	660
tggggctcag	ccttccccctg	agtactagcg	tccctgacag	cgcggaatct	gggtgcagtt	720
cctgcagtac	cccactctac	gatcagggtg	gcccggtgga	aatcctgccc	tttctgtacc	780
tgggcagtgc	gtatcacgct	tcccgaagg	acatgctgga	tgccttgggc	ataactgcct	840
tgatcaacgt	ctcagccaat	tgtcccaacc	atattgaggg	tcactaccag	tacaagagca	900
tccctgtgga	ggacaaccac	aaggcagaca	tcagctcctg	gttcaacgag	gccattgact	960
tcatagactc	catcaagaat	gctggaggaa	gggtgtttgt	ccactgccag	gcaggcattt	1020
cccggtcagc	caccatctgc	cttgcttacc	ttatgaggac	taatcgagtc	aagctggacg	1080
aggcctttga	gtttgtgaag	cagaggcgaa	gcacatctc	tcccaacttc	agcttcatgg	1140

gccagctgct gcagtttgag tcccaggtgc tggctccgca ctgttcggca gaggctggga 1200  
 gccccgccat ggctgtgctc gaccgaggca cctccaccac caccgtgttc aacttccccg 1260  
 tctccatccc tgtccactcc acgaacagtg cgctgagcta ccttcagagc cccattacga 1320  
 cctctcccag ctgctgaaag gccacgggag gtgaggctct tcacatccca ttgggactcc 1380  
 atgctccttg agaggagaaa tgcaataact ctgggagggg ctcgagaggg ctggtcctta 1440  
 tttattttaac ttcacccgag ttctcttggg tttctaagca gttatggtga tgacttagcg 1500  
 tcaagacatt tgctgaactc agcacattcg ggaccaatat atagtgggta catcaagtcc 1560  
 atctgacaaa atggggcaga agagaaaagga ctcagtgtgt gatccggttt ctttttgtct 1620  
 gccctgtttt tttgtagaat ctcttcattg ttgacatacc taccagtatt attcccgcg 1680  
 acacatatac atatgagaat ataccttatt tatttttgtg taggtgtctg ccttcacaaa 1740  
 tgtcattgtc tactcctaga agaaccaaat acctcaattt ttgtttttga gtactgtact 1800  
 atcctgtaaa tatatcttaa gcaggtttgt tttcagcact gatggaaaat accagtgttg 1860  
 ggtttttttt tagttgcaa cagttgtatg tttgctgatt atttatgacc tgaaataata 1920  
 tatttcttct tctaagaaga ctttttgta cataaggatg acttttttat acaatggaat 1980  
 aaattatggc atttctattg 2000

<210> 7

<211> 1110

<212> DNA

<213> Homo sapiens

<400> 7

gacagagccc gggccacgga gctccttgcc agctctcctc ctgcacagc cgctcgaacc 60  
 gcctgctgag ccccatggcc cgcgccacgc tctccgccgc cccagcaat ccccggtcc 120  
 tgccgggtggc gctgctgctc ctgctcctgg tggccgccag ccggcgcgca gcaggagcgc 180  
 ccctggccac tgaactgcgc tgccagtgtc tgcagaccct gcagggaatt cacctcaaga 240  
 acatccaaag tgtgaagggtg aagtcccccg gacccactg cgcccaaacc gaagtcatag 300  
 ccacactcaa gaatgggcag aaagcttgct tcaacccgc atcgcccatg gttaagaaaa 360  
 tcatcgaaaa gatgctgaaa aatggcaaat ccaactgacc agaaggaagg aggaagctta 420  
 ttggtggctg ttctgaagg aggcctgcc ttacaggaac agaagaggaa agagagacac 480  
 agctgcagag gccacctggc ttgcgcctaa tgtgtttgag catacttagg agaagtcttc 540

tattttatttta tttattttatt tatttgtttg ttttagaaga ttctatgtta atattttatg 600  
tgtaaaataa ggttatgatt gaatctactt gcacactctc ccattatatt tattgtttat 660  
tttaggtcaa acccaagtta gttcaatcct gattcatatt taatttgaag atagaagggtt 720  
tgcagatatt ctctagtcac ttgttaatat ttcttcgtga tgacatatca catgtcagcc 780  
actgtgatag aggctgagga atccaagaaa atggccagta agatcaatgt gacggcaggg 840  
aaatgtatgt gtgtctatct tgtaactgta aagatgaatg tcagttgtta tttattgaaa 900  
tgatttcaca gtgtgtgggc aacatttctc atgttgaagc ttttaagaact aaaatgttct 960  
aaatatccct tggcatttta tgtctttctt gtaagatact gccttgttta atgttaatta 1020  
tgcagtgttt ccctctgtgt tagagcagag aggtttcgat atttattgat gttttcacaa 1080  
agaacaggaa aataaaatat ttaaaaatat 1110

<210> 8

<211> 1064

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (27)..(27)

<223> n=a, c, g or t

<220>

<221> misc\_feature

<222> (766)..(766)

<223> n=a, c, g or t

<400> 8

cacagccggg tcgcaggcac ctcccngcc agctctcccg cattctgcac agcttcccga 60  
cgcgctctgt gagcccatg gccacgcca cgctctccgc cgccccagc aatccccggc 120  
tcctgcgggt ggcgctgtg ctctgctcc tgggtgggcag ccggcgcgca gcaggagcgt 180

ccgtgggtcac	tgaactgcgc	tgccagtgct	tgacagacact	gcaggggaatt	cacctcaaga	240
acatccaaag	tgtgaatgta	aggccccccg	gacccccactg	cgcccaaacc	gaagtcatag	300
ccacactcaa	gaatgggaag	aaagcttgct	tcaacccccg	atcccccatg	gttcagaaaa	360
tcatcgaaaa	gatactgaac	aaggggagca	ccaactgaca	ggagagaagt	aagaagctta	420
tcagcgtatc	attgacactt	cctgcagggt	ggccccctgcc	cttaccagag	ctgaaaatga	480
aaaagagaac	agcagctttc	tagggacagc	tggaaaggga	cttaatgtgt	ttgactattt	540
cttacgaggg	ttctacttat	ttatgtattt	atTTTTgaaa	gcttgatatt	taatatttta	600
catgctgtta	tttaaagatg	tgagtgtgtt	tcatcaaaca	tagctcagtc	ctgattattt	660
aattggaata	tgatgggttt	taaatgtgtc	attaaactaa	tatttagtgg	gagaccataa	720
tgtgtcagcc	accttgataa	atgacagggt	ggggaactgg	agggtngggg	gattgaaatg	780
caagcaatta	gtggatcact	gttagggtaa	gggaatgtat	gtacacatct	atTTTTtata	840
ctTTTTTTTT	taaaaaagaa	tgtcagttgt	tatttattca	aattatctca	cattatgtgt	900
tcaacatttt	tatgctgaag	tttcccttag	acattttatg	tcttgcttgt	agggcataat	960
gccttgTTTT	atgtccattc	tgacgcgttt	ctctttccct	tggaaaagag	aatttatcat	1020
tactgtttaca	tttgtacaaa	tgacatgata	ataaaagtgt	tatg		1064

&lt;210&gt; 9

&lt;211&gt; 1172

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 9

gagacattcc	tcaattgctt	agacatattc	tgagcctaca	gcagaggaac	ctccagtctc	60
agcaccatga	atcaaactgc	gattctgatt	tgctgcctta	tctttctgac	tctaagtggc	120
attcaaggag	tacctctctc	tagaaccgta	cgctgtacct	gcatcagcat	tagtaatcaa	180
cctgttaatc	caaggctctt	agaaaaactt	gaaattattc	ctgcaagcca	atTTTgtcca	240
cgtgttgaga	tcattgctac	aatgaaaaag	aagggtgaga	agagatgtct	gaatccagaa	300
tcgaaggcca	tcaagaatTT	actgaaagca	gttagcaagg	aaatgtctaa	aagatctcct	360
taaaaccaga	ggggagcaaa	atcgatgcag	tgcttccaag	gatggaccac	acagaggctg	420
cctctcccat	cacttcccta	catggagtat	atgtcaagcc	ataattgttc	ttagtttgca	480
gttacactaa	aaggtagcca	atgatgggtc	ccaaatcagc	tgctactact	cctgtaggaa	540

```

ggttaatggt catcatccta agctattcag taataactct accctggcac tataatgtaa 600
gctctactga ggtgctatgt tcttagtgga tgttctgacc ctgcttcaaa tatttccttc 660
acctttccca tcttccaagg gtactaagga atctttctgc tttgggggttt atcagaattc 720
tcagaatctc aaataactaa aaggatgca atcaaactctg ctttttaaaag aatgctcttt 780
acttcatgga cttccactgc catcctccca agggggccca attctttcag tggctaccta 840
catacaattc caaacacata caggaaggta gaaatatctg aaaatgtatg tgtaagtatt 900
cttatttaat gaaagactgt acaaagtata agtcttagat gtatatatctt cctatatgtt 960
tttcagtgt catggaataa catgtaatta agtactatgt atcaatgagt aacaggaaaa 1020
ttttaaaaa acagatagat atatgctctg catgttacat aagataaatg tgctgaatgg 1080
ttttcaaata aaaatgaggt actctcctgg aaatatataag aaagactatc taaatgttga 1140
aagatcaaaa ggttaataaa gtaattataa ct 1172

```

<210> 10

<211> 4068

<212> DNA

<213> Homo sapiens

<400> 10

```

gaattcctcc tctcttcacc ccgttagctg ttttcaatgt aatgctgccc tccttctctt 60
gcactgcctt ctgcgctaac acctccattc ctgtttataa ccgtgtattt attacttaat 120
gtatataatg taatgttttg taagttatta atttatatat ctaacattgc ctgccaatgg 180
tggtgttaaa tttgtgtaga aaactctgcc taagagttac gactttttct tgtaatgttt 240
tgtattgtgt attatataac ccaaacgtca cttagtagag acatatggcc cccttggcag 300
agaggacagg ggtgggcttt tgttcaaagg gtctgccctt tccctgcctg agttgctact 360
tctgcacaac ccctttatga accagttttc acccgaattt tgactgtttc atttagaaga 420
aaagcaaaat gagaaaaagc tttcctcatt tctccttgag atggcaaagc actcagaaat 480
gacatcacat accctaaaga accctgggat gactaaggca gagagagtct gagaaaactc 540
tttgggtgctt ctgcctttag ttttaggaca catttatgca gatgagctta taagagaccg 600
ttccctccgc cttcttcttc agaggaagtt tcttggtaga tcaccgacac ctcattccagg 660
cgggggggttg gggggaaact tggcaccagc catcccaggc agagcaccac tgtgatttgt 720

```

tctcctggtg	gagagagctg	gaaggaagga	gccagcgtgc	aaataatgaa	ggagcacggg	780
ggcaccttca	gtagcaccgg	aatcagcggg	ggtagcgggtg	actctgctat	ggacagcctg	840
cagccgctcc	agcctaacta	catgcctgtg	tgtttgtttg	cagaagaatc	ttatcaaaaa	900
ttagcaatgg	aaacgctgga	ggaattagac	tggtgttttag	accagctaga	gaccatacag	960
acctaccggt	ctgtcagtga	gatggcttct	aacaagttca	aaagaatgct	gaaccgggag	1020
ctgacacacc	tctcagagat	gagccgatca	gggaaccagg	tgtctgaata	cattttcaaat	1080
actttcttag	acaagcagaa	tgatgtggag	atcccatctc	ctaccagaa	agacagggag	1140
aaaaagaaaa	agcagcagct	catgaccag	ataagtggag	tgaagaaatt	aatgcatagt	1200
tcaagcctaa	acaatacaag	catctcacgc	tttggagtca	acactgaaaa	tgaagatcac	1260
ctggccaagg	agctggaaga	cctgaacaaa	tggggcttta	acatctttta	tgtggctgga	1320
tattctcaca	atagaccctt	aacatgcac	atgtatgcta	tattccagga	aagagacctc	1380
ctaaagacat	tcagaatctc	atctgacaca	tttataacct	acatgatgac	tttagaagac	1440
cattaccatt	ctgacgtggc	atatcacaac	agcctgcacg	ctgctgatgt	agcccagtcg	1500
acctatgttc	tcctttctac	accagcatta	gacgctgtct	tcacagattt	ggagatcctg	1560
gctgccatth	ttgcagctgc	catccatgac	gttgatcatc	ctggagtctc	caatcagtht	1620
ctcatcaaca	caaattcaga	acttgctttg	atgtataatg	atgaatctgt	gttggaaaat	1680
catcaccttg	ctgtgggttt	caaactgctg	caagaagaac	actgtgacat	cttcatgaat	1740
ctcaccaaga	agcagcgtca	gacactcagg	aagatgggta	ttgacatggg	gttagcaact	1800
gatatgtcta	aacatatgag	cctgctggca	gacctgaaga	caatggtaga	aacgaagaaa	1860
gttacaagtt	caggcgtht	tctcctagac	aactataccg	atcgcatthca	ggthcttcgc	1920
aacatgggtac	actgtgcaga	cctgagcaac	cccaccaagt	ccttggaatt	gtatcggcaa	1980
tggacagacc	gcatcatgga	ggaatttttc	cagcagggag	acaaagagcg	ggagagggga	2040
atggaaatta	gccaatgtg	tgataaacac	acagcttctg	tggaaaaatc	ccaggthggg	2100
ttcatcgact	acattgtcca	tcattgtgg	gagacatggg	cagattthgg	acagcctgat	2160
gctcaggaca	ttctcgatac	cttagaagat	aacaggaact	ggtatcagag	catgatacct	2220
caaagtccct	caccaccact	ggacgagcag	aacagggact	gccagggtht	gatggagaag	2280
tttcagthtg	aactgactct	cgatgaggaa	gattctgaag	gacctgagaa	ggagggagag	2340
ggacacagct	atttcagcag	cacaaagacg	ctthgtgtga	ttgatccaga	aaacagagat	2400
tcctggggag	agactgacat	agacattgca	acagaagaca	agthccccgt	ggatacataa	2460
thccccctct	cctgtggaga	tgaacattct	atcctthgatg	agcatgccag	ctatthggta	2520

gggccagccc accatggggg ccaagacctg cacaggacaa gggccacctg gcctttcagt	2580
tacttgagtt tggagtcaga aagcaagacc aggaagcaaa tagcagctca ggaaatccca	2640
cggttgactt gccttgatgg caagcttggt ggagagggct gaagctgttg ctggggggccg	2700
attctgatca agacacatgg cttgaaaatg gaagacacaa aactgagaga tcattctgca	2760
ctaagtttcg ggaacttatc cccgacagtg actgaactca ctgactaata acttcattta	2820
tgaatcttct cacttgcccc tttgtctgcc aacctgtgtg ccttttttgt aaaacatttt	2880
catgtcttta aaatgcctgt tgaataacctg gagtttagta tcaacttcta cacagataag	2940
ctttcaaagt tgacaaaactt ttttgactct ttctggaaaa gggaaagaaa atagtcttcc	3000
ttctttcttg ggcaatatcc ttcactttac tacagttact tttgcaaaca gacagaaagg	3060
atacacttct aaccacattt tacttccttc ccctgttgtc cagtccaact ccacagtcac	3120
tcttaaaaact tctctctgtt tgccctgcctc caacagtact ttttaactttt tgctgtaaac	3180
agaataaaaat tgaacaaatt agggggtaga aaggagcagt ggtgtcgttc accgtgagag	3240
tctgcataga actcagcagt gtgccctgct gtgtcttgga ccctgcccc cacaggagtt	3300
gctacagtcc ctggccctgc tccccatct cctctcttca ccccgtagc tgttttcaat	3360
gtaatgctgc cgtccttctc ttgcactgcc ttctgcgcta acacctccat tcctgtttat	3420
aaccgtgtat ttattactta atgtatataa tgtaatgttt tgtaagttat taatttatat	3480
atctaacatt gcctgccaat ggtgggtgta aatttggtga gaaaactctg cctaagagtt	3540
acgacttttt cttgtaatgt tttgtattgt gtattatata acccaaactg cacttagtag	3600
agacatatgg ccccttggc agagaggaca ggggtgggct tttgttcaaa ggggtctgcc	3660
tttccctgcc tgagttgcta cttctgcaca acccctttat gaaccagttt tggaaacaat	3720
attctcacat tagatactaa atgggtttata ctgagtcttt tacttttgta tagcttgata	3780
ggggcagggg caatgggatg tagtttttac ccaggttcta tccaaatcta tgtgggcatg	3840
agttgggtta taactggatc ctactatcat tgtggctttg gttcaaaagg aaacactaca	3900
tttgctcaca gatgattctt ctgattcttc tgaatgctcc cgaactactg actttgaaga	3960
ggtagcctcc tgccctgcat taagcaggaa tgtcatgttc cagttcatta caaaagaaaa	4020
caataaaaaca atgtgaattt ttataataaa aaaaaaaaaa aggaattc	4068

&lt;210&gt; 11

&lt;211&gt; 2900

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

atgagtgtca gaagtgtgaa ggggtgcctgt tctgaatccc agagcctcct ctccctctgt	60
gaggctggca ggtgaggaag ggtttaacct cactggaagg aatccctgga gctagcggct	120
gctgaaggcg tcgaggtgtg ggggcacttg gacagaacag tcaggcagcc gggagctctg	180
ccagctttgg tgaccttggg ccgggctggg agcgctgcgg cgggagccgg aggactatga	240
gctgccgcgc gttgtccaga gccagccca gccctacggc cgcgcccgag agctctgttc	300
cctggaactt tgggcactgc ctctgggacc cctgccggcc agcaggcagg atgggtgcttg	360
cctcgtgccc cttggtgccc gtctgctgat gtgcccagcc tgtgcccgcc atgccgccct	420
ccatctcagc tttccaggcc gcctacatcg gcctcagagg gctcatcgcc ctgggtctctg	480
tgcccgggaa cgtgctggtg atctgggagg tgaagggtgaa ccaggcgctg cgggatgcca	540
ccttctgctt catcgtgtcg ctggcggttg ctgatgtggc cgtgggtgcc ctgggtcatcc	600
ccctcgccat cctcatcaac attgggccac agacctactt ccacacctgc ctcatgggtg	660
cctgtccggg cctcatcctc acccagagct ccatacctggc cctgctggca attgctgtgg	720
accgctacct ccgggtcaag atccctctcc ggtacaagat ggtggtgacc ccccgagggg	780
cggcggtggc catagccggc tgctggatcc tctccttcgt ggtgggactg acccctatgt	840
ttggctggaa caatctgagt gcggtggagc gggcctgggc agccaacggc agcatggggg	900
agcccgtgat caagtgcgag ttcgagaagg tcatcagcat ggagtacatg gtctacttca	960
acttctttgt gtgggtgctg cccccgcttc tctcatgggt cctcatctac ctggaggtct	1020
tctacctaat ccgcaagcag ctcaacaaga aggtgtcggc ctctccggc gaccgcaga	1080
agtactatgg gaaggagctg aagatcgcca agtcgctggc cctcatcctc ttcctctttg	1140
ccctcagctg gctgcctttg cacatcctca actgcatcac cctcttctgc ccgtcctgcc	1200
acaagcccag catccttacc tacattgcca tcttctcac gcacggcaac tcggccatga	1260
acccatttgt ctatgccttc cgcattcaga agttccgcgt caccttcctt aagatttgga	1320
atgaccattt ccgctgccag cctgcacctc ccattgacga ggatctccca gaagagaggc	1380
ctgatgacta gacccgcct tccgctccca ccagcccaca tccagtgggg tctcagtcca	1440



```

gtcctcacat gcccgctgtc ccaggggtct ccctgagcct gcccagctg ggctgttggc 1500
tggtggcatg ggggaggctc tgaagagata cccacagagt gtgggtccctc cactaggagt 1560
taactaccct acacctcttg gccctgcagg aggcctggga gggcaagggt cctacggagg 1620
gaccaggtgt ctagaggcaa cagtgttctg agccccacc tgcctgacca tcccatgagc 1680
agtccagcgc ttcagggctg ggcaggtcct ggggaggctg agactgcaga ggagccacct 1740
gggctgggag aagggtgcttg ggcttctgcg gtgaggcagg ggagtctgct tgtcttagat 1800
gttggtggtg cagccccagg accaagctta aggagaggag agcatctgct ctgagacgga 1860
tggaaggaga gaggttgagg atgcactggc ctgttctgta ggagagactg gccagaggca 1920
gctaaggggc aggaatcaag gagcctccgt tcccacctct gaggactctg gacccaggc 1980
cataccaggt gctaggggtg ctgctctcct tgccctgggc cagcccagga ttgtacgtgg 2040
gagaggcaga aagggtaggt tcagtaatca tttctgatga ttgctggag tgctggctcc 2100
acgccctggg gagtgagctt ggtgcggtag gtgctggcct caaacagcca cgaggtggta 2160
gctctgagcc ctcttcttg ccctgagctt tccggggagg agcctggagt gtaattacct 2220
gtcatctggg ccaccagctc cactggcccc cgttgccggg cctggactgt cctaggtgac 2280
cccatctctg ctgcttcttg gcctgatgga gaggagaaca ctagacatgc caactcggga 2340
gcattctgcc tgccctggga cggggtggac gagggagtgt ctgtaaggac tcagtgttga 2400
ctgtaggcgc ccctggggtg ggtttagcag gctgcagcag gcagaggagg agtaccccc 2460
tgagagcatg tgggggaagg ccttgctgtc atgtgaatcc ctcaataccc ctagtatctg 2520
gctgggtttt caggggcttt ggaagctctg ttgcaggtgt ccgggggtct aggactttag 2580
ggatctggga tctggggaag gaccaacca tgccctgcc agcctggagc ccctgtgttg 2640
gggggcaagg tgggggagcc tggagccct gtgtgggagg gcgaggcggg ggagcctgga 2700
gcccctgtgt gggagggcga ggcgggggat cctggagccc ctgtgtcggg gggcgaggga 2760
ggggaggtgg ccgtcggttg accttctgaa catgagtgtc aactccagga cttgcttcca 2820
agcccttccc tctgttgga attgggtgtg ccctggctcc caagggaggc ccatgtgact 2880
aataaaaaac tgtgaaccct 2900

```

&lt;210&gt; 12

&lt;211&gt; 1716

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

<400> 12  
gccaactccg tgggtggtctg ggtgaatatc caggccaaga ccacaggcta tgacacgcac 60  
tgctacatct tgaacctggc cattgccgac ctgtgggttg tcctcaccat ccagtcctgg 120  
gtggtcagtc tcgtgcagca caaccagtgg cccatgggag agctcacgtg caaagtcaca 180  
cacctcatct tctccatcaa cctcttcagc agcattttct tcctcacgtg catgagcgtg 240  
gaccgctacc tctccatcac ctacttcacc aacaccccca gcagcaggaa gaagatggta 300  
cgccgtgtcg tctgcatcct ggtgtggctg ctggccttct gcgtgtctct gcctgacacc 360  
tactacctga agaccgtcac gtctgcgtcc aacaatgaga cctactgccg gtcccttctac 420  
cccgagcaca gcatcaagga gtggctgata ggcattggag tggctctcgt tgtcttgggc 480  
tttgccgttc ccttctccat tctgctgtc ttctacttcc tgctggccag agccatctcg 540  
gcgtccagtg accaggagaa gcacagcagc cggaagatca tcttctccta cgtggtggtc 600  
ttccttgtct gctggttgcc ctaccacgtg gcggtgtgc tggacatctt ctccatcctg 660  
cactacatcc ctttcacctg ccggtgtggag cagccctctt tcacggccct gcatgtcaca 720  
cagtgcctgt cgctggtgca ctgctgcgtc aaccctgtcc tctacagctt catcaatcgc 780  
aactacaggt acgagctgat gaaggccttc atcttcaagt actcggccaa aacagggctc 840  
accaagctca tcgatgcctc cagagtctca gagacggagt actctgcctt ggagcagagc 900  
accaaagat ctgccctgga gaggtctgtg gacgggttta cttgtttttg aacaggggtga 960  
tgggcccctat ggttttctag agcaaagcaa agtagcttcg ggtcttgatg cttgagtaga 1020  
gtgaagaggg gagcacgtgc cccctgcata cattytctct ttctcttgat gacgcagctg 1080  
tcatttggct gtgcgtgtg acagttttgc aacaggcaga gctgtgtcgc acagcagtg 1140  
tgtgcgtcag agccagctga ggacaggctt gcctggactt ctgtaagata ggattttctg 1200  
tgtttcctga attttttata tgggtgatttg tatttaaatt ttaagacttt attttctcac 1260  
tattggtgta cttataaat gtatttgaaa gttaaataata ttttaaataat tgtttgggag 1320  
gcatagtgtc gacatatatt cagagtgttg tagttttaag gttagcgtga ctttcagttt 1380  
tgactaagga tgacactaat tgtagctgt tttgaaatta tatatatata aatatataaa 1440  
tatatgccag tcttggctga aatgttttat ttaccatagt ttatatctg tgtggtgttt 1500  
tgtaccggca cgggatatgg aacgaaaact gctttgtaat gcagtttgtg acattaatag 1560  
tattgtaaag ttacatttta aaataaaciaaaaactgttc tggactgcaa atctgcacac 1620  
acaacgaaca gttgcatttc agagagttct ctcaatttgt aagttatttt tttttaataa 1680

agattttttgt ttcctaaaaa aaaaaaaaaa aaaaaa 1716

<210> 13

<211> 3565

<212> DNA

<213> Homo sapiens

<400> 13

gcagccggggc ggccgcagaa ggcgccaggc ccgcgcgcca cccctctggc gccaccgtgg 60  
ttgagcccgt gacgtttaca ctcattcata aaacgcttgt tataaaagca gtggctgcgg 120  
cgcctcgtac tccaaccgca tctgcagcga gcaactgaga agccaagact gagccggcgg 180  
ccgcgggcgca gcgaacgagc agtgaccgtg ctcctaccca gctctgcttc acagcgccca 240  
cctgtctccg cccctcggcc cctcgcccgg ctttgcctaa ccgccacgat gatgttctcg 300  
ggcttcaacg cagactacga ggcgtcatcc tcccgtgca gcagcgcgtc cccggccggg 360  
gatagcctct cttactacca ctcaccgcga gactccttct ccagcatggg ctgcgctgtc 420  
aacgcgcagg taaggctggc ttcccgtcgc cgcggggccg ggggcttggg gtcgcggagg 480  
aggagacacc gggcgggacg ctccagtaga tgagtagggg gctcccttgt gcctggaggg 540  
aggctgccgt ggccggagcg gtgccggctc gggggctcgg gacttgctct gagcgcacgc 600  
acgcttgcca tagtaagaat tggttcccc ttcgggaggg aggttcgttc tgagcaacct 660  
ctggtctgca ctccaggacg gatctctgac attagctgga gcagacgtgt cccaagcaca 720  
aactcgctaa ctagagcctg gcttcttcgg ggaggtggca gaaagcggca atccccctc 780  
ccccggcagc ctggagcacg gaggagggat gagggaggag ggtgcagcgg gcgggtgtgt 840  
aaggcagttt cattgataaa aagcgagttc attctggaga ctccggagcg gcgcctgcgt 900  
cagcgcagac gtcagggata ttataacaa accccctttc aagcaagtga tgctgaaggg 960  
ataacgggaa cgcagcggca ggatggaaga gacaggcact gcgctgcgga atgcctggga 1020  
ggaaaagggg gagaccttct atccaggatg agggacattt aagatgaaat gtccgtggca 1080  
ggatcgtttc tcttactgc tgcattcggc actgggaact cgccccacct gtgtccggaa 1140  
cctgctcgct cacgtcggct ttccccctct gttttgttct aggacttctg cacggacctg 1200  
gccgtctcca gtgccaaact cattcccacg gtcactgcca tctcgaccag tccggacctg 1260  
cagtggctgg tgcagccgc cctcgtctcc tctgtggccc catcgcagac cagagccct 1320  
caccctttcg gagtccccgc cccctccgct ggggcttact ccagggctgg cgttgtgaag 1380

accatgacag	gaggccgagc	gcagagcatt	ggcaggaggg	gcaaggtgga	acaggtgagg	1440
aactctagcg	tactcttcct	gggaatgtgg	gggctgggtg	ggaagcagcc	ccggagatgc	1500
aggagcccag	tacagaggat	gaagccactg	atggggctgg	ctgcacatcc	gtaactggga	1560
gccctggctc	caagcccatt	ccatcccaac	tcagactctg	agtctcaccc	taagaagtac	1620
tctcatagtt	tcttcctaa	gtttcttacc	gcattgctttc	agactgggct	cttctttgtt	1680
ctcttgctga	ggatcttatt	ttaaatgcaa	gtcacaccta	ttctgcaact	gcaggtcaga	1740
aatggtttca	cagtgggggtg	ccaggaagca	gggaagctgc	aggagccagt	tctactgggg	1800
tgggtgaatg	gaggtgatgg	cagacacttt	tactgaatgt	cggctctttt	ttgtgattat	1860
tctagttatc	tccagaagaa	gaagagaaaa	ggagaatccg	aagggaaagg	aataagatgg	1920
ctgcagccaa	atgccgcaac	cggaggaggg	agctgactga	tacactccaa	gcggtaggta	1980
ctctgtgggt	tgctcctttt	taaaacttaa	gggaaagttg	gagattgagc	ataagggccc	2040
ttgagtaaga	ctgtgtctta	tgctttcctt	tatccctctg	tatacaggag	acagaccaac	2100
tagaagatga	gaagtctgct	ttgcagaccg	agattgccaa	cctgctgaag	gagaaggaaa	2160
aactagagtt	catcctggca	gctcaccgac	ctgcctgcaa	gatccctgat	gacctgggct	2220
tcccagaaga	gatgtctgtg	gcttcctttg	atctgactgg	gggcctgcca	gaggttgcca	2280
ccccggagtc	tgaggaggcc	ttcaccttgc	ctctcctcaa	tgacctgag	cccaagccct	2340
cagtggaacc	tgtcaagagc	atcagcagca	tggagctgaa	gaccgagccc	tttgatgact	2400
tcttgttccc	agcatcatcc	aggcccagtg	gctctgagac	agcccgtcc	gtgccagaca	2460
tggacctatc	tgggtccttc	tatgcagcag	actgggagcc	tctgcacagt	ggctccctgg	2520
ggatggggcc	catggccaca	gagctggagc	ccctgtgcac	tccggtggtc	acctgtactc	2580
ccagctgcac	tgcttacacg	tcttccttcg	tcttcaccta	ccccgaggct	gactccttcc	2640
ccagctgtgc	agctgcccac	cgcaagggca	gcagcagcaa	tgagccttcc	tctgactcgc	2700
tcagctcacc	cacgctgctg	gccctgtgag	ggggcaggga	aggggaggca	gccggcaccc	2760
acaagtgcc	ctgcccagagc	tggtgcatta	cagagaggag	aaacacatct	tccctagagg	2820
gttcctgtag	acctagggag	gaccttatct	gtgcgtgaaa	cacaccaggc	tgtgggcctc	2880
aaggacttga	aagcatccat	gtgtggactc	aagtccctac	ctcttcggga	gatgtagcaa	2940
aacgcatgga	gtgtgtattg	ttcccagtga	cacttcagag	agctggtagt	tagtagcatg	3000
ttgagccagg	cctgggtctg	tgtctctttt	ctctttctcc	ttagtcttct	catagcatta	3060
actaatctat	tgggttcatt	attggaatta	acctggtgct	ggatattttc	aaattgtatc	3120

tagtgcagct	gattttaaca	ataactactg	tgttcctggc	aatagtgtgt	tctgattaga	3180
aatgaccaat	attatactaa	gaaaagatac	gactttatatt	tctggtagat	agaaataaat	3240
agctatatcc	atgtactgta	gtttttcttc	aacatcaatg	ttcattgtaa	tgttactgat	3300
catgcattgt	tgagggtggc	tgaatgttct	gacattaaca	gttttccatg	aaaacgtttt	3360
attgtgtttt	taatttatatt	attaagatgg	attctcagat	atattatatt	ttattttatt	3420
tttttctacc	ttgaggctct	ttgacatgtg	gaaagtgaat	ttgaatgaaa	aatttaagca	3480
ttgtttgctt	attgttccaa	gacattgtca	ataaaagcat	ttaagttgaa	tgcgaccaac	3540
cttgtgctct	tttcattctg	gaagt				3565

&lt;210&gt; 14

&lt;211&gt; 982

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 14

agacaatttt	cagcaggaag	aagtagaaag	gataaaatgg	atcctgcacc	acgggaacct	60
cacagcacct	cacttctttt	ggttttcttt	ctctttgggg	cacctctgga	ctcactcccc	120
agcatgaagg	cgctgagccc	ggtgcgcggc	tgctacgagg	cggtgtgctg	cctgtcggaa	180
cgcagtctgg	ccatcgcccc	gggccgaggg	aagggcccg	cagctgagga	gccgctgagc	240
ttgctggacg	acatgaacca	ctgctactcc	cgcttgcggg	aactgggtacc	cggagtcccc	300
agaggcactc	agcttagcca	ggtggaaatc	ctacagcgcg	tcacgacta	cattctcgac	360
ctgcaggtag	tcctggccga	gccagcccc	ggacccccct	atggccccca	ccttcccatc	420
cagacagccg	agctcgctcc	ggaacttgtc	atctccaacg	acaaaaggag	cttttgccac	480
tgactcggcc	gtgtcctgac	acctccagaa	cgcagggtgt	ggcgcccgtt	ctgcctggga	540
ccccgggaac	ctctcctgcc	ggaagccgga	cggcagggat	gggccccaac	ttcgccctgc	600
ccacttgact	tcaccaaata	ccttcctgga	gactgaacct	ggtgctcagg	agcgaaggac	660
tgtgaacttg	tggcctgaag	agccagagct	agctctggcc	accagctggg	cgacgtcacc	720
ctgctccac	cccaccccaa	gttctaaggt	cttttcagag	cgtggagggt	tggaaggagt	780
ggctgctctc	caaactatgc	caaggcgggc	gcagagctgg	tcttctggtc	tccttgga	840
aaggttctgt	tgccctgatt	tatgaactct	ataatagagt	atatagggtt	tgtacctttt	900
ttacaggaag	gtgactttct	gtaacaatgc	gatgtatatt	aaacttttta	taaaagttaa	960

cattttgcat aataaacgat tt

982

&lt;210&gt; 15

&lt;211&gt; 3915

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 15

gtgggggtggg gtgggggtgg gggcttgctg ccccttcagg ctccaccctt tgcggagatt	60
ataaatagtc atgatcccag cgagaccag agatgcctgt aatgggtgaga ctttggatcc	120
ttcctgagga cgtggagaaa actttctgct gagaaggaca ttttgaaggt tttgttggt	180
gaaaaagctg tttctggaat caccctaga tctttcttga agacttgaat tagattacag	240
cgatggggac acagaaggct accccagctc tgatatttgc catcacagtt gctacaatcg	300
gctctttcca atttggctac aacactgggg tcatcaatgc tcctgagaag atcataaagg	360
aatttatcaa taaaactttg acggacaagg gaaatgcccc accctctgag gtgctgctca	420
cgtctctctg gtccttgtct gtggccatat tttccgtcgg gggatatgatc ggctcctttt	480
ccgtcggact ctctgtcaac cgctttggca ggcgcaattc aatgctgatt gtcaacctgt	540
tggctgtcac tgggtggctgc tttatgggac tgtgtaaagt agctaagtcg gttgaaatgc	600
tgatcctggg tcgcttggtt attggcctct tctgcccact ctgcacaggt tttgtgccc	660
tgtacattgg agagatctcg cctactgccc tgcgggggtgc ctttggcact ctcaaccagc	720
tgggcacgtg tgttggaatt ctgggtggccc agatcctttgg tctggaattc atccttgggt	780
ctgaagagct atggccgctg ctactgggtt ttaccatcct tcctgctatc ctacaaagtg	840
cagcccttcc attttgccct gaaagtccca gatttttgct cattaacaga aaagaagagg	900
agaatgctaa gcagatcctc cagcgggtgt ggggcaccca ggatgtatcc caagacatcc	960
aggagatgaa agatgagagt gcaaggatgt cacaagaaaa gcaagtcacc gtgctagagc	1020
tcttttagagt gtccagctac cgacagccca tcatcatctc cattgtgctc cagctctctc	1080
agcagctctc tgggatcaat gctgtgttct attactcaac aggaatcttc aaggatgcag	1140
gtgttcaaga gcccatctat gccaccatcg gcgcgggtgt ggttaatact atcttcaactg	1200
tagtttctct atttctgggtg gaaagggcag gaagaaggac tctgcatatg ataggccttg	1260
gagggatggc tttttgttcc acgctcatga ctgtttcttt gttattaaag gataactata	1320

atgggatgag	ctttgtctgt	attgggggcta	tcttgggtctt	tgtagccttc	tttgaaattg	1380
gaccaggccc	cattccctgg	tttattgttg	ccgaactctt	cagccagggc	ccccgccag	1440
ctgcgatggc	agtggccggc	tgctccaact	ggacctccaa	cttcctagtc	ggattgctct	1500
ccccctccgc	tgctcactat	ttaggagcct	acgtttttat	tatcttcacc	ggcttcctca	1560
ttaccttctt	ggctttttacc	ttcttcaaag	tccctgagac	ccgtggcagg	acttttgagg	1620
atatcacacg	ggcctttgaa	gggcaggcac	acggtgcaga	tagatctgga	aaggacggcg	1680
tcatggagat	gaacagcatc	gagcctgcta	aggagaccac	caccaatgtc	taagtcgtgc	1740
ctccttccac	ctccctcccg	gcatgggaaa	gccacctctc	cctcaacaag	ggagagacct	1800
catcaggatg	aaccaggac	gcttctgaat	gctgctactt	aattcctttc	tcatcccacg	1860
cactccatga	gcaccccaag	gctgcggttt	gttggatctt	caatggcttt	ttaaatttta	1920
tttcttgga	atcctcttct	gcttaggaga	gaccgagtga	acctaccttc	atttcaggag	1980
ggattggccg	cttggcacat	gacaactttg	ccagcttttc	ctcccttggg	ttctgatatt	2040
gccgcactag	gggatatagg	agaggaaaag	taagggtgcag	ttcccccaac	ctcagactta	2100
ccaggaagca	gatacatatg	agtgtggaag	ccggaggggtg	tttatgtaag	agcaccttcc	2160
tcacttccat	acagctctac	gtggcaaat	aacttgagtt	ttattttatt	tatcctctgg	2220
tttaattaca	taattttttt	ttttttactt	taagtttcag	gatacatgtg	ccgaatgtgc	2280
aggtttgtta	cataggtata	tatatgccat	gatggaaata	tttatttttt	taagcgtaat	2340
tttgccaaat	aataaaaaaca	gaaggaaatt	gagattagag	ggaggtggtt	aaagagaggt	2400
tatagagtag	aagatttgat	gctggagagg	ttaagggtgca	ataagaattt	agggagaaat	2460
gttgttcatt	attggagggt	aatgatgtg	gtgcctgagg	tctgtacgtt	acctcttaac	2520
aatttctgtc	cttcagatgg	aaactcttta	acttctcgta	aaagtcatat	acctatataa	2580
taaagctact	gatttccttg	gagctttttt	ctttaagata	atagtttaca	tgtagtagta	2640
cttgaaatct	aggattatta	actaatatgg	gcattgtagt	taatgatgg	tgatgggttc	2700
taattttgga	tggagtccag	ggaagagaaa	gtgatttcta	gaaagcctgt	ccccctcact	2760
ggatgaaata	actccttctt	gtagtagtct	cattactttt	gaagtaatcc	cgccacctat	2820
ctcgtgggag	agccatccaa	ataagaaacc	taaaataatt	ggttcttggg	agagattcat	2880
tatttttcca	ctttgttctt	taggagattt	taggtgttga	ttttctgttg	tatttttaact	2940
cataccttta	aaggaattcc	ccaaagaatg	tttatagcaa	acttggaatt	tgtaacctca	3000
gctctgggag	aggatttttt	tctgagcgat	tattatctaa	agtgtgttgt	tgcttttaggc	3060
tcacggcacg	cttgcgatg	tctgttacca	tgtcactgtg	gtcctatgcc	gaatgccttc	3120

```

aggggacttg aatctttcca ataaaccagg tttagacagt atgagtcaat gtgcagtgta 3180
gccacacatt gagaggatga atgtatgtgc actgtcactt tgctctgggt ggaagtacgt 3240
tattgttgac ttattttctc tgtgtttgtt cctacagccc ctttttcata tgttgctcag 3300
tctccctttc ccttcttggg gcttacacat ctcagaccct ttagccaaac ccttgctcagt 3360
gacagtattt tgggttcttag ttctcactgt tccctctgct cctggagcct ttgaataaaa 3420
atgcacgtag ctgaggccgg atgcggtggc tcacgcctgt aatcccagca ctttgggagg 3480
cctaggcggg cgggtcagggg ttcgagacca gtctggccaa catcgtgaaa ccctgtctct 3540
actaaaaatg caaaaattag cggggcgtgg tggcgggcgc ctgtaatccc agctacttgg 3600
gaagctgagg cgggagaatc atgtgaaccg gggacgcagg ggttgcagtg agcggagatc 3660
gcatcattgc actctagcct gggccacagg gcgagactcc gtctcaaaaa aaaaaaaatg 3720
cacatagcta tcgagtgtgc tttagcttga aaaggtgacc ttgcaacttc atgtcaactt 3780
tctggctcct caaacagtag gttggcagta aggcagggtc ccatttctca ctgagaagat 3840
tgtgaatatt tccatatgga ttttctattg ttactctggg tctttgtttt aaaataaaaa 3900
ttctgaatgt acacg 3915

```

&lt;210&gt; 16

&lt;211&gt; 1670

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 16

```

cggcacgagg caagtgacgc cgagggcctg agtgctccag tagccaccgc atctggagaa 60
ccagcgggta ccatggaggg gatcagtata tacacttcag ataactacac cgaggaaatg 120
ggctcagggg actatgactc catgaaggaa ccctgtttcc gtgaagaaaa tgctaatttc 180
aataaaatct tcctgccac catctactcc atcatcttct taactggcat tgtgggcaat 240
ggattggtca tcctgggtcat gggttaccag aagaaactga gaagcatgac ggacaagtac 300
aggctgcacc tgtcagtggc cgaectctctc tttgtcatca cgcttccctt ctgggcagtt 360
gatgccgtgg caaactggta ctttgggaac ttctatgca aggcagtcca tgatcatctac 420
acagtcaacc tctacagcag tgtcctcatc ctggccttca tcagtctgga ccgtacctg 480
gccatcgtec acgccaccaa cagtcagagg ccaaggaagc tgttggctga aaaggtggtc 540

```



tatgttggcg	tctggatccc	tgcctctctg	ctgactattc	ccgacttcat	ctttgccaac	600
gtcagtgagg	cagatgacag	atatatctgt	gaccgcttct	accccaatga	cttgtgggtg	660
gttgtgttcc	agtttcagca	catcatgggt	ggccttatcc	tgcctgggtat	tgtcatcctg	720
tcctgctatt	gcattatcat	ctccaagctg	tcacactcca	agggccacca	gaagcgcaag	780
gccctcaaga	ccacagtcac	cctcatcctg	gctttctctg	cctgttgggt	gccttactac	840
attgggatca	gcacgactc	cttcacctc	ctggaaatca	tcaagcaagg	gtgtgagttt	900
gagaacactg	tgcacaagt	gatttccatc	accgaggccc	tagctttctt	ccactgttgt	960
ctgaacccca	tcctctatgc	tttccttgga	gccaaattta	aaacctctgc	ccagcacgca	1020
ctcacctctg	tgagcagagg	gtccagcctc	aagatcctct	ccaaaggaaa	gcgagggtgga	1080
cattcatctg	tttccactga	gtctgagtct	tcaagttttc	actccagcta	acacagatgt	1140
aagagacttt	tttttatacg	ataaataact	tttttttaag	ttacacattt	ttcagatata	1200
aaagactgac	caatattgta	cagtttttat	tgcttgttgg	atttttgctc	ttgtgtttct	1260
ttagtttttc	gtgaagggtt	aattgactta	tttatataaa	ttttttttgt	ttcatattga	1320
tgtgtgtcta	ggcaggacct	gtggccaagt	tcttagttgc	tgtatgtctc	gtggtaggac	1380
tgtagaaaag	ggaactgaac	attccagagc	gtgtagtgaa	tcacgtaaag	ctagaaatga	1440
tccccagctg	tttatgcata	gataatctct	ccattcccgt	ggaacgtttt	tcctgttctt	1500
aagacgtgat	tttgctgtag	aagatggcac	ttataaccaa	agcccaaagt	ggatatagaaa	1560
tgtgtgtttt	tcagttttca	ggagtgggtt	gatttcagca	cctacagtgt	acagtcttgt	1620
attaagttgt	taataaaaagt	acatgttaaa	cttaaaaaaa	aaaaaaaaaa		1670

&lt;210&gt; 17

&lt;211&gt; 2360

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 17

gctacgcggg	ccacgctgct	ggctggcctg	acctaggcgc	gcggggctcg	gcggccgcgc	60
gggctgggctg	agtgagcaag	acaagacact	caagaagagc	gagctgcgcc	tgggtcccgg	120
ccaggcttgc	acgcagaggc	gggcggcaga	cggtgcccgg	cggaatctcc	tgagctccgc	180
cgcccagctc	tgggtgccagc	gcccagtggc	cgccgcttcg	aaagtgactg	gtgcctcgcc	240
gcctcctctc	ggtgcgggac	catgaagctg	ctgccgtcgg	tgggtgctgaa	gctctttctg	300

gctgcagttc	tctcggcact	ggtgactggc	gagagcctgg	agcggccttcg	gagagggcta	360
gctgctggaa	ccagcaaccc	ggaccctccc	actgtatcca	cggaccagct	gctaccacct	420
ggagggcgcc	gggaccggaa	agtccgtgac	ttgcaagagg	cagatctgga	ccttttgaga	480
gtcactttat	cctccaagcc	acaagcactg	gccacaccaa	acaaggagga	gcacgggaaa	540
agaaagaaga	aaggcaaggg	gctaggggaag	aagagggacc	catgtcttcg	gaaatacaag	600
gacttctgca	tccatggaga	atgcaaatat	gtgaaggagc	tccgggctcc	ctcctgcac	660
tgccacccgg	gttaccatgg	agagaggtgt	catgggctga	gcctcccagt	ggaaaatcgc	720
ttatatacct	atgaccacac	aaccatcctg	gccgtgggtg	ctgtgggtgc	gtcatctgtc	780
tgtctgctgg	tcacgtggg	gcttctcatg	tttaggtacc	ataggagagg	aggttatgat	840
gtggaaaatg	aagagaaagt	gaagttgggc	atgactaatt	cccactgaga	gagacttgtg	900
ctcaaggaat	cggctgggga	ctgctacctc	tgagaagaca	caaggtgatt	tcagactgca	960
gaggggaaag	acttccatct	agtcacaaag	actccttcgt	cccagttgc	cgtctaggat	1020
tgggcctccc	ataattgctt	tgccaaaata	ccagagcctt	caagtgccaa	acagagtatg	1080
tccgatggta	tctgggtaag	aagaaagcaa	aagcaaggga	ccttcatgcc	cttctgatcc	1140
ccctccacca	aacccactt	cccctcataa	gtttgtttta	acacttatct	tctggattag	1200
aatgccgggt	aaattccata	tgctccagga	tctttgactg	aaaaaaaaa	agaagaagaa	1260
gaaggagagc	aagaaggaaa	gatttgtgaa	ctggaagaaa	gcaacaaaga	ttgagaagcc	1320
atgtactcaa	gtaccaccaa	gggatctgcc	attgggaccc	tccagtgcct	gatttgatga	1380
gttaactgtg	aaataccaca	agcctgagaa	ctgaattttg	ggacttctac	ccagatggaa	1440
aaataacaac	tatttttgtt	gttggtgttt	gtaaatgcct	cttaaattat	atatttat	1500
tattctatgt	atgttaattt	atttagtttt	taacaatcta	acaataatat	ttcaagtgcc	1560
tagactgtta	ctttggcaat	ttcctggccc	tccactcctc	atccccacaa	tctggccttag	1620
tgccaccac	ctttgccaca	aagctaggat	ggttctgtga	cccatctgta	gtaatttatt	1680
gtctgtctac	atctctgcag	atcttccgtg	gtcagagtgc	cactgcggga	gctctgtatg	1740
gtcaggatgt	aggggttaac	ttggtcagag	ccactctatg	agttggactt	cagtcttgcc	1800
taggcgattt	tgtctaccat	ttgtgttttg	aaagcccaag	gtgctgatgt	caaagtgtaa	1860
cagatatcag	tgtctccccg	tgtcctctcc	ctgccaaagtc	tcagaagagg	ttgggcttcc	1920
atgcctgtag	ctttcctggg	ccctcacccc	catggcccca	ggccacagcg	tggggaactca	1980
ctttcccttg	tgtcaagaca	tttctctaac	tcctgccatt	cttctgggtgc	tactccatgc	2040

aggggtcagt gcagcagagg acagtctgga gaaggtatta gcaaagcaaa aggctgagaa	2100
ggaacagggga acattggagc tgactgttct tggtaactga ttacctgcca attgctaccg	2160
agaaggttgg aggtggggaa ggctttgtat aatcccaccc acctcaccaa aacgatgaag	2220
gtatgctgtc atggtccttt ctggaagttt ctggtgccat ttctgaactg ttacaacttg	2280
tatttccaaa cctggttcat atttatactt tgcaatccaa ataaagataa cccttattcc	2340
ataaaaaaaaa aaaaaaaaaa	2360

&lt;210&gt; 18

&lt;211&gt; 6082

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 18

caccttgaca cataggccac ttcctgagcc tggaccagct ctcagagctg ggggaactggc	60
ccaatgcaaa agggtcggga gcatctgcaa cagagactga gctctatcag cttcgggtgac	120
atagcctcca ttcacgtcc ccaactcagc agagagagca caccatcaga cttctaagac	180
ttagtagcca agaagtgttg aattaaactc tctgagacct ctctttagtc tgaccctggc	240
agcctcagtc tcccagagcc tgtgggaact cggcagccga gaggcagaag gctgggcgac	300
gtccggagaa gaagaaacgg ggggaagaact tttctcttac gatctggctt tactctcacg	360
cgcacagccg agtccctggg gaccacagcag aggtccgaag cggagcgggg cggggcgggg	420
ctacggaagc tggcgaggcc gagccctcc tagtggttcc ggaccttgct ccctgaacac	480
tcggaggtgg cgggtggatct tactccttcc agccagtgag gatccagcaa cctgctccgt	540
gcctcccgcg cctgttgggtt ggaagtgaag accttgaaga tcggccgggtt ggaagtgaag	600
accttgaaga tcggcggggc cagcggggcc gagggggcgg gtctggcgct aggtccagcc	660
cctgcgtgcc gggaaaccca gaggaggtcg cagttcagcc cagctgaggc ctgtctgcag	720
aatcgacacc aaccagcatc atgtccatga cactggggta ctgggacatc cgcgggggtga	780
gtgaggggtcc gctgcactgt gggaccgggc gcgtgggcgg gaagtgccga gcggctgggg	840
accggctcta gggacgggtc cctccttagg gctatctctc acaggagggc ctgtgcatgc	900
ctgtgtgtgt gtgtgtgtgt gtgtgtgtgt gtgtgtgtgt gcgtgcgccg gggtgggggg	960
ggggtgcagt gcagtgtaga ctaggggctc acctgggtgca gagaaagtca ccaagtcagg	1020
gaccctccat ctctgacacg acctgcgggc catctcttcc agctggccca cgccatgcgc	1080

ctgctcctgg	aatacacaga	ctcaagctac	gaggaaaaga	agtatacgat	gggggacggg	1140
aatgacaccc	ttgtgtccgg	gctctgccc	ctcacgctga	gttggcacca	agcaacccat	1200
ggtggccacc	tgtggctacc	tctgcaggcc	tcccctgctg	gagctgcagg	ctgtcccttc	1260
cctgagcccc	ggtgaggagg	tcctgtggcc	ttgcaaggca	gaatgctggg	gccccgatgct	1320
gggccccctg	tttaattggg	ttgggtgtcc	ctcagagctt	ccctaaaccc	tggaagcctt	1380
agccgtgtgg	ggccagagc	cctcagcggg	attctttgtc	cctgaaccct	gggatgtggg	1440
actgagtgg	cagattctag	atccacctgt	ctcagggatc	ttgccactgg	ttccttggga	1500
gggtccccgg	gaaggagggc	tgggctctgg	ggaggtttgt	tttcacttct	tcttccccac	1560
ggcagctcct	gactatgaca	gaagccagtg	gctgaatgaa	aaattcaagc	tgggcctgga	1620
ctttcccaat	gtaggtgcag	gggaaggggc	ggttttgggg	gaaagtgcga	cgtgtctctg	1680
actgcattct	ctctccccag	attagagggtg	ttcggatcag	gagtcttctg	cccaattcct	1740
ctcactcctg	gttgtctaca	cagccccctg	atgatgttct	gtgtcccagc	tcatttgttc	1800
atgtgacagt	atttctatgt	caggcctgcc	atgagcgggc	acagtgagtg	cctgggtctcc	1860
cctctgccct	tgcatatggg	aaggggatgc	tggggagcct	gctggcccca	actgagcttc	1920
cccggtttcc	catctatcca	gctgcacct	ttgattgatg	gggctcaca	gatcaccag	1980
agcaacgcca	tcctgtgcta	cattgccccg	aagcacaacc	tgtgtgagtg	tgggttgctg	2040
cagtgtgtgg	ggggaagggtg	gcacacctc	tggctggatt	ggggtgctat	gctcagagtg	2100
agtctgtgtt	ttgtgggtgg	cagggtgggga	gacagaagag	gagaagattc	gtgtggacat	2160
tttggagaac	caggctatgg	acgtctccaa	tcagctggcc	agagtctgct	acagccctga	2220
ctttgtgagt	ccctccctgg	tctggaccag	aagccaggct	tgtattccca	tctactctgg	2280
tcctattcac	aatttgaact	cctcactgct	attgatcctc	tgagggttcc	ctgtactgta	2340
gcagagtgac	tgtcatatca	ttaaacttat	aaaataaaaa	cttgaaatca	gtccccacca	2400
atcataggaa	gtcctatgaa	agctagcaat	tcagttccta	gacaataaag	tcatgcgttc	2460
agaattccct	tcacctctga	cccctgacct	ctgccacggg	ccatctaacc	cagctgggttc	2520
atgccatctg	cctgttcggg	gaatacgcag	actcacacta	tggggtaaag	aagtatgtag	2580
tgggatgtaa	tagcaccctc	ctctgtggaa	ttccgtgcta	ttcttttcc	agtgttctc	2640
cttgacacgc	acagggatga	atgcattttc	ctaacaggca	gctcaggggc	ggccagaggg	2700
cccttgttcc	tctgtctcct	tccatgcagc	ctctcaaaat	ctcatctctc	cagaaaactac	2760
tcaatgtcca	ttgtattatt	ctgccactgc	actaggagga	actttcaact	tctttctctg	2820

agctccttta	gttcctttgta	tccttgattt	tgctcgtgtc	tgggtccaga	gcctgccagg	2880
tactcaggtg	ctcctgggct	gaccagagg	aggggtggtt	ggaggtcagt	ggggacagat	2940
tcagggatag	tgttgcatte	ctctctgcct	tcccatcacc	acaaaagcct	ccagctaccc	3000
atctggagtg	taataaatgc	tggatatgtc	agctgaagcc	agttccagct	gtggggaaga	3060
tggctgcttg	ctcgtggcca	gctggggcca	tacacagccc	tggggaggcc	acatctgtgc	3120
agggagcttg	tgtctgaggg	tgggtgacagc	tgttttctgc	ctcaggagaa	actgaagcca	3180
gaatacttgg	aggaacttcc	tacaatgatg	cagcacttct	cacagtccct	ggggaagagg	3240
ccatggtttg	ttggagacaa	ggtaatgggg	gcattgtgatg	aggacactag	agatttgcca	3300
tacatcctat	gttacagaga	ttccagccca	cacattcttg	gccttctgca	gatcaccttt	3360
gtagatttcc	tgccttatga	tgtccttgac	ctccaccgta	tatttgagcc	caactgcttg	3420
gacgccttcc	caaactctga	ggacttcctc	tcccgctttg	aggtgatgcc	cccacctcc	3480
tttctctttg	atgccccttg	ttccgttacc	tcctttcaga	tgctttccca	gtcctggatc	3540
tgcataaaga	ataacttgca	tttattgagt	gctggcttca	tgccaggaac	cttgcccagc	3600
acattatacc	tatcgtgtgg	aatttgaaat	ttccaacatt	cctacagggt	gacagaatta	3660
tcttgcccat	ttagagataa	gaaaactttg	aatgagaggg	tcagtccttt	gttctgggtc	3720
ccagagccag	tggaggctgt	gctgggctcc	ctgtgagcat	ctggatctat	gggtggcagt	3780
cagggctctc	ccttttgtga	caaaagaaag	aagcctcagg	ctcatccagc	ctggatttca	3840
cagcccaggg	cactttggaa	gaggcagaga	actttaggag	catggatgca	gctggcaata	3900
gtaggactga	cacacggtgg	cattgacgtc	gagtacgaaa	cccacaggca	gtattcatag	3960
ctactcccag	aagctttgca	cgatcagacc	cccacgtggg	gaatcctgag	agccagagct	4020
gtggccagag	ctggattagg	gtacatatgt	gggtgcccct	gttgaaggag	tgtatgttga	4080
agtgcctctg	gctggggcac	tctccttctt	tatctttttt	cctctctttt	ttccctccag	4140
tgttccaagt	gttccccctg	tgagatgagt	agcacactga	ttttactgct	attcaccgac	4200
cttctcctct	gcatgaggca	gggtgtgagg	cacagtggga	gttgcataga	tgactgcccc	4260
atcctggaaa	tgagtgcagt	gagaggcctg	caggcagagc	agcctgtgag	gtgtgtggca	4320
ccacctgggt	accaggcctg	gggcctgccc	ctcactcatg	gggaaccatc	cctcaccctg	4380
gctgaatttg	tttgagagca	gcaaactcta	cttttagtac	agatgtgaga	atctgaggca	4440
ttagtccaac	aagtttttca	gcctagaatt	tgttttcctt	tcccactacc	catcaaggga	4500
tctggttact	cagctagttc	ccatcagctc	tggctgtggg	ctcggctgag	tggccttggg	4560
gttatgtaag	aggtagtggg	aggggaagag	agctgagggc	tgcagcatat	ggtccacctg	4620

ggcttggcct ttgggaatag gcagccctgg ctctctctga atccttagaa attacacggc 4680  
tatttgatcc tggaaagatc gtgcagagca cacctgagtg tcatacagcc tggctctgagg 4740  
tagtgggggtt ggagatgagg tgggttgggg cacagtgggtg tttagctcag gtaccagggtg 4800  
gggaggttta gactttctgc tttaaaagga atgattagag cctggctctgg cgtttctttt 4860  
gctgggccaa cacaccttga ccactttcat ccaggttttg ccaggtcctt ggggtgagatc 4920  
tggtctctct tccaggctgc acagacattt tcagagggtcc cctctgtgtg tgcaaacctt 4980  
ggcaagccag gtgcctccct gtgaaacagg agaattgtgt gtagtcagag agtgacagga 5040  
cctcctgagg gatattggggg aggatgggga tttgacagaa agaggccaga actggagaga 5100  
gacagaacca gtctacgttg cagctctgtc ccccttagta gctatttgag tgtgaggaag 5160  
ttactgaact tctgtttccc acatgagaaa tgggtgataat agattcagcc ttgcagagta 5220  
gtcgagtggg ttttctaagc ttatgttgta atttctcttg ggtacagagc acccagcacc 5280  
gtgtagaatc ttcataagtg ttagctgtta ctgtggtaca acattactta aaggaagttg 5340  
gaagagttaa ctccgcaaat ctggggaccc taagaggctg tgtgatgcct cagcacttga 5400  
gcccacgtgg aaaggctgtg gccagggccc tgacctgctg tgtctgcagt ggggttgtcc 5460  
cagccctcat gggcagctga ccttgagttc tggccttatt ttccccctc tcagggtttg 5520  
gagaagatct ctgcctacat gaagtcagc cgcttcctcc caaaacctct gtacacaagg 5580  
gtggctgtct ggggcaacaa gtaatgcctt gaaggccagg aggtgggagt gaggagccca 5640  
tactcagcct gctgccagg ctgtgcagcg cagctggact ctgcatccca gcacctgcct 5700  
cctcgttcc tttctctgtt tattcccatc tttaccccca agactttatt gggcctcttc 5760  
acttccccta aacctctgtc ccatgcaggc cctttgaagc ctcagctacc cactttcctt 5820  
catgaacatc cccctcccaa cactaccctt cctgcacta aagccagcct gaccttcctt 5880  
cctgttagtg gttgtatctg ctttgaaggg cctacctggc ccctcgctg tggagctcag 5940  
ccctgagctg tccccgtgtt gcatgacagc attgactggg ttacaggccc tgctcctgca 6000  
gcatggcccc tgccttaggc ctacctgac aaaataaagc ctcagccaca tttgctatag 6060  
tcttgtctta tttgctcctg gc 6082

<210> 19

<211> 5102

<212> DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 19

gcttctgcga	ctccagttgt	gagagccgca	agggcatggg	aattgacgcc	actcaccgac	60
ccccagtctc	aatctcaacg	ctgtgaggaa	acctcgactt	tgccaggtcc	ccaagggcag	120
cggggctcgg	cgagcgaggc	acccttctcc	gtcccatccc	caatccaagc	gctcctggca	180
ctgacgacgc	caagagactc	gagtgggagt	taaagcttcc	agtgagggca	gcaggtgtcc	240
aggccggggc	tgcggttcc	tggtgacgtc	ttgccctagg	caaagggtccc	agttccttct	300
cggagccggc	tgtcccgcg	cactggaaac	cgcacctccc	cgcagcatgg	gcaccagcct	360
cagcccgaac	gacccttggc	cgctaaaccc	gctgtccatc	cagcagacca	cgctcctgct	420
actcctgtcg	gtgtgggcca	ctgtgcatgt	gggccagcgg	ctgttgaggc	aacggaggcg	480
gcagctccgg	tccgcgcccc	cgggcccggt	tgcgtggcca	ctgatcggaa	acgcggcggc	540
ggtggggccag	gcggctcacc	tctcgttcgc	tcgcctggcg	cggcgctacg	gcgacgtttt	600
ccagatccgc	ctgggcagct	gccccatagt	ggtgtgtaat	ggcgagcgcg	ccatccacca	660
ggccctggtg	cagcagggtc	cggccttcgc	cgaccggccg	gccttcgcct	ccttcctgtg	720
ggtgtccggc	ggccgcagca	tggctttcgg	ccactactcg	gagcactgga	aggtgcagcg	780
gcgcgcagcc	cacagcatga	tgcgcaactt	cttcacgcgc	cagccgcgca	gccgccaaat	840
cctcgagggc	cacgtgctga	gcgaggcgcg	cgagctgggt	gcgctgctgg	tgcgcggcag	900
cgcggacggc	gccttcctcg	acccgaggcc	gctgaccgtc	gtggccgtgg	ccaacgtcat	960
gagtgccgtg	tgtttcggct	gccgctacag	ccacgacgac	cccaggttcc	gtgagctgct	1020
cagccacaac	gaagagtctg	ggcgcacggc	gggcgcgggc	agcctggtgg	acgtgatgcc	1080
ctggctgcag	tacttcccca	acccggtgcg	caccgttttc	cgcgaattcg	agcagctcaa	1140
ccgcaacttc	agcaacttca	tcctggacaa	gttcttgagg	cactgcgaaa	gccttcggcc	1200
cggggccgcc	ccccgcgaca	tgatggacgc	ctttatcctc	tctgcggaaa	agaaggcggc	1260
cggggactcg	cacggtggtg	gcgcgcggct	ggatttggag	aacgtaccgg	ccactatcac	1320
tgacatcttc	ggcgccagcc	aggacaccct	gtccaccgcg	ctgcagtggc	tgctcctcct	1380
cttcaccagg	tatcctgatg	tgacagactcg	agtgcaggca	gaattggatc	aggtcgtggg	1440
gagggaccgt	ctgccttgta	tgggtgacca	gcccacacct	ccctatgtcc	tggccttcct	1500
ttatgaagcc	atgcgcttct	ccagctttgt	gcctgtcact	attcctcatg	ccaccactgc	1560
caacacctct	gtcttgggct	accacattcc	caaggacact	gtgggtttttg	tcaaccagtg	1620

gtctgtgaat catgacccag tgaagtggcc taacccggag aactttgatc cagctcgatt	1680
cttggacaag gatggcctca tcaacaagga cctgaccagc agagtgatga ttttttcagt	1740
gggcaaaagg cgggtgcattg gcgaagaact ttctaagatg cagctttttc tcttcatctc	1800
catcctggct caccagtgcg atttcagggc caacccaaat gagcctgcga aaatgaattt	1860
cagttatggc ctaaccatta aacccaagtc atttaaagtc aatgtcactc tcagagagtc	1920
catggagctc cttgatagtg ctgtccaaaa tttacaagcc aaggaaactt gccataaga	1980
agcaagaggc aagctgaaat tttagaaata ttcacatctt cggagatgag gagtaaaatt	2040
cagttttttt ccagttcctc ttttgtgctg cttctcaatt agcgtttaag gtgagcataa	2100
atcaactgtc catcagggtga ggtgtgctcc ataccagcg gttcttcatg agtagtgggc	2160
tatgcaggag cttctgggag atttttttga gtcaaagact taaagggccc aatgaattat	2220
tatatacata ctgcatcttg gttatttctg aaggtagcat tctttggagt taaaatgcac	2280
atatagacac atacacccaa acacttacac caaactactg aatgaagaag tatttttggt	2340
accaggccat ttttggtggg aatccaagat tggctctcca tatgcagaaa tagacaaaaa	2400
gtatattaaa caaagtttca gagtatattg ttgaagagac agagacaagt aatttcagt	2460
taaagtgtgt gattgaaggt gataagggaa aagataaaga ccagaaattc ctttttcacc	2520
ttttcaggaa aataacttag actctagtat ttatgggtgg atttatectt ttgccttctg	2580
gtatacttcc ttacttttaa ggataaatca taaagtcagt tgctcaaaaa gaaatcaata	2640
gttgaattag tgagtatagt ggggttccat gagttatcat gaattttaaa gtatgcatta	2700
ttaaattgta aaactccaag gtgatgttgt acctcttttg cttgccaag tacagaattt	2760
gaattatcag caaagaaaaa aaaaaagcc agccaagctt taaattatgt gaccataatg	2820
tactgatttc agtaagtctc ataggttaaa aaaaaagtc accaaatagt gtgaaatata	2880
ttacttaact gtccgtaagc agtatattag tattatcttg ttcaggaaaa ggttgaataa	2940
tatatgcctt gtgtaatatt gaaaattgaa aagtacaact aacgcaacca agtgtgctaa	3000
aaatgagctt gattaaatca accacctatt ttgacatgg aaatgaagca gggtttcttt	3060
tcttcaactc aattttggcg aatctcaaaa ttagatccta agatgtgttc ttatttttat	3120
aacatcttta ttgaaattct atttataata cagaatcttg ttttgaaaat aacctaat	3180
atatattaaa attccaaatt catggcatgc tttaaatttta actaaatttt aaagccattc	3240
tgattattga gttccagttg aagttagtgg aaatctgaac attctcctgt ggaaggcaga	3300
gaaatctaag ctgtgtctgc ccaatgaata atggaaaatg ccatgaatta cctggatgtt	3360
ctttttacga ggtgacaaga gttggggaca gaactcccat tacaactgac caagtttctc	3420



ttctagatga	ttttttgaaa	gttaacatta	atgcctgctt	tttggaagt	cagaatcaga	3480
agatagtctt	ggaagctgtt	tggaaaagac	agtggagatg	aggtcagttg	tgttttttaa	3540
gatggcaatt	actttggtag	ctgggaaagc	ataaagctca	aatgaaatgt	atgcattcac	3600
atttagaaaa	gtgaattgaa	gtttcaagtt	ttaaagttca	ttgcaattaa	acttccaaag	3660
aaagtctctac	agtgtcctaa	gtgctaagtg	cttattacat	tttattaagc	tttttggaat	3720
ctttgtacca	aaatttttaa	aaagggagtt	tttgatagtt	gtgtgtatgt	gtgtgtgggg	3780
tgggggggatg	gtaagagaaa	agagagaaac	actgaaaaga	aggaaagatg	gttaaacatt	3840
ttcccactca	ttctgaatta	attaatttgg	agcacaaaat	tcaaagcatg	gacatttaga	3900
agaaagatgt	ttggcgtagc	agagttaaatt	ctcaaataagg	ctattaaaaa	agtctacaac	3960
atagcagatc	tgttttgtgg	tttggaatat	taaaaaactt	catgtaattt	tatttttaaaa	4020
tttcatagct	gtacttcttg	aatataaaaa	atcatgccag	tatttttaaa	ggcattagag	4080
tcaactacac	aaagcaggct	tgcccagtac	atttaaattt	tttggcactt	gccattccaa	4140
aatattatgc	cccaccaagg	ctgagacagt	gaatttgggc	tgctgtagcc	tattttttta	4200
gattgagaaa	tgtgtagctg	caaaaataat	catgaaccaa	tctggatgcc	tcattatgtc	4260
aaccagggtcc	agatgtgcta	taatctgttt	ttacgtatgt	aggcccagtc	gtcatcagat	4320
gcttgcgcca	aaagaaagct	gtgtttatat	ggaagaaagt	aagggtgctt	gagtttacct	4380
ggcttattta	atatgcttat	aacctagtta	aagaaaggaa	aagaaaacaa	aaaacgaatg	4440
aaaataactg	aatttggagg	ctggagtaat	cagattactg	ctttaatcag	aaaccctcat	4500
tgtgtttcta	ccggagagag	aatgtatttg	ctgacaacca	ttaaagtcag	aagttttact	4560
ccaggttatt	gcaataaagt	ataatgttta	ttaaattgctt	cattttgtatg	tcaaagcttt	4620
gactctataa	gcaaattgct	tttttccaaa	acaaaaagat	gtctcagggt	tgttttgtga	4680
attttctaaa	agctttcatg	tcccagaact	tagcctttac	ctgtgaagtg	ttactacagc	4740
cttaatatatt	tcctagtaga	tctatattag	atcaaatagt	tgcatagcag	tatatgttaa	4800
tttgtgtgtt	tttagctgtg	acacaactgt	gtgattaaaa	ggtatacttt	agtagacatt	4860
tataactcaa	ggataccttc	ttatttaatc	ttttcttatt	tttgtacttt	atcatgaatg	4920
cttttagtgt	gtgcataata	gctacagtgc	atagttgtag	acaaagtaca	ttctggggaa	4980
acaacattta	tatgtagcct	ttactgtttg	atataccaaa	ttaaaaaaaaa	attgtatctc	5040
attacttata	ctgggacacc	attaccaaaa	taataaaaat	cactttcata	atcttgaaaa	5100
aa						5102

&lt;210&gt; 20

&lt;211&gt; 1524

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 20

```

gcagagcaca gcatcgtcgg gaccagactc gtctcaggcc agttgcagcc ttctcagcca      60
aacgccgacc aaggaaaact cactaccatg agaattgcag tgatttgctt ttgcctccta      120
ggcatcacct gtgccatacc agttaaacag gctgattctg gaagttctga ggaaaagcag      180
ctttacaaca aatacccaga tgctgtggcc acatggctaa accctgacct atctcagaag      240
cagaatctcc tagccccaca gacccttcca agtaagtcca acgaaagcca tgaccacatg      300
gatgatatgg atgatgaaga tgatgatgac catgtggaca gccaggactc cattgactcg      360
aacgactctg atgatgtaga tgacactgat gattctcacc agtctgatga gtctcaccat      420
tctgatgaat ctgatgaact ggtcactgat tttcccacgg acctgccagc aaccgaagtt      480
ttcactccag ttgtccccac agtagacaca tatgatggcc gaggtgatag tgtgggttat      540
ggactgaggt caaaatctaa gaagtttcgc agacctgaca tccagtacct tgatgctaca      600
gacgaggaca tcacctcaca catggaaagc gaggagttga atggtgcata caaggccatc      660
cccgttgccc aggacctgaa cgcgccttct gattgggaca gccgtgggaa ggacagttat      720
gaaacgagtc agctggatga ccagagtgtc gaaaccacaca gccacaagca gtccagatta      780
tataagcgga aagccaatga tgagagcaat gagcattccg atgtgattga tagtcaggaa      840
ctttccaaag tcagccgtga attccacagc catgaatttc acagccatga agatatgctg      900
gttgtagacc caaaagtaa ggaagaagat aaacacctga aatttcgtat ttctcatgaa      960
ttagatagtg catcttctga ggtcaattaa aaggagaaaa aatacaattt ctcaactttgc     1020
atttagtcaa aagaaaaaat gctttatagc aaaatgaaag agaacatgaa atgcttcttt     1080
ctcagtttat tggttgaatg tgtatctatt tgagtctgga aataactaat gtgtttgata     1140
attagtttag tttgtggctt catggaaact ccctgtaaac taaaagcttc agggttatgt     1200
ctatgttcat tctatagaag aaatgcaaac tatcactgta ttttaatatatt tgttattctc     1260
tcatgaatag aaatttatgt agaagcaaac aaaatacttt taccactta aaaagagaat     1320
ataacatttt atgtcactat aatcttttgt tttttaagtt agtgtatatt ttgttgtgat     1380
tatctttttg tgggtgtgaat aaatctttta tcttgaatgt aataagaatt tgggtggtgtc     1440

```

aattgcttat ttgttttccc acggttgtcc agcaattaat aaaacataac cttttttact 1500  
gcctaataaaaa aaaaaaaaaa aaaa 1524

<210> 21

<211> 478

<212> DNA

<213> Homo sapiens

<400> 21  
cttcaagacc attgtggcca aggagatctg tgctgacccc aagcagaagt gggttcagga 60  
ttccatggac cacctggaca agcaaaccga aactccgaag acttgaacac ccactccaca 120  
acccaagaat ctgcagctaa cttattttcc cctagctttc ccagacacc ttgttttatt 180  
ttattataat gaatttggtt gttgatgtga aacattatgc ctttaagtaat gttaattctt 240  
atttaagtta ttgatgtttt aagtatcttt catggtacta gtgtttttta gatacagaga 300  
cttgggggaaa ttgcttttcc tcttttgaac cacagttcta cccctgggat gttttgaggg 360  
tctttgcaag aatcattaat acaaagaatt ttttttaaca ttccaatgca ttgctaaaaat 420  
attattgtgg aaatgaatat tttgtaacta ttacaccaaa taaatatatt tttgtaca 478

<210> 22

<211> 1778

<212> DNA

<213> Homo sapiens

<400> 22  
tagaagttta caatgaagtt tcttctaata ctgctcctgc aggccactgc ttctggagct 60  
cttcccctga acagctctac aagcctggaa aaaaataatg tgctatattg tgagagatac 120  
ttagaaaaat tttatggcct tgagataaac aaacttccag tgacaaaaat gaaatatagt 180  
ggaaacttaa tgaaggaaaa aatccaagaa atgcagcact tcttgggtct gaaagtgacc 240  
gggcaactgg acacatctac cctggagatg atgcacgcac ctcgatgtgg agtccccgat 300  
ctccatcatt tcagggaat gccagggggg cccgtatgga ggaaacatta tatcacctac 360  
agaatcaata attacacacc tgacatgaac cgtgaggatg ttgactacgc aatccggaaa 420

```

gctttccaag tatggagtaa tgttaccccc ttgaaattca gcaagattaa cacaggcatg      480
gctgacathtt tgggtggthtt tgcccgtgga gctcatggag acttccatgc ttttgatggc      540
aaagggtggaa tcctagccca tgcttttgga cctggatctg gcattggagg ggatgcacat      600
ttcgatgagg acgaattctg gactacacat tcaggaggca caaacttggt cctcactgct      660
gttcacgaga ttggccattc cttaggtctt ggccattcta gtgatccaaa ggctgtaatg      720
ttccccacct acaaatatgt cgacatcaac acatttcgcc tctctgctga tgacatacgt      780
ggcattcagt ccctgtatgg agacccaaaa gagaaccaac gcttgccaaa tcctgacaat      840
tcagaaccag ctctctgtga ccccaatttg agttttgatg ctgtcactac cgtgggaaat      900
aagatctttt tcttcaaaga caggttcttc tggctgaagg tttctgagag accaaagacc      960
agtgttaatt taatttcttc cttatggcca accttgccat ctggcattga agctgcttat     1020
gaaattgaag ccagaaatca agtttttctt tttaaagatg acaaatactg gttaattagc     1080
aatttaagac cagagccaaa ttatcccaag agcatacatt cttttggtht tcctaacttt     1140
gtgaaaaaaaa ttgatgcagc tgthtttaac ccacgtthtt ataggaccta cttctttgta     1200
gataaccagt attggaggta tgatgaaagg agacagatga tggaccctgg ttatcccaaa     1260
ctgattacca agaacttcca aggaatcggg cctaaaattg atgcagtctt ctattctaaa     1320
aacaaatact actatttctt ccaaggatct aaccaatttg aatatgactt cctactccaa     1380
cgtatcacca aaacactgaa aagcaatagc tggthttggtt gttagaaatg gtgtaattaa     1440
tggthtttgt tagttcactt cagcttaata agtatthatt gcatathtgc tatgtctca     1500
gtgtaccact acttagagat atgtatcata aaaataaaat ctgtaaacca taggtaatga     1560
ttatataaaa tacataatat ttttcaattt tgaaaactct aattgtccat tcttgcttga     1620
ctctactatt aagthtgaaa atagttacct tcaaagcaag ataattctat ttgaagcatg     1680
ctctgtaagt tgcttcctaa catccttgga ctgagaaatt atacttactt ctggcataac     1740
taaaattaag tatatatatt ttggctcaaa taaaattg                                1778

```

&lt;210&gt; 23

&lt;211&gt; 599

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

```

tttttttttt tgctgttacg attatttatt gttactattc actgcttht tcccatggac      60

```

ccatgggtta gagctccata cataaagtag gtgctcatta aatgttggct ggactgagtc 120  
agccctcca tcttggccct tgagtcctca gaaaagcagt gccatccatg agaagcaact 180  
tgaattgtat tttttattga aaagaattca ggctagagtt gggaggagga tgcaagagct 240  
actgggaagg gggagctcag tctgaacctg ggggatcagg ggagtagggg actctccct 300  
tgtccactga tgggggggtct ggctgttact cctctccctt cagcacagaa agaacttggt 360  
cagtaaaaaat gcctgtgtaa gtgctcatgg ctgctgtgct tttgctgtac aagtccctga 420  
gtttctcatc tacagcgggc aggtatgtct tctcgtacag gttctgggag gctgtctttg 480  
ctgactccca gtaactggag agagattcct tcacctgggt gaggaaggtc gggctaggca 540  
tctcatcttg ctggggctgt tgggtccctg gactcaaact ccaataccag gaggacaag 599

<210> 24

<211> 1270

<212> DNA

<213> Homo sapiens

<400> 24

gcggacgtgg gcaggagggc tggaaaagcc ggcgctggag cggaacggga gtagctgcct 60  
gggcgccaaa ggccgcggca ctcccacgtg gaccccgaag tcccgaacct ggggatgggc 120  
ccgcggctgc gaggggatct tctctggatc aagcaatggt ggtgaaaaat gtttcgcaag 180  
ggcaaaaaac gacacagtag tagcagttcc caaagtagcg aaatcagtac taagagcaag 240  
tctgtggatt ctagccttgg gggcttttca cgatccagca ctgtggccag cctcgacaca 300  
gattccacca aaagctcagg acaaagcaac aataattcag atacctgtgc agaatttcga 360  
ataaaatatg ttggtgccat tgagaaactg aaactctccg agggaaaagg ccttgaaggg 420  
ccattagacc tgataaatta tatagacgtt gccagcaag atggaaagtt gccttttgtt 480  
cctccggagg aagaatttat tatgggagtt tccaagtatg gcataaaagt atcaacatca 540  
gatcaatatg aacaagcaca agccatttgc aaggttttat ccaccgcttt tgactctgta 600  
ttaacatctg agaaacctg aatcctgcaa tcaagtagaa gtcaacttca tctgaaagtt 660  
cagctgtttt caaactgcaa tgctgaaatg ttatgcaaat aatgaagtta tcccttgctc 720  
tagattttct gaagaaaatg gattgtgtaa aatgctgatc atttgtttat taaaatgtgt 780  
cctattacac agtgagttaa ctctcaatga agtcatctat tttctgggct aaaaaacttc 840

```

atttgtcttt ttcaacttct aataagctta acctaagtgt cacgaagacg agatgtcaca      900
gaggteccact cagtgcacaca cactgaaggc ctgagggaag actgaggaca tgggctcagt      960
ggtggctttcc cagtcattggt atcactggca tggacctctg tccggcagag gtgtggactg    1020
gagaccagga ttcattgctgg tctggaacaa tgacattgcc aacttaagac acacaaagca    1080
gattttcaga agtgtctggt caagataaca tgctggccaa ccacaattcc tagagttaag    1140
agaaccttaa aagattaccg ctcatgctaa aagtatgtaa agatcccatg tacagtatga    1200
tagtgtacttt tttttaaagg actgtcaata taaaaaactt taaagattaa aaacattaaa    1260
aataaaaaaaa                                     1270

```

&lt;210&gt; 25

&lt;211&gt; 2390

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 25

```

ccagcctcgg agggaggggat tagaagccgc tagacttttt ttctctccct ctcagtagca      60
cggagtcgga attaattgga tttcattcac tggggaggaa caaaaactat ctgggcagct    120
tcattgagag agattcattg acactaagag ccagcgtgc agctggtgca gagagaacct    180
ccggctttga cttctgtctc gtctgcccc aaggccgtag cctcggcttg ggaaggcgag    240
gcggaattaa accccgctcc gagagcgcac gtctgcgcgc ggtgcgtcgg ccattgcctg    300
ccccgagggg cgtctggtag gcaccccgcc ctctcccgca gctcgacccc catgatagat    360
acgctcagac ccgtgccctt cgcgtcgga atggcgatca gcaagacggt ggcgtggctc    420
aacgagcagc tggagctggg caacgagcgg ctgctgctga tggactgccg gccgcaggag    480
ctatacgagt cgtcgcacat cgagtcggcc atcaacgtgg ccatccccgg catcatgctg    540
cggcgcctgc agaagggtaa cctgccggtg cgcgcgtctt tcacgcgcgg cgaggaccgg    600
gaccgcttca cccggcgctg tggcaccgac acagtgggtc tctacgacga gagcagcagc    660
gactggaacg agaatacggg cggcgagtcg ttgctcgggc tgctgctcaa gaagctcaag    720
gacgagggct gccgggcgtt ctacctgga ggtggcttca gtaagttcca agccgagttc    780
tccctgcatt gcgagaccaa tctagacggc tcgtgtagca gcagctcgcc gccgttgcca    840
gtgctggggc tcgggggcct gcggatcagc tctgactctt cctcggacat cgagtctgac    900
cttgaccgag accccaatag tgcaacagac tcggatggta gtccgctgtc caacagccag    960

```

ccttccttcc cagtggagat cttgcccttc ctctacttgg gctgtgccaa agactccacc 1020  
aacttggacg tgttggagga attcggcatc aagtacatct tgaacgtcac cccaatttg 1080  
ccgaatctct ttgagaacgc aggagagttt aaatacaagc aaatcccatc ctgggatcac 1140  
tgagagccaaa acctgtccca gtttttccct gagggcattt ctttcataga tgaagcccg 1200  
ggcaagaact gtggtgtctt ggtacattgc ttggctggca ttagccgctc agtcaactgtg 1260  
actgtggctt accttatgca gaagctcaat ctgtcgatga acgatgccta tgacattgtc 1320  
aaaatgaaaa aatccaacat atcccctaac ttcaacttca tgggtcagct gctggacttc 1380  
gagaggacgc tgggactcag cagcccatgt gacaacaggg ttccagcaca gcagctgtat 1440  
tttaccaccc cttccaacca gaatgtatac caggtggact ctctgcaatc tacgtgaaag 1500  
acccacaccc cctccttget ggaatgtgtc tggcccttca gcagtttctc ttggcagcat 1560  
cagctgggct gctttctttg tgtgtggccc caggtgtcaa aatgacacca gctgtctgta 1620  
ctagacaagg ttaccaagtg cggaattggt taatactaac agagagattt gctccattct 1680  
ctttggaata acaggacatg ctgtatagat acaggcagta ggtttgctct gtacccatgt 1740  
gtacagccta cccatgcagg gactgggatt cgaggacttc caggcgcata gggtagaacc 1800  
aaatgatagg gtaggagcat gtgttcttta gggccttgta aggetgtttc cttttgcac 1860  
tggaactgac tatataattg tcttcaagtg aagactaatt caattttgca tatagaggag 1920  
ccaaagagag atttcagctc tgtatttgtg gtatcagtgt tggaaaagag aaatctgata 1980  
ctccatttgg attattgtaa atatttgatc ttgaatcact tgacagtgtt tgtttgaatt 2040  
gtgtttgttt tttcctttga tgggcttaaa agaaattatc caaagggaga aagagcagta 2100  
tgccacttct taaaacagaa caaaacaaaa aaagaaaatt gtgctctgtt gtaatccaaa 2160  
ggggagattt gcagcatgct tgactttacc aattctgatg acatctttac ggacactatt 2220  
atcactaaga ccttgttatg gcgaagtctt tagtcttttt catgtatttt cctcatgatt 2280  
ttttctcttt atgtagtttg agtatgcctt acctttgtaa atatttttgc ttgtgttgtc 2340  
gcaaagggga taatctggga aagacaccaa atcatgggct cactttaaaa 2390

<210> 26

<211> 3871

<212> DNA

<213> Homo sapiens

<400> 26  
 gcggattgtt gctccgttgt acctgctggg gaattcacct cgttactgct tgatatcttc . 60  
 ccccccttac aaaatcagaa aagttgtgtt ttctaatacc aaagaggagg tttggctttc 120  
 tgtgggtgat tcccagacac tgaagtgcaa agaagagacc ctccatagaaa agtaaaatat 180  
 gactaaaagc aatggagaag agcccaagat ggggggagcagg atggagagat tccagcaggg 240  
 agtccgtaaa cgcacacttt tggccaagaa gaaagtgcag aacattacaa aggaggatgt 300  
 taaaagttac ctgttttcgga atgcttttgt gctgctcaca gtcaccgctg tcatttgtggg 360  
 tacaatcctt ggattttacc tccgaccata cagaatgagc taccgggaag tcaagtactt 420  
 ctccctttcct ggggaacttc tgatgaggat gttacagatg ctgggtcttac cacttatcat 480  
 ctccagtctt gtcacaggaa tggcggcgct agatagtaag gcatcaggga agatgggaat 540  
 gcgagctgta gtctattata tgactaccac catcattgct gtggtgattg gcataatcat 600  
 tgtcatcatc atccatcctg ggaagggcac aaaggaaaac atgcacagag aaggcaaaat 660  
 tgtacgagtg acagctgcag atgccttcct ggacttgatc aggaacatgt tccctccaaa 720  
 tctggtagaa gcctgcttta aacagtttaa aaccaactat gagaagagaa gcttttaaagt 780  
 gcccatccag gccaacgaaa cgcttggtggg tgctgtgata aacaatgtgt ctgaggccat 840  
 ggagactctt acccgaatca cagaggagct ggtcccagtt ccaggatctg tgaatggagt 900  
 caatgccctg ggtctagttg tcttctccat gtgcttcggg tttgtgattg gaaacatgaa 960  
 ggaacagggg caggccctga gagagtctt tgattctctt aacgaagcca tcatgagact 1020  
 ggtagcagta ataatgtggg atgccccgt gggattcttc ttcttgattg ctgggaagat 1080  
 tgtggagatg gaagacatgg gtgtgattgg ggggagcctt gccatgtaca ccgtgactgt 1140  
 cattgttggc ttactcatte acgcagtcac cgtcttgcca ctctctact tcttggtaac 1200  
 acggaaaaac ccttggggtt ttattggagg gttgctgcaa gcactcatca ccgctctggg 1260  
 gacctcttca agttctgcca ccctacccat caccttcaag tgcctggaag agaacaatgg 1320  
 cgtggacaag cgcgtcacca gattcgtgct ccccgtagga gccaccatta acatggatgg 1380  
 gactgccctc tatgaggctt tggctgccat ttccattgct caagttaaca actttgaact 1440  
 gaacttcgga caaattatta caatcagcat cacagccaca gctgccagta ttggggcagc 1500  
 tggaattcct caggcgggac tggctactat ggtcattgtg ctgacatctg tcggcctgcc 1560  
 cactgacgac atcacgctca tcacgcggg ggactgggtc ctggatcgcc tccggaccac 1620  
 caccaacgta ctgggagact ccctgggagc tgggattgtg gagcacttgt cacgacatga 1680



actgaagaac	agagatgttg	aatgggtaa	ctcagtgatt	gaagagaatg	aatgaagaa	1740
accatatcaa	ctgattgcac	aggacaatga	aactgagaaa	cccatcgaca	gtgaaaccaa	1800
gatgtagact	aacataaaga	aacactttct	tgagcaccag	gtgttaaaaa	ccattataaa	1860
atctttccat	ctcattacag	ctcattcgct	ccagcaagcc	cgtcactctc	cctttcctcc	1920
cttctgataa	gactggaaaa	tagtcctcca	aaacacaagg	gaggattttg	gggtggccaaa	1980
gtgtacaatt	ttcatccac	aattgaaatt	tttaaatacat	ttcatgttag	tcttaccgaa	2040
taaggtagca	agatcacaaa	tagtggtgat	cagatcttac	aagtttatgt	ggcacacaat	2100
tcctataaat	gtgatttttt	tatataagtt	aaagagacaa	atagtaggct	aaaaacattt	2160
taaaatcaac	ttttgaaatt	taaaaatctt	tcagaataca	attcagtttt	agtttcaaaa	2220
tgттаacaac	ttgaattaca	accggttatc	agttggacag	taagatttta	tccctttctc	2280
ttctgactgg	tatacctatt	tcattagtag	ctaggtgcac	atatacatct	agcacagctg	2340
tgaggacaga	cagaaggcaa	agtttccatg	tggccttgag	caagtcccat	ctcacctcta	2400
ggcctcagtg	tcctcatcta	taaaatgagg	gacttcctta	gaagtcttca	tggctctctc	2460
cagcccagac	atcctgtgat	gtcatgaaag	cacctgccct	ctgtttcccc	tcagaacacc	2520
ctgtaccatc	catggagcac	gaggccttca	gaaaagacac	ttcaatggga	gtgaacattt	2580
ctaactaagg	acaggatggc	tgtgtgtggt	ggtcaccagg	tcctgtgagc	aaagtgcagg	2640
ttatgcaagt	cgccaggcag	gaggccattc	caggagtggg	attattcatc	aaactctttg	2700
cccagttcat	cccaatgggg	gaagtattcc	cttctttcct	actctgggaa	gaatgtctcc	2760
tgccactcct	caactgatga	tagacttcga	aaacagatga	gaagactagc	agctagcaag	2820
gggtgcttgta	gtcacactgt	ggaacactaa	agagctagga	aagagttgag	cacaggcaac	2880
attacaaaca	aaggatttga	aaacaccaag	agtacaggtc	ttctttaagg	aagaataaaa	2940
aagaagaggt	tcatttttct	ggcttttttt	ttcacctgaa	acactttttc	tcgagtccaa	3000
aatcattccc	cccgtgaagt	ctgcttacca	aaacataaga	cgacttatat	atttgaaaga	3060
agtcaaatga	atgagctctc	taatagaagt	ccatgagttg	agtgggtatt	tcttatttga	3120
aagtgttttt	ctttaatcaa	aagtccttag	aatgagggaa	acaaaatatt	tatttgtttt	3180
ggaatcccac	ttatcaaact	attcaaaact	ttcagctgga	gtggggtttg	cttttgtttt	3240
gtttgtgtcc	ataagagaaa	tggtagaaga	tgaatcagta	tgaagacact	gtcaatgagg	3300
ttatgagaaa	aaaacagcag	gggcattagt	ttcaggcaag	gcagctccca	ggtttagaga	3360
ttaattttta	ccccctaagg	aatatccagt	caaagacgct	gagtgggagc	tgtcaggcag	3420
tagcagctgt	gtttgagttt	ctggetgaaa	atgggtgaaga	atggacttaa	ttatgctaac	3480

```

aaactgaaaa atctagacat agatcctctg atatacaatt agagatattt ttatatagac 3540
cccaagcatt ctgtgcataa aagttaacat taggctgtgg tgcagtaacc atttaatgtc 3600
gaggctctat ttcggaaata cactacaaat gttaaagtac gtggctgtcc tcttaagaca 3660
ctagtagagc aaagacttaa tcatatcaac ttaattctgt tacacaatat gtgtttttta 3720
atatactaac catttcttat ggaaaggctc tgtggggagc ccatcctctc gccaaagccat 3780
cacaggctct gcatacacat gcactcagtg tggactggga agcattactt tgtagatgta 3840
ttttcaataa agaaaaaaat agttttacat t 3871

```

<210> 27

<211> 1433

<212> DNA

<213> Homo sapiens

<400> 27

```

attcggggcg agggaggagg aagaagcgga ggaggcggct cccgctcgca gggccgtgca 60
cctgcccgcc cgcccgtctg ctgctctgcc cgccgcgccg cgctgccgac cgccagcatg 120
ctgccgagag tgggctgccc cgcgctgccg ctgccgccgc cgccgtgctt gccgctgctg 180
ccgctgctgc tgctgctact gggcgcgagt ggcggcgccg gcggggcgcg cgcgagggtg 240
ctgttccgct gcccgccttg cacacccgag cgcctggccg cctgcgggce cccgcgggtt 300
gcgcgcgccg ccgcgggtggc cgcagtgccc ggaggcgccc gcatgccatg cgcgagctc 360
gtccgggagc cgggctgcgg ctgctgctcg gtgtgcgccc ggctggaggg cgaggcgtgc 420
ggcgtctaca ccccgcgctg cggccagggg ctgcgctgct atccccaccc gggctccgag 480
ctgcccctgc aggcgctggt catgggcgag ggcacttgtg agaagcgccg ggacgccgag 540
tatggcgcca gcccgagca ggttgccagc aatggcgatg accactcaga aggaggcctg 600
gtggagaacc acgtggacag caccatgaac atgttgggag ggggaggcag tgctggccgg 660
aagcccctca agtcgggtat gaaggagctg gccgtgttcc gggagaaggt cactgagcag 720
caccggcaga tgggcaaggg tggcaagcat caccttggcc tggaggagcc caagaagctg 780
cgaccacccc ctgccaggac tccctgcaa caggaactgg accaggtcct ggagcggatc 840
tccaccatgc gccttccgga tgagcggggc cctctggagc acctctactc cctgcacatc 900
cccaactgtg acaagcatgg cctgtacaac ctcaaacagt gcaagatgtc tctgaacggg 960

```

cagcgtgggg agtgctggtg tgtgaacccc aacaccggga agctgatcca gggagccccc 1020  
 accatccggg gggaccccgga gtgtcatctc ttctacaatg agcagcagga ggcttgccggg 1080  
 gtgcacaccc agcgggatgca gtagaccgca gccagccggg gcctggcgcc cctgcccccc 1140  
 gccctctctc aaacaccggc agaaaacgga gagtgcttgg gtggtgggtg ctggaggatt 1200  
 ttccagttct gacacacgta tttatatattg gaaagagacc agcaccgagc tcggcacctc 1260  
 cccggcctct ctcttccag ctgcagatgc cacacctgct ccttcttgct tccccgggg 1320  
 gaggaagggg gttgtggtcg gggagctggg gtacaggttt ggggaggggg aagagaaatt 1380  
 tttatatttg aaccctgtg tcccttttgc ataagattaa aggaaggaaa agt 1433

<210> 28

<211> 1011

<212> DNA

<213> Homo sapiens

<400> 28

ggtttatattt ccagatgcaa tcaatgcccc agtcacctgc tgttataact tcaccaatag 60  
 gaagatctca gtgcagaggc tcgcgagcta tagaagaatc accagcagca agtgtcccaa 120  
 acaagctgtg atgtgagttc agcacaccaa ccttccctgg cctgaagttc ttcttgtgg 180  
 agcaagggac aagcctcata aacctagagt cagagagtgc actatttaac ttaatgtaca 240  
 aaggttccca atgggaaaac tgaggcacca agggaaaaag tgaaccccaa catcactctc 300  
 cacctgggtg cctattcaga acacccaatt tctttagctt gaagtcagga tggctccacc 360  
 tggacaccta taggagcagt ttgccctggg ttccctcctt ccacctgcgt tcctcctcta 420  
 gctcccatgg cagccctttg gtgcagaatg ggctgcactt ctagaccaa actgcaaagg 480  
 aacttcatct aactctgtcc tccctcccca cagcttacag accattgtgg caaggagatc 540  
 tgtgctgacc ccaagcagaa gtgggttcag gattccatgg accacctgga caagcaaacc 600  
 caaactccga agacttgaac actcactcca caacccaaga atctgcagct aacttatattt 660  
 tccctagctt tccccagaca ccttgtttat tttattataa tgaattttgt ttgttgatgt 720  
 gaaacattat gccttaagta atgttaattc ttatttaagt tattgatgtt ttaagtttat 780  
 ctttcatggg actagtgttt tttagataga gagacttggg gaaattgctt ttctcttga 840  
 accacagttc taccctggg atgttttgag ggtctttgca agaatacatta atacaaagaa 900  
 ttttttttaa cattccaatg cattgctaaa atattattgt ggaaatgaat attttgtaac 960

tattacacca aataaatata tttttgtaca aaaaaaaaaa aaaaaaaaaa a 1011

<210> 29

<211> 211

<212> PRT

<213> Homo sapiens

<400> 29

Met Gln Ser Ala Met Phe Leu Ala Val Gln His Asp Cys Arg Pro Met  
1 5 10 15

Asp Lys Ser Ala Gly Ser Gly His Lys Ser Glu Glu Lys Arg Glu Lys  
20 25 30

Met Lys Arg Thr Leu Leu Lys Asp Trp Lys Thr Arg Leu Ser Tyr Phe  
35 40 45

Leu Gln Asn Ser Ser Thr Pro Gly Lys Pro Lys Thr Gly Lys Lys Ser  
50 55 60

Lys Gln Gln Ala Phe Ile Lys Pro Ser Pro Glu Glu Ala Gln Leu Trp  
65 70 75 80

Ser Glu Ala Phe Asp Glu Leu Leu Ala Ser Lys Tyr Gly Leu Ala Ala  
85 90 95

Phe Arg Ala Phe Leu Lys Ser Glu Phe Cys Glu Glu Asn Ile Glu Phe  
100 105 110

Trp Leu Ala Cys Glu Asp Phe Lys Lys Thr Lys Ser Pro Gln Lys Leu  
115 120 125

Ser Ser Lys Ala Arg Lys Ile Tyr Thr Asp Phe Ile Glu Lys Glu Ala  
130 135 140

Pro Lys Glu Ile Asn Ile Asp Phe Gln Thr Lys Thr Leu Ile Ala Gln  
145 150 155 160

Asn Ile Gln Glu Ala Thr Ser Gly Cys Phe Thr Thr Ala Gln Lys Arg  
165 170 175

Val Tyr Ser Leu Met Glu Asn Asn Ser Tyr Pro Arg Phe Leu Glu Ser  
 180 185 190

Glu Phe Tyr Gln Asp Leu Cys Lys Lys Pro Gln Ile Thr Thr Glu Pro  
 195 200 205

His Ala Thr  
 210

<210> 30

<211> 1170

<212> PRT

<213> Homo sapiens

<400> 30

Met Gly Leu Ala Trp Gly Leu Gly Val Leu Phe Leu Met His Val Cys  
 1 5 10 15

Gly Thr Asn Arg Ile Pro Glu Ser Gly Gly Asp Asn Ser Val Phe Asp  
 20 25 30

Ile Phe Glu Leu Thr Gly Ala Ala Arg Lys Gly Ser Gly Arg Arg Leu  
 35 40 45

Val Lys Gly Pro Asp Pro Ser Ser Pro Ala Phe Arg Ile Glu Asp Ala  
 50 55 60

Asn Leu Ile Pro Pro Val Pro Asp Asp Lys Phe Gln Asp Leu Val Asp  
 65 70 75 80

Ala Val Arg Ala Glu Lys Gly Phe Leu Leu Leu Ala Ser Leu Arg Gln  
 85 90 95

Met Lys Lys Thr Arg Gly Thr Leu Leu Ala Leu Glu Arg Lys Asp His  
 100 105 110

Ser Gly Gln Val Phe Ser Val Val Ser Asn Gly Lys Ala Gly Thr Leu  
 115 120 125

Asp Leu Ser Leu Thr Val Gln Gly Lys Gln His Val Val Ser Val Glu  
 130 135 140

Glu Ala Leu Leu Ala Thr Gly Gln Trp Lys Ser Ile Thr Leu Phe Val  
 145 150 155 160

Gln Glu Asp Arg Ala Gln Leu Tyr Ile Asp Cys Glu Lys Met Glu Asn  
 165 170 175

Ala Glu Leu Asp Val Pro Ile Gln Ser Val Phe Thr Arg Asp Leu Ala  
 180 185 190

Ser Ile Ala Arg Leu Arg Ile Ala Lys Gly Gly Val Asn Asp Asn Phe  
 195 200 205

Gln Gly Val Leu Gln Asn Val Arg Phe Val Phe Gly Thr Thr Pro Glu  
 210 215 220

Asp Ile Leu Arg Asn Lys Gly Cys Ser Ser Ser Thr Ser Val Leu Leu  
 225 230 235 240

Thr Leu Asp Asn Asn Val Val Asn Gly Ser Ser Pro Ala Ile Arg Thr  
 245 250 255

Asn Tyr Ile Gly His Lys Thr Lys Asp Leu Gln Ala Ile Cys Gly Ile  
 260 265 270

Ser Cys Asp Glu Leu Ser Ser Met Val Leu Glu Leu Arg Gly Leu Arg  
 275 280 285

Thr Ile Val Thr Thr Leu Gln Asp Ser Ile Arg Lys Val Thr Glu Glu  
 290 295 300

Asn Lys Glu Leu Ala Asn Glu Leu Arg Arg Pro Pro Leu Cys Tyr His  
 305 310 315 320

Asn Gly Val Gln Tyr Arg Asn Asn Glu Glu Trp Thr Val Asp Ser Cys  
 325 330 335

Thr Glu Cys His Cys Gln Asn Ser Val Thr Ile Cys Lys Lys Val Ser  
 340 345 350

Cys Pro Ile Met Pro Cys Ser Asn Ala Thr Val Pro Asp Gly Glu Cys  
 355 360 365

Cys Pro Arg Cys Trp Pro Ser Asp Ser Ala Asp Asp Gly Trp Ser Pro  
 370 375 380

Trp Ser Glu Trp Thr Ser Cys Ser Thr Ser Cys Gly Asn Gly Ile Gln  
 385 390 395 400

Gln Arg Gly Arg Ser Cys Asp Ser Leu Asn Asn Arg Cys Glu Gly Ser  
 405 410 415

Ser Val Gln Thr Arg Thr Cys His Ile Gln Glu Cys Asp Lys Arg Phe  
 420 425 430

Lys Gln Asp Gly Gly Trp Ser His Trp Ser Pro Trp Ser Ser Cys Ser  
 435 440 445

Val Thr Cys Gly Asp Gly Val Ile Thr Arg Ile Arg Leu Cys Asn Ser  
 450 455 460

Pro Ser Pro Gln Met Asn Gly Lys Pro Cys Glu Gly Glu Ala Arg Glu  
 465 470 475 480

Thr Lys Ala Cys Lys Lys Asp Ala Cys Pro Ile Asn Gly Gly Trp Gly  
 485 490 495

Pro Trp Ser Pro Trp Asp Ile Cys Ser Val Thr Cys Gly Gly Gly Val  
 500 505 510

Gln Lys Arg Ser Arg Leu Cys Asn Asn Pro Ala Pro Gln Phe Gly Gly  
 515 520 525

Lys Asp Cys Val Gly Asp Val Thr Glu Asn Gln Ile Cys Asn Lys Gln  
 530 535 540

Asp Cys Pro Ile Asp Gly Cys Leu Ser Asn Pro Cys Phe Ala Gly Val  
 545 550 555 560

Lys Cys Thr Ser Tyr Pro Asp Gly Ser Trp Lys Cys Gly Ala Cys Pro  
 565 570 575

Pro Gly Tyr Ser Gly Asn Gly Ile Gln Cys Thr Asp Val Asp Glu Cys  
 580 585 590

Lys Glu Val Pro Asp Ala Cys Phe Asn His Asn Gly Glu His Arg Cys  
 595 600 605

Glu Asn Thr Asp Pro Gly Tyr Asn Cys Leu Pro Cys Pro Pro Arg Phe  
 610 615 620

Thr Gly Ser Gln Pro Phe Gly Gln Gly Val Glu His Ala Thr Ala Asn  
 625 630 635 640

Lys Gln Val Cys Lys Pro Arg Asn Pro Cys Thr Asp Gly Thr His Asp  
 645 650 655

Cys Asn Lys Asn Ala Lys Cys Asn Tyr Leu Gly His Tyr Ser Asp Pro  
 660 665 670

Met Tyr Arg Cys Glu Cys Lys Pro Gly Tyr Ala Gly Asn Gly Ile Ile  
 675 680 685

Cys Gly Glu Asp Thr Asp Leu Asp Gly Trp Pro Asn Glu Asn Leu Val  
 690 695 700

Cys Val Ala Asn Ala Thr Tyr His Cys Lys Lys Asp Asn Cys Pro Asn  
 705 710 715 720

Leu Pro Asn Ser Gly Gln Glu Asp Tyr Asp Lys Asp Gly Ile Gly Asp  
 725 730 735

Ala Cys Asp Asp Asp Asp Asn Asp Lys Ile Pro Asp Asp Arg Asp  
 740 745 750

Asn Cys Pro Phe His Tyr Asn Pro Ala Gln Tyr Asp Tyr Asp Arg Asp  
 755 760 765

Asp Val Gly Asp Arg Cys Asp Asn Cys Pro Tyr Asn His Asn Pro Asp  
 770 775 780

Gln Ala Asp Thr Asp Asn Asn Gly Glu Gly Asp Ala Cys Ala Ala Asp  
 785 790 795 800

Ile Asp Gly Asp Gly Ile Leu Asn Glu Arg Asp Asn Cys Gln Tyr Val  
 805 810 815

Tyr Asn Val Asp Gln Arg Asp Thr Asp Met Asp Gly Val Gly Asp Gln  
 820 825 830

Cys Asp Asn Cys Pro Leu Glu His Asn Pro Asp Gln Leu Asp Ser Asp  
 835 840 845



Ser Asp Arg Ile Gly Asp Thr Cys Asp Asn Asn Gln Asp Ile Asp Glu  
 850 855 860

Asp Gly His Gln Asn Asn Leu Asp Asn Cys Pro Tyr Val Pro Asn Ala  
 865 870 875 880

Asn Gln Ala Asp His Asp Lys Asp Gly Lys Gly Asp Ala Cys Asp His  
 885 890 895

Asp Asp Asp Asn Asp Gly Ile Pro Asp Asp Lys Asp Asn Cys Arg Leu  
 900 905 910

Val Pro Asn Pro Asp Gln Lys Asp Ser Asp Gly Asp Gly Arg Gly Asp  
 915 920 925

Ala Cys Lys Asp Asp Phe Asp His Asp Ser Val Pro Asp Ile Asp Asp  
 930 935 940

Ile Cys Pro Glu Asn Val Asp Ile Ser Glu Thr Asp Phe Arg Arg Phe  
 945 950 955 960

Gln Met Ile Pro Leu Asp Pro Lys Gly Thr Ser Gln Asn Asp Pro Asn  
 965 970 975

Trp Val Val Arg His Gln Gly Lys Glu Leu Val Gln Thr Val Asn Cys  
 980 985 990

Asp Pro Gly Leu Ala Val Gly Tyr Asp Glu Phe Asn Ala Val Asp Phe  
 995 1000 1005

Ser Gly Thr Phe Phe Ile Asn Thr Glu Arg Asp Asp Asp Tyr Ala  
 1010 1015 1020

Gly Phe Val Phe Gly Tyr Gln Ser Ser Ser Arg Phe Tyr Val Val  
 1025 1030 1035

Met Trp Lys Gln Val Thr Gln Ser Tyr Trp Asp Thr Asn Pro Thr  
 1040 1045 1050

Arg Ala Gln Gly Tyr Ser Gly Leu Ser Val Lys Val Val Asn Ser  
 1055 1060 1065

Thr Thr Gly Pro Gly Glu His Leu Arg Asn Ala Leu Trp His Thr  
 1070 1075 1080

Gly Asn Thr Pro Gly Gln Val Arg Thr Leu Trp His Asp Pro Arg  
 1085 1090 1095

His Ile Gly Trp Lys Asp Phe Thr Ala Tyr Arg Trp Arg Leu Ser  
 1100 1105 1110

His Arg Pro Lys Thr Gly Phe Ile Arg Val Val Met Tyr Glu Gly  
 1115 1120 1125

Lys Lys Ile Met Ala Asp Ser Gly Pro Ile Tyr Asp Lys Thr Tyr  
 1130 1135 1140

Ala Gly Gly Arg Leu Gly Leu Phe Val Phe Ser Gln Glu Met Val  
 1145 1150 1155

Phe Phe Ser Asp Leu Lys Tyr Glu Cys Arg Asp Pro  
 1160 1165 1170

<210> 31

<211> 380

<212> PRT

<213> Homo sapiens

<400> 31

Met Met Phe Ser Gly Phe Asn Ala Asp Tyr Glu Ala Ser Ser Ser Arg  
 1 5 10 15

Cys Ser Ser Ala Ser Pro Ala Gly Asp Ser Leu Ser Tyr Tyr His Ser  
 20 25 30

Pro Ala Asp Ser Phe Ser Ser Met Gly Ser Pro Val Asn Ala Gln Asp  
 35 40 45

Phe Cys Thr Asp Leu Ala Val Ser Ser Ala Asn Phe Ile Pro Thr Val  
 50 55 60

Thr Ala Ile Ser Thr Ser Pro Asp Leu Gln Trp Leu Val Gln Pro Ala  
 65 70 75 80

Leu Val Ser Ser Val Ala Pro Ser Gln Thr Arg Ala Pro His Pro Phe  
 85 90 95

Gly Val Pro Ala Pro Ser Ala Gly Ala Tyr Ser Arg Ala Gly Val Val  
 100 105 110

Lys Thr Met Thr Gly Gly Arg Ala Gln Ser Ile Gly Arg Arg Gly Lys  
 115 120 125

Val Glu Gln Leu Ser Pro Glu Glu Glu Glu Lys Arg Arg Ile Arg Arg  
 130 135 140

Glu Arg Asn Lys Met Ala Ala Ala Lys Cys Arg Asn Arg Arg Arg Glu  
 145 150 155 160

Leu Thr Asp Thr Leu Gln Ala Glu Thr Asp Gln Leu Glu Asp Glu Lys  
 165 170 175

Ser Ala Leu Gln Thr Glu Ile Ala Asn Leu Leu Lys Glu Lys Glu Lys  
 180 185 190

Leu Glu Phe Ile Leu Ala Ala His Arg Pro Ala Cys Lys Ile Pro Asp  
 195 200 205

Asp Leu Gly Phe Pro Glu Glu Met Ser Val Ala Ser Leu Asp Leu Thr  
 210 215 220

Gly Gly Leu Pro Glu Val Ala Thr Pro Glu Ser Glu Glu Ala Phe Thr  
 225 230 235 240

Leu Pro Leu Leu Asn Asp Pro Glu Pro Lys Pro Ser Val Glu Pro Val  
 245 250 255

Lys Ser Ile Ser Ser Met Glu Leu Lys Thr Glu Pro Phe Asp Asp Phe  
 260 265 270

Leu Phe Pro Ala Ser Ser Arg Pro Ser Gly Ser Glu Thr Ala Arg Ser  
 275 280 285

Val Pro Asp Met Asp Leu Ser Gly Ser Phe Tyr Ala Ala Asp Trp Glu  
 290 295 300

Pro Leu His Ser Gly Ser Leu Gly Met Gly Pro Met Ala Thr Glu Leu  
 305 310 315 320

Tyr Thr Ser Ser Phe Val Phe Thr Tyr Pro Glu Ala Asp Ser Phe Pro  
340 345 350

Ser Cys Ala Ala Ala His Arg Lys Gly Ser Ser Ser Asn Glu Pro Ser  
355 360 365

Ser Asp Ser Leu Ser Ser Pro Thr Leu Leu Ala Leu  
370 375 380

<210> 32

<211> 347

<212> PRT

<213> Homo sapiens

<400> 32

Met Cys Thr Lys Met Glu Gln Pro Phe Tyr His Asp Asp Ser Tyr Thr  
1 5 10 15

Ala Thr Gly Tyr Gly Arg Ala Pro Gly Gly Leu Ser Leu His Asp Tyr  
20 25 30

Lys Leu Leu Lys Pro Ser Leu Ala Val Asn Leu Ala Asp Pro Tyr Arg  
35 40 45

Ser Leu Lys Ala Pro Gly Ala Arg Gly Pro Gly Pro Glu Gly Gly Gly  
50 55 60

Gly Gly Ser Tyr Phe Ser Gly Gln Gly Ser Asp Thr Gly Ala Ser Leu  
65 70 75 80

Lys Leu Ala Ser Ser Glu Leu Glu Arg Leu Ile Val Pro Asn Ser Asn  
85 90 95

Gly Val Ile Thr Thr Thr Pro Thr Pro Pro Gly Gln Tyr Phe Tyr Pro  
100 105 110

Arg Gly Gly Gly Ser Gly Gly Gly Ala Gly Gly Ala Gly Gly Gly Val  
 115 120 125

Thr Glu Glu Gln Glu Gly Phe Ala Asp Gly Phe Val Lys Ala Leu Asp  
 130 135 140

Asp Leu His Lys Met Asn His Val Thr Pro Pro Asn Val Ser Leu Gly  
 145 150 155 160

Ala Thr Gly Gly Pro Pro Ala Gly Pro Gly Gly Val Tyr Ala Gly Pro  
 165 170 175

Glu Pro Pro Pro Val Tyr Thr Asn Leu Ser Ser Tyr Ser Pro Ala Ser  
 180 185 190

Ala Ser Ser Gly Gly Ala Gly Ala Ala Val Gly Thr Gly Ser Ser Tyr  
 195 200 205

Pro Thr Thr Thr Ile Ser Tyr Leu Pro His Ala Pro Pro Phe Ala Gly  
 210 215 220

Gly His Pro Ala Gln Leu Gly Leu Gly Arg Gly Ala Ser Thr Phe Lys  
 225 230 235 240

Glu Glu Pro Gln Thr Val Pro Glu Ala Arg Ser Arg Asp Ala Thr Pro  
 245 250 255

Pro Val Ser Pro Ile Asn Met Glu Asp Gln Glu Arg Ile Lys Val Glu  
 260 265 270

Arg Lys Arg Leu Arg Asn Arg Leu Ala Ala Thr Lys Cys Arg Lys Arg  
 275 280 285

Lys Leu Glu Arg Ile Ala Arg Leu Glu Asp Lys Val Lys Thr Leu Lys  
 290 295 300

Ala Glu Asn Ala Gly Leu Ser Ser Thr Ala Gly Leu Leu Arg Glu Gln  
 305 310 315 320

Val Ala Gln Leu Lys Gln Lys Val Met Thr His Val Ser Asn Gly Cys  
 325 330 335

Gln Leu Leu Leu Gly Val Lys Gly His Ala Phe  
 340 345

&lt;210&gt; 33

&lt;211&gt; 196

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 33

Met Pro Gly Met Phe Phe Ser Ala Asn Pro Lys Glu Leu Lys Gly Thr  
 1 5 10 15

Thr His Ser Leu Leu Asp Asp Lys Met Gln Lys Arg Arg Pro Lys Thr  
 20 25 30

Phe Gly Met Asp Met Lys Ala Tyr Leu Arg Ser Met Ile Pro His Leu  
 35 40 45

Glu Ser Gly Met Lys Ser Ser Lys Ser Lys Asp Val Leu Ser Ala Ala  
 50 55 60

Glu Val Met Gln Trp Ser Gln Ser Leu Glu Lys Leu Leu Ala Asn Gln  
 65 70 75 80

Thr Gly Gln Asn Val Phe Gly Ser Phe Leu Lys Ser Glu Phe Ser Glu  
 85 90 95

Glu Asn Ile Glu Phe Trp Leu Ala Cys Glu Asp Tyr Lys Lys Thr Glu  
 100 105 110

Ser Asp Leu Leu Pro Cys Lys Ala Glu Glu Ile Tyr Lys Ala Phe Val  
 115 120 125

His Ser Asp Ala Ala Lys Gln Ile Asn Ile Asp Phe Arg Thr Arg Glu  
 130 135 140

Ser Thr Ala Lys Lys Ile Lys Ala Pro Thr Pro Thr Cys Phe Asp Glu  
 145 150 155 160

Ala Gln Lys Val Ile Tyr Thr Leu Met Glu Lys Asp Ser Tyr Pro Arg  
 165 170 175

Phe Leu Lys Ser Asp Ile Tyr Leu Asn Leu Leu Asn Asp Leu Gln Ala  
 180 185 190

Asn Ser Leu Lys  
195

<210> 34

<211> 367

<212> PRT

<213> Homo sapiens

<400> 34

Met Val Met Glu Val Gly Thr Leu Asp Ala Gly Gly Leu Arg Ala Leu  
1 5 10 15

Leu Gly Glu Arg Ala Ala Gln Cys Leu Leu Leu Asp Cys Arg Ser Phe  
20 25 30

Phe Ala Phe Asn Ala Gly His Ile Ala Gly Ser Val Asn Val Arg Phe  
35 40 45

Ser Thr Ile Val Arg Arg Arg Ala Lys Gly Ala Met Gly Leu Glu His  
50 55 60

Ile Val Pro Asn Ala Glu Leu Arg Gly Arg Leu Leu Ala Gly Ala Tyr  
65 70 75 80

His Ala Val Val Leu Leu Asp Glu Arg Ser Ala Ala Leu Asp Gly Ala  
85 90 95

Lys Arg Asp Gly Thr Leu Ala Leu Ala Ala Gly Ala Leu Cys Arg Glu  
100 105 110

Ala Arg Ala Ala Gln Val Phe Phe Leu Lys Gly Gly Tyr Glu Ala Phe  
115 120 125

Ser Ala Ser Cys Pro Glu Leu Cys Ser Lys Gln Ser Thr Pro Met Gly  
130 135 140

Leu Ser Leu Pro Leu Ser Thr Ser Val Pro Asp Ser Ala Glu Ser Gly  
145 150 155 160

Cys Ser Ser Cys Ser Thr Pro Leu Tyr Asp Gln Gly Gly Pro Val Glu  
                           165                          170                          175  
  
 Ile Leu Pro Phe Leu Tyr Leu Gly Ser Ala Tyr His Ala Ser Arg Lys  
                           180                          185                          190  
  
 Asp Met Leu Asp Ala Leu Gly Ile Thr Ala Leu Ile Asn Val Ser Ala  
                           195                          200                          205  
  
 Asn Cys Pro Asn His Phe Glu Gly His Tyr Gln Tyr Lys Ser Ile Pro  
                           210                          215                          220  
  
 Val Glu Asp Asn His Lys Ala Asp Ile Ser Ser Trp Phe Asn Glu Ala  
   225                          230                          235                          240  
  
 Ile Asp Phe Ile Asp Ser Ile Lys Asn Ala Gly Gly Arg Val Phe Val  
                           245                          250                          255  
  
 His Cys Gln Ala Gly Ile Ser Arg Ser Ala Thr Ile Cys Leu Ala Tyr  
                           260                          265                          270  
  
 Leu Met Arg Thr Asn Arg Val Lys Leu Asp Glu Ala Phe Glu Phe Val  
                           275                          280                          285  
  
 Lys Gln Arg Arg Ser Ile Ile Ser Pro Asn Phe Ser Phe Met Gly Gln  
                           290                          295                          300  
  
 Leu Leu Gln Phe Glu Ser Gln Val Leu Ala Pro His Cys Ser Ala Glu  
   305                          310                          315                          320  
  
 Ala Gly Ser Pro Ala Met Ala Val Leu Asp Arg Gly Thr Ser Thr Thr  
                           325                          330                          335  
  
 Thr Val Phe Asn Phe Pro Val Ser Ile Pro Val His Ser Thr Asn Ser  
                           340                          345                          350  
  
 Ala Leu Ser Tyr Leu Gln Ser Pro Ile Thr Thr Ser Pro Ser Cys  
                           355                          360                          365

&lt;210&gt; 35

&lt;211&gt; 107

&lt;212&gt; PRT



&lt;213&gt; Homo sapiens

&lt;400&gt; 35

Met Ala Arg Ala Thr Leu Ser Ala Ala Pro Ser Asn Pro Arg Leu Leu  
 1 5 10 15

Arg Val Ala Leu Leu Leu Leu Leu Val Ala Ala Ser Arg Arg Ala  
 20 25 30

Ala Gly Ala Pro Leu Ala Thr Glu Leu Arg Cys Gln Cys Leu Gln Thr  
 35 40 45

Leu Gln Gly Ile His Leu Lys Asn Ile Gln Ser Val Lys Val Lys Ser  
 50 55 60

Pro Gly Pro His Cys Ala Gln Thr Glu Val Ile Ala Thr Leu Lys Asn  
 65 70 75 80

Gly Gln Lys Ala Cys Leu Asn Pro Ala Ser Pro Met Val Lys Lys Ile  
 85 90 95

Ile Glu Lys Met Leu Lys Asn Gly Lys Ser Asn  
 100 105

&lt;210&gt; 36

&lt;211&gt; 106

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 36

Met Ala His Ala Thr Leu Ser Ala Ala Pro Ser Asn Pro Arg Leu Leu  
 1 5 10 15

Arg Val Ala Leu Leu Leu Leu Leu Val Gly Ser Arg Arg Ala Ala  
 20 25 30

Gly Ala Ser Val Val Thr Glu Leu Arg Cys Gln Cys Leu Gln Thr Leu  
 35 40 45

Gln Gly Ile His Leu Lys Asn Ile Gln Ser Val Asn Val Arg Ser Pro  
 50 55 60

Gly Pro His Cys Ala Gln Thr Glu Val Ile Ala Thr Leu Lys Asn Gly  
 65 70 75 80

Lys Lys Ala Cys Leu Asn Pro Ala Ser Pro Met Val Gln Lys Ile Ile  
 85 90 95

Glu Lys Ile Leu Asn Lys Gly Ser Thr Asn  
 100 105

<210> 37

<211> 98

<212> PRT

<213> Homo sapiens

<400> 37

Met Asn Gln Thr Ala Ile Leu Ile Cys Cys Leu Ile Phe Leu Thr Leu  
 1 5 10 15

Ser Gly Ile Gln Gly Val Pro Leu Ser Arg Thr Val Arg Cys Thr Cys  
 20 25 30

Ile Ser Ile Ser Asn Gln Pro Val Asn Pro Arg Ser Leu Glu Lys Leu  
 35 40 45

Glu Ile Ile Pro Ala Ser Gln Phe Cys Pro Arg Val Glu Ile Ile Ala  
 50 55 60

Thr Met Lys Lys Lys Gly Glu Lys Arg Cys Leu Asn Pro Glu Ser Lys  
 65 70 75 80

Ala Ile Lys Asn Leu Leu Lys Ala Val Ser Lys Glu Met Ser Lys Arg  
 85 90 95

Ser Pro

&lt;210&gt; 38

&lt;211&gt; 564

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 38

Met Lys Glu His Gly Gly Thr Phe Ser Ser Thr Gly Ile Ser Gly Gly  
 1 5 10 15

Ser Gly Asp Ser Ala Met Asp Ser Leu Gln Pro Leu Gln Pro Asn Tyr  
 20 25 30

Met Pro Val Cys Leu Phe Ala Glu Glu Ser Tyr Gln Lys Leu Ala Met  
 35 40 45

Glu Thr Leu Glu Glu Leu Asp Trp Cys Leu Asp Gln Leu Glu Thr Ile  
 50 55 60

Gln Thr Tyr Arg Ser Val Ser Glu Met Ala Ser Asn Lys Phe Lys Arg  
 65 70 75 80

Met Leu Asn Arg Glu Leu Thr His Leu Ser Glu Met Ser Arg Ser Gly  
 85 90 95

Asn Gln Val Ser Glu Tyr Ile Ser Asn Thr Phe Leu Asp Lys Gln Asn  
 100 105 110

Asp Val Glu Ile Pro Ser Pro Thr Gln Lys Asp Arg Glu Lys Lys Lys  
 115 120 125

Lys Gln Gln Leu Met Thr Gln Ile Ser Gly Val Lys Lys Leu Met His  
 130 135 140

Ser Ser Ser Leu Asn Asn Thr Ser Ile Ser Arg Phe Gly Val Asn Thr  
 145 150 155 160

Glu Asn Glu Asp His Leu Ala Lys Glu Leu Glu Asp Leu Asn Lys Trp  
 165 170 175

Gly Leu Asn Ile Phe Asn Val Ala Gly Tyr Ser His Asn Arg Pro Leu  
 180 185 190

Thr Cys Ile Met Tyr Ala Ile Phe Gln Glu Arg Asp Leu Leu Lys Thr  
 195 200 205

Phe Arg Ile Ser Ser Asp Thr Phe Ile Thr Tyr Met Met Thr Leu Glu  
 210 215 220

Asp His Tyr His Ser Asp Val Ala Tyr His Asn Ser Leu His Ala Ala  
 225 230 235 240

Asp Val Ala Gln Ser Thr His Val Leu Leu Ser Thr Pro Ala Leu Asp  
 245 250 255

Ala Val Phe Thr Asp Leu Glu Ile Leu Ala Ala Ile Phe Ala Ala Ala  
 260 265 270

Ile His Asp Val Asp His Pro Gly Val Ser Asn Gln Phe Leu Ile Asn  
 275 280 285

Thr Asn Ser Glu Leu Ala Leu Met Tyr Asn Asp Glu Ser Val Leu Glu  
 290 295 300

Asn His His Leu Ala Val Gly Phe Lys Leu Leu Gln Glu Glu His Cys  
 305 310 315 320

Asp Ile Phe Met Asn Leu Thr Lys Lys Gln Arg Gln Thr Leu Arg Lys  
 325 330 335

Met Val Ile Asp Met Val Leu Ala Thr Asp Met Ser Lys His Met Ser  
 340 345 350

Leu Leu Ala Asp Leu Lys Thr Met Val Glu Thr Lys Lys Val Thr Ser  
 355 360 365

Ser Gly Val Leu Leu Leu Asp Asn Tyr Thr Asp Arg Ile Gln Val Leu  
 370 375 380

Arg Asn Met Val His Cys Ala Asp Leu Ser Asn Pro Thr Lys Ser Leu  
 385 390 395 400

Glu Leu Tyr Arg Gln Trp Thr Asp Arg Ile Met Glu Glu Phe Phe Gln  
 405 410 415

Gln Gly Asp Lys Glu Arg Glu Arg Gly Met Glu Ile Ser Pro Met Cys  
 420 425 430

Asp Lys His Thr Ala Ser Val Glu Lys Ser Gln Val Gly Phe Ile Asp  
 435 440 445

Tyr Ile Val His Pro Leu Trp Glu Thr Trp Ala Asp Leu Val Gln Pro  
 450 455 460

Asp Ala Gln Asp Ile Leu Asp Thr Leu Glu Asp Asn Arg Asn Trp Tyr  
 465 470 475 480

Gln Ser Met Ile Pro Gln Ser Pro Ser Pro Leu Asp Glu Gln Asn  
 485 490 495

Arg Asp Cys Gln Gly Leu Met Glu Lys Phe Gln Phe Glu Leu Thr Leu  
 500 505 510

Asp Glu Glu Asp Ser Glu Gly Pro Glu Lys Glu Gly Glu Gly His Ser  
 515 520 525

Tyr Phe Ser Ser Thr Lys Thr Leu Cys Val Ile Asp Pro Glu Asn Arg  
 530 535 540

Asp Ser Leu Gly Glu Thr Asp Ile Asp Ile Ala Thr Glu Asp Lys Ser  
 545 550 555 560

Pro Val Asp Thr

<210> 39

<211> 326

<212> PRT

<213> Homo sapiens

<400> 39

Met Pro Pro Ser Ile Ser Ala Phe Gln Ala Ala Tyr Ile Gly Ile Glu  
 1 5 10 15

Val Leu Ile Ala Leu Val Ser Val Pro Gly Asn Val Leu Val Ile Trp  
 20 25 30

Ala Val Lys Val Asn Gln Ala Leu Arg Asp Ala Thr Phe Cys Phe Ile  
 35 40 45  
 Val Ser Leu Ala Val Ala Asp Val Ala Val Gly Ala Leu Val Ile Pro  
 50 55 60  
 Leu Ala Ile Leu Ile Asn Ile Gly Pro Gln Thr Tyr Phe His Thr Cys  
 65 70 75 80  
 Leu Met Val Ala Cys Pro Val Leu Ile Leu Thr Gln Ser Ser Ile Leu  
 85 90 95  
 Ala Leu Leu Ala Ile Ala Val Asp Arg Tyr Leu Arg Val Lys Ile Pro  
 100 105 110  
 Leu Arg Tyr Lys Met Val Val Thr Pro Arg Arg Ala Ala Val Ala Ile  
 115 120 125  
 Ala Gly Cys Trp Ile Leu Ser Phe Val Val Gly Leu Thr Pro Met Phe  
 130 135 140  
 Gly Trp Asn Asn Leu Ser Ala Val Glu Arg Ala Trp Ala Ala Asn Gly  
 145 150 155 160  
 Ser Met Gly Glu Pro Val Ile Lys Cys Glu Phe Glu Lys Val Ile Ser  
 165 170 175  
 Met Glu Tyr Met Val Tyr Phe Asn Phe Phe Val Trp Val Leu Pro Pro  
 180 185 190  
 Leu Leu Leu Met Val Leu Ile Tyr Leu Glu Val Phe Tyr Leu Ile Arg  
 195 200 205  
 Lys Gln Leu Asn Lys Lys Val Ser Ala Ser Ser Gly Asp Pro Gln Lys  
 210 215 220  
 Tyr Tyr Gly Lys Glu Leu Lys Ile Ala Lys Ser Leu Ala Leu Ile Leu  
 225 230 235 240  
 Phe Leu Phe Ala Leu Ser Trp Leu Pro Leu His Ile Leu Asn Cys Ile  
 245 250 255  
 Thr Leu Phe Cys Pro Ser Cys His Lys Pro Ser Ile Leu Thr Tyr Ile  
 260 265 270

Ala Ile Phe Leu Thr His Gly Asn Ser Ala Met Asn Pro Ile Val Tyr  
 275 280 285

Ala Phe Arg Ile Gln Lys Phe Arg Val Thr Phe Leu Lys Ile Trp Asn  
 290 295 300

Asp His Phe Arg Cys Gln Pro Ala Pro Pro Ile Asp Glu Asp Leu Pro  
 305 310 315 320

Glu Glu Arg Pro Asp Asp  
 325

<210> 40

<211> 302

<212> PRT

<213> Homo sapiens

<400> 40

Ala Asn Ser Val Val Val Trp Val Asn Ile Gln Ala Lys Thr Thr Gly  
 1 5 10 15

Tyr Asp Thr His Cys Tyr Ile Leu Asn Leu Ala Ile Ala Asp Leu Trp  
 20 25 30

Val Val Leu Thr Ile Pro Val Trp Val Val Ser Leu Val Gln His Asn  
 35 40 45

Gln Trp Pro Met Gly Glu Leu Thr Cys Lys Val Thr His Leu Ile Phe  
 50 55 60

Ser Ile Asn Leu Phe Ser Ser Ile Phe Phe Leu Thr Cys Met Ser Val  
 65 70 75 80

Asp Arg Tyr Leu Ser Ile Thr Tyr Phe Thr Asn Thr Pro Ser Ser Arg  
 85 90 95

Lys Lys Met Val Arg Arg Val Val Cys Ile Leu Val Trp Leu Leu Ala  
 100 105 110

Phe Cys Val Ser Leu Pro Asp Thr Tyr Tyr Leu Lys Thr Val Thr Ser  
 115 120 125

Ala Ser Asn Asn Glu Thr Tyr Cys Arg Ser Phe Tyr Pro Glu His Ser  
 130 135 140

Ile Lys Glu Trp Leu Ile Gly Met Glu Leu Val Ser Val Val Leu Gly  
 145 150 155 160

Phe Ala Val Pro Phe Ser Ile Ile Ala Val Phe Tyr Phe Leu Leu Ala  
 165 170 175

Arg Ala Ile Ser Ala Ser Ser Asp Gln Glu Lys His Ser Ser Arg Lys  
 180 185 190

Ile Ile Phe Ser Tyr Val Val Val Phe Leu Val Cys Trp Leu Pro Tyr  
 195 200 205

His Val Ala Val Leu Leu Asp Ile Phe Ser Ile Leu His Tyr Ile Pro  
 210 215 220

Phe Thr Cys Arg Leu Glu His Ala Leu Phe Thr Ala Leu His Val Thr  
 225 230 235 240

Gln Cys Leu Ser Leu Val His Cys Cys Val Asn Pro Val Leu Tyr Ser  
 245 250 255

Phe Ile Asn Arg Asn Tyr Arg Tyr Glu Leu Met Lys Ala Phe Ile Phe  
 260 265 270

Lys Tyr Ser Ala Lys Thr Gly Leu Thr Lys Leu Ile Asp Ala Ser Arg  
 275 280 285

Val Ser Glu Thr Glu Tyr Ser Ala Leu Glu Gln Ser Thr Lys  
 290 295 300

<210> 41

<211> 380

<212> PRT

<213> Homo sapiens



&lt;400&gt; 41

Met Met Phe Ser Gly Phe Asn Ala Asp Tyr Glu Ala Ser Ser Ser Arg  
 1 5 10 15

Cys Ser Ser Ala Ser Pro Ala Gly Asp Ser Leu Ser Tyr Tyr His Ser  
 20 25 30

Pro Ala Asp Ser Phe Ser Ser Met Gly Ser Pro Val Asn Ala Gln Asp  
 35 40 45

Phe Cys Thr Asp Leu Ala Val Ser Ser Ala Asn Phe Ile Pro Thr Val  
 50 55 60

Thr Ala Ile Ser Thr Ser Pro Asp Leu Gln Trp Leu Val Gln Pro Ala  
 65 70 75 80

Leu Val Ser Ser Val Ala Pro Ser Gln Thr Arg Ala Pro His Pro Phe  
 85 90 95

Gly Val Pro Ala Pro Ser Ala Gly Ala Tyr Ser Arg Ala Gly Val Val  
 100 105 110

Lys Thr Met Thr Gly Gly Arg Ala Gln Ser Ile Gly Arg Arg Gly Lys  
 115 120 125

Val Glu Gln Leu Ser Pro Glu Glu Glu Glu Lys Arg Arg Ile Arg Arg  
 130 135 140

Glu Arg Asn Lys Met Ala Ala Ala Lys Cys Arg Asn Arg Arg Arg Glu  
 145 150 155 160

Leu Thr Asp Thr Leu Gln Ala Glu Thr Asp Gln Leu Glu Asp Glu Lys  
 165 170 175

Ser Ala Leu Gln Thr Glu Ile Ala Asn Leu Leu Lys Glu Lys Glu Lys  
 180 185 190

Leu Glu Phe Ile Leu Ala Ala His Arg Pro Ala Cys Lys Ile Pro Asp  
 195 200 205

Asp Leu Gly Phe Pro Glu Glu Met Ser Val Ala Ser Leu Asp Leu Thr  
 210 215 220

Gly Gly Leu Pro Glu Val Ala Thr Pro Glu Ser Glu Glu Ala Phe Thr  
 225 230 235 240

Leu Pro Leu Leu Asn Asp Pro Glu Pro Lys Pro Ser Val Glu Pro Val  
 245 250 255

Lys Ser Ile Ser Ser Met Glu Leu Lys Thr Glu Pro Phe Asp Asp Phe  
 260 265 270

Leu Phe Pro Ala Ser Ser Arg Pro Ser Gly Ser Glu Thr Ala Arg Ser  
 275 280 285

Val Pro Asp Met Asp Leu Ser Gly Ser Phe Tyr Ala Ala Asp Trp Glu  
 290 295 300

Pro Leu His Ser Gly Ser Leu Gly Met Gly Pro Met Ala Thr Glu Leu  
 305 310 315 320

Glu Pro Leu Cys Thr Pro Val Val Thr Cys Thr Pro Ser Cys Thr Ala  
 325 330 335

Tyr Thr Ser Ser Phe Val Phe Thr Tyr Pro Glu Ala Asp Ser Phe Pro  
 340 345 350

Ser Cys Ala Ala Ala His Arg Lys Gly Ser Ser Ser Asn Glu Pro Ser  
 355 360 365

Ser Asp Ser Leu Ser Ser Pro Thr Leu Leu Ala Leu  
 370 375 380

<210> 42

<211> 119

<212> PRT

<213> Homo sapiens

<400> 42

Met Lys Ala Leu Ser Pro Val Arg Gly Cys Tyr Glu Ala Val Cys Cys  
 1 5 10 15

Leu Ser Glu Arg Ser Leu Ala Ile Ala Arg Gly Arg Gly Lys Gly Pro  
 20 25 30

Ala Ala Glu Glu Pro Leu Ser Leu Leu Asp Asp Met Asn His Cys Tyr  
35 40 45

Ser Arg Leu Arg Glu Leu Val Pro Gly Val Pro Arg Gly Thr Gln Leu  
50 55 60

Ser Gln Val Glu Ile Leu Gln Arg Val Ile Asp Tyr Ile Leu Asp Leu  
65 70 75 80

Gln Val Val Leu Ala Glu Pro Ala Pro Gly Pro Pro Asp Gly Pro His  
85 90 95

Leu Pro Ile Gln Thr Ala Glu Leu Thr Pro Glu Leu Val Ile Ser Asn  
100 105 110

Asp Lys Arg Ser Phe Cys His  
115

<210> 43

<211> 496

<212> PRT

<213> Homo sapiens

<400> 43

Met Gly Thr Gln Lys Val Thr Pro Ala Leu Ile Phe Ala Ile Thr Val  
1 5 10 15

Ala Thr Ile Gly Ser Phe Gln Phe Gly Tyr Asn Thr Gly Val Ile Asn  
20 25 30

Ala Pro Glu Lys Ile Ile Lys Glu Phe Ile Asn Lys Thr Leu Thr Asp  
35 40 45

Lys Gly Asn Ala Pro Pro Ser Glu Val Leu Leu Thr Ser Leu Trp Ser  
50 55 60

Leu Ser Val Ala Ile Phe Ser Val Gly Gly Met Ile Gly Ser Phe Ser  
65 70 75 80

Val Gly Leu Phe Val Asn Arg Phe Gly Arg Arg Asn Ser Met Leu Ile  
 85 90 95

Val Asn Leu Leu Ala Val Thr Gly Gly Cys Phe Met Gly Leu Cys Lys  
 100 105 110

Val Ala Lys Ser Val Glu Met Leu Ile Leu Gly Arg Leu Val Ile Gly  
 115 120 125

Leu Phe Cys Gly Leu Cys Thr Gly Phe Val Pro Met Tyr Ile Gly Glu  
 130 135 140

Ile Ser Pro Thr Ala Leu Arg Gly Ala Phe Gly Thr Leu Asn Gln Leu  
 145 150 155 160

Gly Ile Val Val Gly Ile Leu Val Ala Gln Ile Phe Gly Leu Glu Phe  
 165 170 175

Ile Leu Gly Ser Glu Glu Leu Trp Pro Leu Leu Leu Gly Phe Thr Ile  
 180 185 190

Leu Pro Ala Ile Leu Gln Ser Ala Ala Leu Pro Phe Cys Pro Glu Ser  
 195 200 205

Pro Arg Phe Leu Leu Ile Asn Arg Lys Glu Glu Glu Asn Ala Lys Gln  
 210 215 220

Ile Leu Gln Arg Leu Trp Gly Thr Gln Asp Val Ser Gln Asp Ile Gln  
 225 230 235 240

Glu Met Lys Asp Glu Ser Ala Arg Met Ser Gln Glu Lys Gln Val Thr  
 245 250 255

Val Leu Glu Leu Phe Arg Val Ser Ser Tyr Arg Gln Pro Ile Ile Ile  
 260 265 270

Ser Ile Val Leu Gln Leu Ser Gln Gln Leu Ser Gly Ile Asn Ala Val  
 275 280 285

Phe Tyr Tyr Ser Thr Gly Ile Phe Lys Asp Ala Gly Val Gln Glu Pro  
 290 295 300

Ile Tyr Ala Thr Ile Gly Ala Gly Val Val Asn Thr Ile Phe Thr Val  
 305 310 315 320

Val Ser Leu Phe Leu Val Glu Arg Ala Gly Arg Arg Thr Leu His Met  
                                   325                                  330                                  335

Ile Gly Leu Gly Gly Met Ala Phe Cys Ser Thr Leu Met Thr Val Ser  
                                   340                                  345                                  350

Leu Leu Leu Lys Asp Asn Tyr Asn Gly Met Ser Phe Val Cys Ile Gly  
                                   355                                  360                                  365

Ala Ile Leu Val Phe Val Ala Phe Phe Glu Ile Gly Pro Gly Pro Ile  
                                   370                                  375                                  380

Pro Trp Phe Ile Val Ala Glu Leu Phe Ser Gln Gly Pro Arg Pro Ala  
                                   385                                  390                                  395                                  400

Ala Met Ala Val Ala Gly Cys Ser Asn Trp Thr Ser Asn Phe Leu Val  
                                   405                                  410                                  415

Gly Leu Leu Phe Pro Ser Ala Ala His Tyr Leu Gly Ala Tyr Val Phe  
                                   420                                  425                                  430

Ile Ile Phe Thr Gly Phe Leu Ile Thr Phe Leu Ala Phe Thr Phe Phe  
                                   435                                  440                                  445

Lys Val Pro Glu Thr Arg Gly Arg Thr Phe Glu Asp Ile Thr Arg Ala  
                                   450                                  455                                  460

Phe Glu Gly Gln Ala His Gly Ala Asp Arg Ser Gly Lys Asp Gly Val  
                                   465                                  470                                  475                                  480

Met Glu Met Asn Ser Ile Glu Pro Ala Lys Glu Thr Thr Thr Asn Val  
                                   485                                  490                                  495

<210> 44

<211> 352

<212> PRT

<213> Homo sapiens

&lt;400&gt; 44

```

Met Glu Gly Ile Ser Ile Tyr Thr Ser Asp Asn Tyr Thr Glu Glu Met
1          5          10          15

Gly Ser Gly Asp Tyr Asp Ser Met Lys Glu Pro Cys Phe Arg Glu Glu
20          25          30

Asn Ala Asn Phe Asn Lys Ile Phe Leu Pro Thr Ile Tyr Ser Ile Ile
35          40          45

Phe Leu Thr Gly Ile Val Gly Asn Gly Leu Val Ile Leu Val Met Gly
50          55          60

Tyr Gln Lys Lys Leu Arg Ser Met Thr Asp Lys Tyr Arg Leu His Leu
65          70          75          80

Ser Val Ala Asp Leu Leu Phe Val Ile Thr Leu Pro Phe Trp Ala Val
85          90          95

Asp Ala Val Ala Asn Trp Tyr Phe Gly Asn Phe Leu Cys Lys Ala Val
100         105         110

His Val Ile Tyr Thr Val Asn Leu Tyr Ser Ser Val Leu Ile Leu Ala
115         120         125

Phe Ile Ser Leu Asp Arg Tyr Leu Ala Ile Val His Ala Thr Asn Ser
130         135         140

Gln Arg Pro Arg Lys Leu Leu Ala Glu Lys Val Val Tyr Val Gly Val
145         150         155         160

Trp Ile Pro Ala Leu Leu Leu Thr Ile Pro Asp Phe Ile Phe Ala Asn
165         170         175

Val Ser Glu Ala Asp Asp Arg Tyr Ile Cys Asp Arg Phe Tyr Pro Asn
180         185         190

Asp Leu Trp Val Val Val Phe Gln Phe Gln His Ile Met Val Gly Leu
195         200         205

Ile Leu Pro Gly Ile Val Ile Leu Ser Cys Tyr Cys Ile Ile Ile Ser
210         215         220

```

Lys Leu Ser His Ser Lys Gly His Gln Lys Arg Lys Ala Leu Lys Thr  
 225 230 235 240

Thr Val Ile Leu Ile Leu Ala Phe Phe Ala Cys Trp Leu Pro Tyr Tyr  
 245 250 255

Ile Gly Ile Ser Ile Asp Ser Phe Ile Leu Leu Glu Ile Ile Lys Gln  
 260 265 270

Gly Cys Glu Phe Glu Asn Thr Val His Lys Trp Ile Ser Ile Thr Glu  
 275 280 285

Ala Leu Ala Phe Phe His Cys Cys Leu Asn Pro Ile Leu Tyr Ala Phe  
 290 295 300

Leu Gly Ala Lys Phe Lys Thr Ser Ala Gln His Ala Leu Thr Ser Val  
 305 310 315 320

Ser Arg Gly Ser Ser Leu Lys Ile Leu Ser Lys Gly Lys Arg Gly Gly  
 325 330 335

His Ser Ser Val Ser Thr Glu Ser Glu Ser Ser Ser Phe His Ser Ser  
 340 345 350

<210> 45

<211> 208

<212> PRT

<213> Homo sapiens

<400> 45

Met Lys Leu Leu Pro Ser Val Val Leu Lys Leu Phe Leu Ala Ala Val  
 1 5 10 15

Leu Ser Ala Leu Val Thr Gly Glu Ser Leu Glu Arg Leu Arg Arg Gly  
 20 25 30

Leu Ala Ala Gly Thr Ser Asn Pro Asp Pro Pro Thr Val Ser Thr Asp  
 35 40 45

Gln Leu Leu Pro Leu Gly Gly Gly Arg Asp Arg Lys Val Arg Asp Leu  
 50 55 60

Gln Glu Ala Asp Leu Asp Leu Leu Arg Val Thr Leu Ser Ser Lys Pro  
65 70 75 80

Gln Ala Leu Ala Thr Pro Asn Lys Glu Glu His Gly Lys Arg Lys Lys  
85 90 95

Lys Gly Lys Gly Leu Gly Lys Lys Arg Asp Pro Cys Leu Arg Lys Tyr  
100 105 110

Lys Asp Phe Cys Ile His Gly Glu Cys Lys Tyr Val Lys Glu Leu Arg  
115 120 125

Ala Pro Ser Cys Ile Cys His Pro Gly Tyr His Gly Glu Arg Cys His  
130 135 140

Gly Leu Ser Leu Pro Val Glu Asn Arg Leu Tyr Thr Tyr Asp His Thr  
145 150 155 160

Thr Ile Leu Ala Val Val Ala Val Val Leu Ser Ser Val Cys Leu Leu  
165 170 175

Val Ile Val Gly Leu Leu Met Phe Arg Tyr His Arg Arg Gly Gly Tyr  
180 185 190

Asp Val Glu Asn Glu Glu Lys Val Lys Leu Gly Met Thr Asn Ser His  
195 200 205

<210> 46

<211> 218

<212> PRT

<213> Homo sapiens

<400> 46

Met Ser Met Thr Leu Gly Tyr Trp Asp Ile Arg Gly Leu Ala His Ala  
1 5 10 15

Met Arg Leu Leu Leu Glu Tyr Thr Asp Ser Ser Tyr Glu Glu Lys Lys  
20 25 30

Tyr Thr Met Gly Asp Ala Pro Asp Tyr Asp Arg Ser Gln Trp Leu Asn  
35 40 45



Glu Lys Phe Lys Leu Gly Leu Asp Phe Pro Asn Leu Pro Tyr Leu Ile  
 50 55 60

Asp Gly Ala His Lys Ile Thr Gln Ser Asn Ala Ile Leu Cys Tyr Ile  
 65 70 75 80

Ala Arg Lys His Asn Leu Cys Gly Glu Thr Glu Glu Glu Lys Ile Arg  
 85 90 95

Val Asp Ile Leu Glu Asn Gln Ala Met Asp Val Ser Asn Gln Leu Ala  
 100 105 110

Arg Val Cys Tyr Ser Pro Asp Phe Glu Lys Leu Lys Pro Glu Tyr Leu  
 115 120 125

Glu Glu Leu Pro Thr Met Met Gln His Phe Ser Gln Phe Leu Gly Lys  
 130 135 140

Arg Pro Trp Phe Val Gly Asp Lys Ile Thr Phe Val Asp Phe Leu Ala  
 145 150 155 160

Tyr Asp Val Leu Asp Leu His Arg Ile Phe Glu Pro Asn Cys Leu Asp  
 165 170 175

Ala Phe Pro Asn Leu Lys Asp Phe Ile Ser Arg Phe Glu Gly Leu Glu  
 180 185 190

Lys Ile Ser Ala Tyr Met Lys Ser Ser Arg Phe Leu Pro Lys Pro Leu  
 195 200 205

Tyr Thr Arg Val Ala Val Trp Gly Asn Lys  
 210 215

<210> 47

<211> 543

<212> PRT

<213> Homo sapiens

&lt;400&gt; 47

Met Gly Thr Ser Leu Ser Pro Asn Asp Pro Trp Pro Leu Asn Pro Leu  
1 5 10 15

Ser Ile Gln Gln Thr Thr Leu Leu Leu Leu Leu Ser Val Leu Ala Thr  
20 25 30

Val His Val Gly Gln Arg Leu Leu Arg Gln Arg Arg Arg Gln Leu Arg  
35 40 45

Ser Ala Pro Pro Gly Pro Phe Ala Trp Pro Leu Ile Gly Asn Ala Ala  
50 55 60

Ala Val Gly Gln Ala Ala His Leu Ser Phe Ala Arg Leu Ala Arg Arg  
65 70 75 80

Tyr Gly Asp Val Phe Gln Ile Arg Leu Gly Ser Cys Pro Ile Val Val  
85 90 95

Leu Asn Gly Glu Arg Ala Ile His Gln Ala Leu Val Gln Gln Gly Ser  
100 105 110

Ala Phe Ala Asp Arg Pro Ala Phe Ala Ser Phe Arg Val Val Ser Gly  
115 120 125

Gly Arg Ser Met Ala Phe Gly His Tyr Ser Glu His Trp Lys Val Gln  
130 135 140

Arg Arg Ala Ala His Ser Met Met Arg Asn Phe Phe Thr Arg Gln Pro  
145 150 155 160

Arg Ser Arg Gln Val Leu Glu Gly His Val Leu Ser Glu Ala Arg Glu  
165 170 175

Leu Val Ala Leu Leu Val Arg Gly Ser Ala Asp Gly Ala Phe Leu Asp  
180 185 190

Pro Arg Pro Leu Thr Val Val Ala Val Ala Asn Val Met Ser Ala Val  
195 200 205

Cys Phe Gly Cys Arg Tyr Ser His Asp Asp Pro Glu Phe Arg Glu Leu  
210 215 220

Leu Ser His Asn Glu Glu Phe Gly Arg Thr Val Gly Ala Gly Ser Leu  
 225 230 235 240

Val Asp Val Met Pro Trp Leu Gln Tyr Phe Pro Asn Pro Val Arg Thr  
 245 250 255

Val Phe Arg Glu Phe Glu Gln Leu Asn Arg Asn Phe Ser Asn Phe Ile  
 260 265 270

Leu Asp Lys Phe Leu Arg His Cys Glu Ser Leu Arg Pro Gly Ala Ala  
 275 280 285

Pro Arg Asp Met Met Asp Ala Phe Ile Leu Ser Ala Glu Lys Lys Ala  
 290 295 300

Ala Gly Asp Ser His Gly Gly Gly Ala Arg Leu Asp Leu Glu Asn Val  
 305 310 315 320

Pro Ala Thr Ile Thr Asp Ile Phe Gly Ala Ser Gln Asp Thr Leu Ser  
 325 330 335

Thr Ala Leu Gln Trp Leu Leu Leu Leu Phe Thr Arg Tyr Pro Asp Val  
 340 345 350

Gln Thr Arg Val Gln Ala Glu Leu Asp Gln Val Val Gly Arg Asp Arg  
 355 360 365

Leu Pro Cys Met Gly Asp Gln Pro Asn Leu Pro Tyr Val Leu Ala Phe  
 370 375 380

Leu Tyr Glu Ala Met Arg Phe Ser Ser Phe Val Pro Val Thr Ile Pro  
 385 390 395 400

His Ala Thr Thr Ala Asn Thr Ser Val Leu Gly Tyr His Ile Pro Lys  
 405 410 415

Asp Thr Val Val Phe Val Asn Gln Trp Ser Val Asn His Asp Pro Val  
 420 425 430

Lys Trp Pro Asn Pro Glu Asn Phe Asp Pro Ala Arg Phe Leu Asp Lys  
 435 440 445

Asp Gly Leu Ile Asn Lys Asp Leu Thr Ser Arg Val Met Ile Phe Ser  
 450 455 460

Val Gly Lys Arg Arg Cys Ile Gly Glu Glu Leu Ser Lys Met Gln Leu  
465 470 475 480

Phe Leu Phe Ile Ser Ile Leu Ala His Gln Cys Asp Phe Arg Ala Asn  
485 490 495

Pro Asn Glu Pro Ala Lys Met Asn Phe Ser Tyr Gly Leu Thr Ile Lys  
500 505 510

Pro Lys Ser Phe Lys Val Asn Val Thr Leu Arg Glu Ser Met Glu Leu  
515 520 525

Leu Asp Ser Ala Val Gln Asn Leu Gln Ala Lys Glu Thr Cys Gln  
530 535 540

<210> 48

<211> 300

<212> PRT

<213> Homo sapiens

<400> 48

Met Arg Ile Ala Val Ile Cys Phe Cys Leu Leu Gly Ile Thr Cys Ala  
1 5 10 15

Ile Pro Val Lys Gln Ala Asp Ser Gly Ser Ser Glu Glu Lys Gln Leu  
20 25 30

Tyr Asn Lys Tyr Pro Asp Ala Val Ala Thr Trp Leu Asn Pro Asp Pro  
35 40 45

Ser Gln Lys Gln Asn Leu Leu Ala Pro Gln Thr Leu Pro Ser Lys Ser  
50 55 60

Asn Glu Ser His Asp His Met Asp Asp Met Asp Asp Glu Asp Asp Asp  
65 70 75 80

Asp His Val Asp Ser Gln Asp Ser Ile Asp Ser Asn Asp Ser Asp Asp  
85 90 95

Val Asp Asp Thr Asp Asp Ser His Gln Ser Asp Glu Ser His His Ser  
100 105 110

Asp Glu Ser Asp Glu Leu Val Thr Asp Phe Pro Thr Asp Leu Pro Ala  
 115 120 125  
 Thr Glu Val Phe Thr Pro Val Val Pro Thr Val Asp Thr Tyr Asp Gly  
 130 135 140  
 Arg Gly Asp Ser Val Val Tyr Gly Leu Arg Ser Lys Ser Lys Lys Phe  
 145 150 155 160  
 Arg Arg Pro Asp Ile Gln Tyr Pro Asp Ala Thr Asp Glu Asp Ile Thr  
 165 170 175  
 Ser His Met Glu Ser Glu Glu Leu Asn Gly Ala Tyr Lys Ala Ile Pro  
 180 185 190  
 Val Ala Gln Asp Leu Asn Ala Pro Ser Asp Trp Asp Ser Arg Gly Lys  
 195 200 205  
 Asp Ser Tyr Glu Thr Ser Gln Leu Asp Asp Gln Ser Ala Glu Thr His  
 210 215 220  
 Ser His Lys Gln Ser Arg Leu Tyr Lys Arg Lys Ala Asn Asp Glu Ser  
 225 230 235 240  
 Asn Glu His Ser Asp Val Ile Asp Ser Gln Glu Leu Ser Lys Val Ser  
 245 250 255  
 Arg Glu Phe His Ser His Glu Phe His Ser His Glu Asp Met Leu Val  
 260 265 270  
 Val Asp Pro Lys Ser Lys Glu Glu Asp Lys His Leu Lys Phe Arg Ile  
 275 280 285  
 Ser His Glu Leu Asp Ser Ala Ser Ser Glu Val Asn  
 290 295 300

&lt;210&gt; 49

&lt;211&gt; 99

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 49

Met Lys Val Ser Ala Ala Leu Leu Cys Leu Leu Leu Ile Ala Ala Thr  
 1 5 10 15

Phe Ile Pro Gln Gly Leu Ala Gln Pro Asp Ala Ile Asn Ala Pro Val  
 20 25 30

Thr Cys Cys Tyr Asn Phe Thr Asn Arg Lys Ile Ser Val Gln Arg Leu  
 35 40 45

Ala Ser Tyr Arg Arg Ile Thr Ser Ser Lys Cys Pro Lys Glu Ala Val  
 50 55 60

Ile Phe Lys Thr Ile Val Ala Lys Glu Ile Cys Ala Asp Pro Lys Gln  
 65 70 75 80

Lys Trp Val Gln Asp Ser Met Asp His Leu Asp Lys Gln Thr Gln Thr  
 85 90 95

Pro Lys Thr

&lt;210&gt; 50

&lt;211&gt; 470

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 50

Met Lys Phe Leu Leu Ile Leu Leu Leu Gln Ala Thr Ala Ser Gly Ala  
 1 5 10 15

Leu Pro Leu Asn Ser Ser Thr Ser Leu Glu Lys Asn Asn Val Leu Phe  
 20 25 30

Gly Glu Arg Tyr Leu Glu Lys Phe Tyr Gly Leu Glu Ile Asn Lys Leu  
 35 40 45

Pro Val Thr Lys Met Lys Tyr Ser Gly Asn Leu Met Lys Glu Lys Ile  
 50 55 60

Gln Glu Met Gln His Phe Leu Gly Leu Lys Val Thr Gly Gln Leu Asp  
65 70 75 80

Thr Ser Thr Leu Glu Met Met His Ala Pro Arg Cys Gly Val Pro Asp  
85 90 95

Leu His His Phe Arg Glu Met Pro Gly Gly Pro Val Trp Arg Lys His  
100 105 110

Tyr Ile Thr Tyr Arg Ile Asn Asn Tyr Thr Pro Asp Met Asn Arg Glu  
115 120 125

Asp Val Asp Tyr Ala Ile Arg Lys Ala Phe Gln Val Trp Ser Asn Val  
130 135 140

Thr Pro Leu Lys Phe Ser Lys Ile Asn Thr Gly Met Ala Asp Ile Leu  
145 150 155 160

Val Val Phe Ala Arg Gly Ala His Gly Asp Phe His Ala Phe Asp Gly  
165 170 175

Lys Gly Gly Ile Leu Ala His Ala Phe Gly Pro Gly Ser Gly Ile Gly  
180 185 190

Gly Asp Ala His Phe Asp Glu Asp Glu Phe Trp Thr Thr His Ser Gly  
195 200 205

Gly Thr Asn Leu Phe Leu Thr Ala Val His Glu Ile Gly His Ser Leu  
210 215 220

Gly Leu Gly His Ser Ser Asp Pro Lys Ala Val Met Phe Pro Thr Tyr  
225 230 235 240

Lys Tyr Val Asp Ile Asn Thr Phe Arg Leu Ser Ala Asp Asp Ile Arg  
245 250 255

Gly Ile Gln Ser Leu Tyr Gly Asp Pro Lys Glu Asn Gln Arg Leu Pro  
260 265 270

Asn Pro Asp Asn Ser Glu Pro Ala Leu Cys Asp Pro Asn Leu Ser Phe  
275 280 285

Asp Ala Val Thr Thr Val Gly Asn Lys Ile Phe Phe Phe Lys Asp Arg  
290 295 300

Phe Phe Trp Leu Lys Val Ser Glu Arg Pro Lys Thr Ser Val Asn Leu  
 305 310 315 320

Ile Ser Ser Leu Trp Pro Thr Leu Pro Ser Gly Ile Glu Ala Ala Tyr  
 325 330 335

Glu Ile Glu Ala Arg Asn Gln Val Phe Leu Phe Lys Asp Asp Lys Tyr  
 340 345 350

Trp Leu Ile Ser Asn Leu Arg Pro Glu Pro Asn Tyr Pro Lys Ser Ile  
 355 360 365

His Ser Phe Gly Phe Pro Asn Phe Val Lys Lys Ile Asp Ala Ala Val  
 370 375 380

Phe Asn Pro Arg Phe Tyr Arg Thr Tyr Phe Phe Val Asp Asn Gln Tyr  
 385 390 395 400

Trp Arg Tyr Asp Glu Arg Arg Gln Met Met Asp Pro Gly Tyr Pro Lys  
 405 410 415

Leu Ile Thr Lys Asn Phe Gln Gly Ile Gly Pro Lys Ile Asp Ala Val  
 420 425 430

Phe Tyr Ser Lys Asn Lys Tyr Tyr Tyr Phe Phe Gln Gly Ser Asn Gln  
 435 440 445

Phe Glu Tyr Asp Phe Leu Leu Gln Arg Ile Thr Lys Thr Leu Lys Ser  
 450 455 460

Asn Ser Trp Phe Gly Cys  
 465 470

<210> 51

<211> 101

<212> PRT

<213> Homo sapiens



&lt;400&gt; 51

Met Gly Thr Arg Leu Leu Pro Ala Leu Phe Leu Val Leu Leu Val Leu  
 1 5 10 15

Gly Phe Glu Val Gln Gly Thr Gln Gln Pro Gln Gln Asp Glu Met Pro  
 20 25 30

Ser Pro Thr Phe Leu Thr Gln Val Lys Glu Ser Leu Ser Ser Tyr Trp  
 35 40 45

Glu Ser Ala Lys Thr Ala Ala Gln Asn Leu Tyr Glu Lys Thr Tyr Leu  
 50 55 60

Pro Ala Val Asp Glu Lys Leu Arg Asp Leu Tyr Ser Lys Ser Thr Ala  
 65 70 75 80

Ala Met Ser Thr Tyr Thr Gly Ile Phe Thr Asp Gln Val Leu Ser Val  
 85 90 95

Leu Lys Gly Glu Glu  
 100

&lt;210&gt; 52

&lt;211&gt; 150

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 52

Met Phe Arg Lys Gly Lys Lys Arg His Ser Ser Ser Ser Ser Gln Ser  
 1 5 10 15

Ser Glu Ile Ser Thr Lys Ser Lys Ser Val Asp Ser Ser Leu Gly Gly  
 20 25 30

Leu Ser Arg Ser Ser Thr Val Ala Ser Leu Asp Thr Asp Ser Thr Lys  
 35 40 45

Ser Ser Gly Gln Ser Asn Asn Asn Ser Asp Thr Cys Ala Glu Phe Arg  
 50 55 60

Ile Lys Tyr Val Gly Ala Ile Glu Lys Leu Lys Leu Ser Glu Gly Lys  
65 70 75 80

Gly Leu Glu Gly Pro Leu Asp Leu Ile Asn Tyr Ile Asp Val Ala Gln  
85 90 95

Gln Asp Gly Lys Leu Pro Phe Val Pro Pro Glu Glu Glu Phe Ile Met  
100 105 110

Gly Val Ser Lys Tyr Gly Ile Lys Val Ser Thr Ser Asp Gln Tyr Glu  
115 120 125

Gln Ala Gln Ala Ile Cys Lys Val Leu Ser Thr Ala Phe Asp Ser Val  
130 135 140

Leu Thr Ser Glu Lys Pro  
145 150

<210> 53

<211> 381

<212> PRT

<213> Homo sapiens

<400> 53

Met Ile Asp Thr Leu Arg Pro Val Pro Phe Ala Ser Glu Met Ala Ile  
1 5 10 15

Ser Lys Thr Val Ala Trp Leu Asn Glu Gln Leu Glu Leu Gly Asn Glu  
20 25 30

Arg Leu Leu Leu Met Asp Cys Arg Pro Gln Glu Leu Tyr Glu Ser Ser  
35 40 45

His Ile Glu Ser Ala Ile Asn Val Ala Ile Pro Gly Ile Met Leu Arg  
50 55 60

Arg Leu Gln Lys Gly Asn Leu Pro Val Arg Ala Leu Phe Thr Arg Gly  
65 70 75 80

Glu Asp Arg Asp Arg Phe Thr Arg Arg Cys Gly Thr Asp Thr Val Val  
85 90 95

Leu Tyr Asp Glu Ser Ser Ser Asp Trp Asn Glu Asn Thr Gly Gly Glu  
 100 105 110

Ser Leu Leu Gly Leu Leu Leu Lys Lys Leu Lys Asp Glu Gly Cys Arg  
 115 120 125

Ala Phe Tyr Leu Glu Gly Gly Phe Ser Lys Phe Gln Ala Glu Phe Ser  
 130 135 140

Leu His Cys Glu Thr Asn Leu Asp Gly Ser Cys Ser Ser Ser Ser Pro  
 145 150 155 160

Pro Leu Pro Val Leu Gly Leu Gly Gly Leu Arg Ile Ser Ser Asp Ser  
 165 170 175

Ser Ser Asp Ile Glu Ser Asp Leu Asp Arg Asp Pro Asn Ser Ala Thr  
 180 185 190

Asp Ser Asp Gly Ser Pro Leu Ser Asn Ser Gln Pro Ser Phe Pro Val  
 195 200 205

Glu Ile Leu Pro Phe Leu Tyr Leu Gly Cys Ala Lys Asp Ser Thr Asn  
 210 215 220

Leu Asp Val Leu Glu Glu Phe Gly Ile Lys Tyr Ile Leu Asn Val Thr  
 225 230 235 240

Pro Asn Leu Pro Asn Leu Phe Glu Asn Ala Gly Glu Phe Lys Tyr Lys  
 245 250 255

Gln Ile Pro Ile Ser Asp His Trp Ser Gln Asn Leu Ser Gln Phe Phe  
 260 265 270

Pro Glu Ala Ile Ser Phe Ile Asp Glu Ala Arg Gly Lys Asn Cys Gly  
 275 280 285

Val Leu Val His Cys Leu Ala Gly Ile Ser Arg Ser Val Thr Val Thr  
 290 295 300

Val Ala Tyr Leu Met Gln Lys Leu Asn Leu Ser Met Asn Asp Ala Tyr  
 305 310 315 320

Asp Ile Val Lys Met Lys Lys Ser Asn Ile Ser Pro Asn Phe Asn Phe  
 325 330 335

Met Gly Gln Leu Leu Asp Phe Glu Arg Thr Leu Gly Leu Ser Ser Pro  
                   340                  345                  350

Cys Asp Asn Arg Val Pro Ala Gln Gln Leu Tyr Phe Thr Thr Pro Ser  
           355                  360                  365

Asn Gln Asn Val Tyr Gln Val Asp Ser Leu Gln Ser Thr  
       370                  375                  380

<210> 54

<211> 542

<212> PRT

<213> Homo sapiens

<400> 54

Met Thr Lys Ser Asn Gly Glu Glu Pro Lys Met Gly Gly Arg Met Glu  
   1                  5                  10                  15

Arg Phe Gln Gln Gly Val Arg Lys Arg Thr Leu Leu Ala Lys Lys Lys  
           20                  25                  30

Val Gln Asn Ile Thr Lys Glu Asp Val Lys Ser Tyr Leu Phe Arg Asn  
       35                  40                  45

Ala Phe Val Leu Leu Thr Val Thr Ala Val Ile Val Gly Thr Ile Leu  
       50                  55                  60

Gly Phe Thr Leu Arg Pro Tyr Arg Met Ser Tyr Arg Glu Val Lys Tyr  
       65                  70                  75                  80

Phe Ser Phe Pro Gly Glu Leu Leu Met Arg Met Leu Gln Met Leu Val  
           85                  90                  95

Leu Pro Leu Ile Ile Ser Ser Leu Val Thr Gly Met Ala Ala Leu Asp  
           100                  105                  110

Ser Lys Ala Ser Gly Lys Met Gly Met Arg Ala Val Val Tyr Tyr Met  
       115                  120                  125

Thr Thr Thr Ile Ile Ala Val Val Ile Gly Ile Ile Ile Val Ile Ile  
 130 135 140

Ile His Pro Gly Lys Gly Thr Lys Glu Asn Met His Arg Glu Gly Lys  
 145 150 155 160

Ile Val Arg Val Thr Ala Ala Asp Ala Phe Leu Asp Leu Ile Arg Asn  
 165 170 175

Met Phe Pro Pro Asn Leu Val Glu Ala Cys Phe Lys Gln Phe Lys Thr  
 180 185 190

Asn Tyr Glu Lys Arg Ser Phe Lys Val Pro Ile Gln Ala Asn Glu Thr  
 195 200 205

Leu Val Gly Ala Val Ile Asn Asn Val Ser Glu Ala Met Glu Thr Leu  
 210 215 220

Thr Arg Ile Thr Glu Glu Leu Val Pro Val Pro Gly Ser Val Asn Gly  
 225 230 235 240

Val Asn Ala Leu Gly Leu Val Val Phe Ser Met Cys Phe Gly Phe Val  
 245 250 255

Ile Gly Asn Met Lys Glu Gln Gly Gln Ala Leu Arg Glu Phe Phe Asp  
 260 265 270

Ser Leu Asn Glu Ala Ile Met Arg Leu Val Ala Val Ile Met Trp Tyr  
 275 280 285

Ala Pro Val Gly Ile Leu Phe Leu Ile Ala Gly Lys Ile Val Glu Met  
 290 295 300

Glu Asp Met Gly Val Ile Gly Gly Gln Leu Ala Met Tyr Thr Val Thr  
 305 310 315 320

Val Ile Val Gly Leu Leu Ile His Ala Val Ile Val Leu Pro Leu Leu  
 325 330 335

Tyr Phe Leu Val Thr Arg Lys Asn Pro Trp Val Phe Ile Gly Gly Leu  
 340 345 350

Leu Gln Ala Leu Ile Thr Ala Leu Gly Thr Ser Ser Ser Ser Ala Thr  
 355 360 365

Leu Pro Ile Thr Phe Lys Cys Leu Glu Glu Asn Asn Gly Val Asp Lys  
 370 375 380

Arg Val Thr Arg Phe Val Leu Pro Val Gly Ala Thr Ile Asn Met Asp  
 385 390 395 400

Gly Thr Ala Leu Tyr Glu Ala Leu Ala Ala Ile Phe Ile Ala Gln Val  
 405 410 415

Asn Asn Phe Glu Leu Asn Phe Gly Gln Ile Ile Thr Ile Ser Ile Thr  
 420 425 430

Ala Thr Ala Ala Ser Ile Gly Ala Ala Gly Ile Pro Gln Ala Gly Leu  
 435 440 445

Val Thr Met Val Ile Val Leu Thr Ser Val Gly Leu Pro Thr Asp Asp  
 450 455 460

Ile Thr Leu Ile Ile Ala Val Asp Trp Phe Leu Asp Arg Leu Arg Thr  
 465 470 475 480

Thr Thr Asn Val Leu Gly Asp Ser Leu Gly Ala Gly Ile Val Glu His  
 485 490 495

Leu Ser Arg His Glu Leu Lys Asn Arg Asp Val Glu Met Gly Asn Ser  
 500 505 510

Val Ile Glu Glu Asn Glu Met Lys Lys Pro Tyr Gln Leu Ile Ala Gln  
 515 520 525

Asp Asn Glu Thr Glu Lys Pro Ile Asp Ser Glu Thr Lys Met  
 530 535 540

<210> 55

<211> 328

<212> PRT

<213> Homo sapiens

<400> 55

Met Leu Pro Arg Val Gly Cys Pro Ala Leu Pro Leu Pro Pro Pro Pro  
 1 5 10 15

Leu Leu Pro Leu Leu Pro Leu Leu Leu Leu Leu Leu Gly Ala Ser Gly  
 20 25 30

Gly Gly Gly Gly Ala Arg Ala Glu Val Leu Phe Arg Cys Pro Pro Cys  
 35 40 45

Thr Pro Glu Arg Leu Ala Ala Cys Gly Pro Pro Pro Val Ala Pro Pro  
 50 55 60

Ala Ala Val Ala Ala Val Ala Gly Gly Ala Arg Met Pro Cys Ala Glu  
 65 70 75 80

Leu Val Arg Glu Pro Gly Cys Gly Cys Cys Ser Val Cys Ala Arg Leu  
 85 90 95

Glu Gly Glu Ala Cys Gly Val Tyr Thr Pro Arg Cys Gly Gln Gly Leu  
 100 105 110

Arg Cys Tyr Pro His Pro Gly Ser Glu Leu Pro Leu Gln Ala Leu Val  
 115 120 125

Met Gly Glu Gly Thr Cys Glu Lys Arg Arg Asp Ala Glu Tyr Gly Ala  
 130 135 140

Ser Pro Glu Gln Val Ala Asp Asn Gly Asp Asp His Ser Glu Gly Gly  
 145 150 155 160

Leu Val Glu Asn His Val Asp Ser Thr Met Asn Met Leu Gly Gly Gly  
 165 170 175

Gly Ser Ala Gly Arg Lys Pro Leu Lys Ser Gly Met Lys Glu Leu Ala  
 180 185 190

Val Phe Arg Glu Lys Val Thr Glu Gln His Arg Gln Met Gly Lys Gly  
 195 200 205

Gly Lys His His Leu Gly Leu Glu Glu Pro Lys Lys Leu Arg Pro Pro  
 210 215 220

Pro Ala Arg Thr Pro Cys Gln Gln Glu Leu Asp Gln Val Leu Glu Arg  
 225 230 235 240

Ile Ser Thr Met Arg Leu Pro Asp Glu Arg Gly Pro Leu Glu His Leu  
                           245                          250                          255

Tyr Ser Leu His Ile Pro Asn Cys Asp Lys His Gly Leu Tyr Asn Leu  
                           260                          265                          270

Lys Gln Cys Lys Met Ser Leu Asn Gly Gln Arg Gly Glu Cys Trp Cys  
                           275                          280                          285

Val Asn Pro Asn Thr Gly Lys Leu Ile Gln Gly Ala Pro Thr Ile Arg  
                           290                          295                          300

Gly Asp Pro Glu Cys His Leu Phe Tyr Asn Glu Gln Gln Glu Ala Cys  
  305                          310                          315                          320

Gly Val His Thr Gln Arg Met Gln  
                           325

<210> 56

<211> 38

<212> PRT

<213> Homo sapiens

<400> 56

Met Gln Ser Met Pro Gln Ser Pro Ala Val Ile Thr Ser Pro Ile Gly  
  1                          5                          10                          15

Arg Ser Gln Cys Arg Gly Ser Arg Ala Ile Glu Glu Ser Pro Ala Ala  
                           20                          25                          30

Ser Val Pro Asn Lys Leu  
                           35

<210> 57

<211> 29

<212> DNA

<213> Artificial DNA



<400> 57  
acgttcagac cacctattcc cttcttgcg 29

<210> 58

<211> 24

<212> DNA

<213> Artificial DNA

<400> 58  
attcctgctt ttcagagtga gaca 24

<210> 59

<211> 24

<212> DNA

<213> Artificial DNA

<400> 59  
gaacagcttc cctcactgtg taca 24

<210> 60

<211> 19

<212> DNA

<213> Artificial DNA

<400> 60  
tcgccaggcc ctgtggtgg 19

<210> 61

<211> 16

<212> DNA

<213> Artificial DNA

<400> 61  
agggcggttc gcaggt 16

<210> 62

<211> 23

<212> DNA

<213> Artificial DNA

<400> 62

gggcctttct gtgaaagttg taa

23

<210> 63

<211> 23

<212> DNA

<213> Artificial DNA

<400> 63

ttcccagcat catccaggcc cag

23

<210> 64

<211> 21

<212> DNA

<213> Artificial DNA

<400> 64

cgagcccttt gatgacttcc t

21

<210> 65

<211> 20

<212> DNA

<213> Artificial DNA

<400> 65

ggctcccagt ctgctgcata

20

<210> 66

<211> 24

<212> DNA

<213> Artificial DNA

<400> 66

cgcgtctctc aagctcgccct cttc

24

<210> 67

<211> 22

<212> DNA

<213> Artificial DNA

<400> 67

cagctacttt tctggtcagg gc

22

<210> 68

<211> 20

<212> DNA

<213> Artificial DNA

<400> 68

acaatcaggc gttccagctc

20

<210> 69

<211> 31

<212> DNA

<213> Artificial DNA

<400> 69

acagtggagc tctgtattag aaagcccctc a

31

<210> 70

<211> 29

<212> DNA

<213> Artificial DNA

<400> 70

gaggaagata ctgtggtact gtcataaaa

29

<210> 71

<211> 22

<212> DNA

<213> Artificial DNA

<400> 71

gagttacctg gccttcccag tt

22

<210> 72

<211> 25

<212> DNA

<213> Artificial DNA

<400> 72

ccaacttcag cttcatgggc cagct

25

<210> 73

<211> 20

<212> DNA

<213> Artificial DNA

<400> 73

gcagaggcga agcatcatct

20

<210> 74

<211> 21

<212> DNA

<213> Artificial DNA

<400> 74

cagcacctgg gactcaaact g

21

<210> 75

<211> 22

<212> DNA

<213> Artificial DNA

<400> 75

tgccagtget tgcagaccct gc

22

<210> 76

<211> 18

<212> DNA

<213> Artificial DNA

<400> 76

cctggccact gaactgcg

18

<210> 77

<211> 23

<212> DNA

<213> Artificial DNA

<400> 77

tg gatgttct tg aggtgaat tcc

23

<210> 78

<211> 26

<212> DNA

<213> Artificial DNA

<400> 78

tgaattccct gcagtgtctg caagca

26

<210> 79

<211> 25

<212> DNA

<213> Artificial DNA

<400> 79

acattcacac tttggatggt cttga

25

<210> 80

<211> 19

<212> DNA

<213> Artificial DNA

<400> 80

gagcgtcggt ggtcactga

19

<210> 81

<211> 35

<212> DNA

<213> Artificial DNA

<400> 81

ttctgactct aagtggcatt caaggagtac ctctc

35

<210> 82

<211> 23

<212> DNA

<213> Artificial DNA

<400> 82

cgattctgat ttgctgcctt atc

23

<210> 83

<211> 21

<212> DNA

<213> Artificial DNA

<400> 83

gcaggtagag cgtacgggtt t

21

<210> 84

<211> 25

<212> DNA

<213> Artificial DNA

<400> 84

tgatgcgggc tgtccattgc cgata

25

<210> 85

<211> 22

<212> DNA

<213> Artificial DNA

<400> 85

ttgtctccct gctggaaaaa tt

22

<210> 86

<211> 19

<212> DNA

<213> Artificial DNA

<400> 86

tgagcaaccc caccaagtc

19

<210> 87

<211> 22

<212> DNA

<213> Artificial DNA

<400> 87

ccgcatccag aagttccgcg tc

22

<210> 88

<211> 18

<212> DNA

<213> Artificial DNA

<400> 88

gcaactcggc catgaacc

18

<210> 89

<211> 24

<212> DNA

<213> Artificial DNA

<400> 89

ggtcattcca aatcttaagg aagg

24



<210> 90

<211> 23

<212> DNA

<213> Artificial DNA

<400> 90

ccggtccttc taccgccgagc aca

23

<210> 91

<211> 22

<212> DNA

<213> Artificial DNA

<400> 91

ctgcgtccaa caatgagacc ta

22

<210> 92

<211> 20

<212> DNA

<213> Artificial DNA

<400> 92

ccgatcagcc actccttgat

20

<210> 93

<211> 29

<212> DNA

<213> Artificial DNA

<400> 93

aggaccttat ctgtgcgtga aacacacca

29

<210> 94

<211> 25

<212> DNA

<213> Artificial DNA

<400> 94

cctagagggt tcctgtagac ctagg

25

<210> 95

<211> 19

<212> DNA

<213> Artificial DNA

<400> 95

agtccttgag gcccacagc

19

<210> 96

<211> 22

<212> DNA

<213> Artificial DNA

<400> 96

cgccgccttg gcatagtttg ga

22

<210> 97

<211> 21

<212> DNA

<213> Artificial DNA

<400> 97

tccaaggaga ccagaagacc a

21

<210> 98

<211> 20

<212> DNA

<213> Artificial DNA

<400> 98

gtgtggaagg agtggctgct

20

<210> 99

<211> 27

<212> DNA

<213> Artificial DNA

<400> 99

tagatctgga aaggacggcg tcatgga

27

<210> 100

<211> 16

<212> DNA

<213> Artificial DNA

<400> 100

gcaggcacac ggtgca

16

<210> 101

<211> 20

<212> DNA

<213> Artificial DNA

<400> 101

gcaggctcga tgctgttcat

20

<210> 102

<211> 29

<212> DNA

<213> Artificial DNA

<400> 102

ctccaaagga aagcgaggtg gacattcat

29

<210> 103

<211> 19

<212> DNA

<213> Artificial DNA

<400> 103

gggtccagcc tcaagatcc

19

<210> 104

<211> 26

<212> DNA

<213> Artificial DNA

<400> 104

acttgaagac tcagactcag tggaaa

26

<210> 105

<211> 26

<212> DNA

<213> Artificial DNA

<400> 105

cgtgtcctct ccctgccaaag tctcag

26

<210> 106

<211> 29

<212> DNA

<213> Artificial DNA

<400> 106

gtcaaagtgt aacagatatc agtgtctcc

29

<210> 107

<211> 20

<212> DNA

<213> Artificial DNA

<400> 107

tacaggcatg gaagcccaac

20

<210> 108

<211> 31

<212> DNA

<213> Artificial DNA

<400> 108

cctacaatga tgcagcactt ctcacagttc c

31

<210> 109

<211> 24

<212> DNA

<213> Artificial DNA

<400> 109

gaagccagaa tacttggagg aact

24

<210> 110

<211> 22

<212> DNA

<213> Artificial DNA

<400> 110

atacgggtgga ggtcaaggac at

22

<210> 111

<211> 25

<212> DNA

<213> Artificial DNA

<400> 111

attcctcatg ccaccactgc caaca

25

<210> 112

<211> 20

<212> DNA

<213> Artificial DNA

<400> 112

tccagctttg tgectgtcac

20

<210> 113

<211> 20

<212> DNA

<213> Artificial DNA

<400> 113

gggaatgtgg tagcccaaga

20

<210> 114

<211> 27

<212> DNA

<213> Artificial DNA

<400> 114

tcacctcaca catggaaagc gaggagt

27

<210> 115

<211> 23

<212> DNA

<213> Artificial DNA

<400> 115

ccagtaccct gatgctacag acg

23

<210> 116

<211> 20

<212> DNA

<213> Artificial DNA

<400> 116

tggccttgta tgcaccattc

20

<210> 117

<211> 29

<212> DNA

<213> Artificial DNA

<400> 117

acccaaactc cgaagacttg aacacccac

29

<210> 118

<211> 19

<212> DNA

<213> Artificial DNA

<400> 118

tggaccacct ggacaagca

19

<210> 119

<211> 21

<212> DNA

<213> Artificial DNA

<400> 119

gctgcagatt cttgggttgt g

21

<210> 120

<211> 25

<212> DNA

<213> Artificial DNA

<400> 120

ccttatggcc aaccttgcca tctgg

25

<210> 121

<211> 26

<212> DNA

<213> Artificial DNA

<400> 121

ctgagagacc aaagaccagt gttaat

26



<210> 122

<211> 24

<212> DNA

<213> Artificial DNA

<400> 122

tttctggctt caatttcata agca

24

<210> 123

<211> 25

<212> DNA

<213> Artificial DNA

<400> 123

acagccgccc agaacctgta cgaga

25

<210> 124

<211> 22

<212> DNA

<213> Artificial DNA

<400> 124

tccagttact gggagtcagc aa

22

<210> 125

<211> 21

<212> DNA

<213> Artificial DNA

<400> 125

tctacagcgg gcaggatatgt c

21

<210> 126

<211> 23

<212> DNA

<213> Artificial DNA

<400> 126

ttcacgatcc agcactgtgg cca

23

<210> 127

<211> 24

<212> DNA

<213> Artificial DNA

<400> 127

agcaagtctg tggattctag cctt

24

<210> 128

<211> 22

<212> DNA

<213> Artificial DNA

<400> 128

cctgagcttt tgggtggaatc tg

22

<210> 129

<211> 36

<212> DNA

<213> Artificial DNA

<400> 129

atgcttgact ttaccaattc tgatgacatc tttacg

36

<210> 130

<211> 25

<212> DNA

<213> Artificial DNA

<400> 130

gaaaattgtg ctctgttgta atcca

25

<210> 131

<211> 29

<212> DNA

<213> Artificial DNA

<400> 131

gccataacaa ggtcttagtg ataatagtg

29

<210> 132

<211> 24

<212> DNA

<213> Artificial DNA

<400> 132

tggactgggtt cctggatcgc ctcc

24

<210> 133

<211> 21

<212> DNA

<213> Artificial DNA

<400> 133

cactgacgac atcacgctca t

21

<210> 134

<211> 20

<212> DNA

<213> Artificial DNA

<400> 134

cccagtacgt tggtggtggt

20

<210> 135

<211> 31

<212> DNA

<213> Artificial DNA

<400> 135

aacctcaaac agtgcaagat gtctctgaac g

31

<210> 136

<211> 21

<212> DNA

<213> Artificial DNA

<400> 136

ctgtgacaag catggcctgt a

21

<210> 137

<211> 16

<212> DNA

<213> Artificial DNA

<400> 137

gcactcccca cgctgc

16

<210> 138

<211> 21

<212> DNA

<213> Artificial DNA

<400> 138

ccatgggcagc ccttttggtgc a

21

<210> 139

<211> 21

<212> DNA

<213> Artificial DNA

<400> 139

acctgcgttc ctctctctagc t

21

<210> 140

<211> 22

<212> DNA

<213> Artificial DNA

<400> 140

tggtctagaa gtgcagccca tt

22

<210> 141

<211> 24

<212> DNA

<213> Artificial DNA

<400> 141

gaacagcttc cctcactgtg taca

24

<210> 142

<211> 25

<212> DNA

<213> Artificial DNA

<400> 142

attcctgctt ttcagagtga gacac

25

<210> 143

<211> 29

<212> DNA

<213> Artificial DNA

<400> 143

acgttcagac cacctattcc cttcttgcg

29

<210> 144

<211> 21

<212> DNA

<213> Artificial DNA

<400> 144

tctttggcta ccagtccagc a

21

<210> 145

<211> 20

<212> DNA

<213> Artificial DNA

<400> 145

aaaggcccga gtatccctga

20

<210> 146

<211> 29

<212> DNA

<213> Artificial DNA

<400> 146

caagtcaccc agtcctactg ggacaccaa

29

<210> 147

<211> 22

<212> DNA

<213> Artificial DNA

<400> 147

tctaggactt ctgcacggac ct

22

<210> 148

<211> 20

<212> DNA

<213> Artificial DNA

<400> 148

aggtccggac tggtcgagat

20

<210> 149

<211> 27

<212> DNA

<213> Artificial DNA

<400> 149

cagtgccaac ttcattccca cggtcac

27

<210> 150

<211> 22

<212> DNA

<213> Artificial DNA

<400> 150

cagctacttt tctgggcagg gc

22

<210> 151

<211> 20

<212> DNA

<213> Artificial DNA

<400> 151

acaatcaggc gttccagctc

20

<210> 152

<211> 24

<212> DNA

<213> Artificial DNA

<400> 152

cgcgtctctc aagctcgctt cttc

24

<210> 153

<211> 25

<212> DNA

<213> Artificial DNA

<400> 153

tggctgaagg gaattaacag atagt

25



<210> 154

<211> 27

<212> DNA

<213> Artificial DNA

<400> 154

ttcatgttaa tccattctga gtcattt

27

<210> 155

<211> 26

<212> DNA

<213> Artificial DNA

<400> 155

tcacccaggg agccatactg gcacat

26

<210> 156

<211> 20

<212> DNA

<213> Artificial DNA

<400> 156

gcagaggcga agcatcatct

20

<210> 157

<211> 21

<212> DNA

<213> Artificial DNA

<400> 157

cagcacctgg gactcaaact g

21

<210> 158

<211> 25

<212> DNA

<213> Artificial DNA

<400> 158

ccaacttcag cttcatgggc cagct

25

<210> 159

<211> 18

<212> DNA

<213> Artificial DNA

<400> 159

tgctgctcct gctcctgg

18

<210> 160

<211> 20

<212> DNA

<213> Artificial DNA

<400> 160

aggtgaattc cctgcagggt

20

<210> 161

<211> 25

<212> DNA

<213> Artificial DNA

<400> 161

cactgaactg cgctgccagt gcttg

25

<210> 162

<211> 19

<212> DNA

<213> Artificial DNA

<400> 162

gagcgtccgt ggtcactga

19

<210> 163

<211> 25

<212> DNA

<213> Artificial DNA

<400> 163

acattcacac tttggatgtt cttga

25

<210> 164

<211> 26

<212> DNA

<213> Artificial DNA

<400> 164

tgaattccct gcagtgtctg caagca

26

<210> 165

<211> 20

<212> DNA

<213> Artificial DNA

<400> 165

tgccagctat gtggtagggc

20

<210> 166

<211> 20

<212> DNA

<213> Artificial DNA

<400> 166

gtcaaccgtg ggatttcctg

20

<210> 167

<211> 31

<212> DNA

<213> Artificial DNA

<400> 167

cctggccttt cagttacttg agtttggagt c

31

<210> 168

<211> 22

<212> DNA

<213> Artificial DNA

<400> 168

cgtgcaaagt cacacacctc at

22

<210> 169

<211> 23

<212> DNA

<213> Artificial DNA

<400> 169

ggtgttggtg aagtaggtga tgg

23

<210> 170

<211> 25

<212> DNA

<213> Artificial DNA

<400> 170

cttcctcacg tgcattgagcg tggac

25

<210> 171

<211> 19

<212> DNA

<213> Artificial DNA

<400> 171

cctttgaagg gcaggcaca

19

<210> 172

<211> 20

<212> DNA

<213> Artificial DNA

<400> 172

gcaggctcga tgctgttcat

20

<210> 173

<211> 27

<212> DNA

<213> Artificial DNA

<400> 173

tccatgacgc cgtcctttcc agatcta

27

<210> 174

<211> 24

<212> DNA

<213> Artificial DNA

<400> 174

ctgagaagca tgacggacaa gtac

24

<210> 175

<211> 22

<212> DNA

<213> Artificial DNA

<400> 175

aggggaagcgt gatgacaaag ag

22

<210> 176

<211> 24

<212> DNA

<213> Artificial DNA

<400> 176

ctgcacctgt cagtggccga cctc

24

<210> 177

<211> 22

<212> DNA

<213> Artificial DNA

<400> 177

aggaacaggg aacattggag ct

22

<210> 178

<211> 19

<212> DNA

<213> Artificial DNA

<400> 178

ttggtgaggt ggggtgggat

19

<210> 179

<211> 33

<212> DNA

<213> Artificial DNA

<400> 179

aactgattac ctgccaattg ctaccgagaa ggt

33

<210> 180

<211> 20

<212> DNA

<213> Artificial DNA

<400> 180

agaggaggtc gcagttcagc

20

<210> 181

<211> 22

<212> DNA

<213> Artificial DNA

<400> 181

cagtacccca gtgtcatgga ca

22

<210> 182

<211> 27

<212> DNA

<213> Artificial DNA

<400> 182

tctgcagaat cgacaccaac cagcatc

27



CLAIMS

1. A method for the prediction, diagnosis or prognosis of chronic lung disease by the detection of:

5

a) a polynucleotide comprising at least one of the sequences of SEQ ID NO. 1 to 28;

10

b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;

15

c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;

20

d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

in a biological sample comprising the following steps:

hybridizing at least one polynucleotide specified in (a) to (d) to a nucleic acid material of a biological sample, thereby forming a hybridization complex; and

25

detecting said hybridization complex.

2. The method of claim 1, wherein before hybridization, the nucleic acid material of the biological sample is amplified.

30

3. A method for the prediction, diagnosis or prognosis of chronic lung disease by the detection of:
- a) a polynucleotide comprising at least one of the sequences of SEQ ID NO. 1 to 28;
  - b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;
  - c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;
  - d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);
  - e) a purified polypeptide encoded by a polynucleotide sequence specified in (a) to (d);
  - f) a purified polypeptide comprising at least one of the sequences of SEQ ID NO. 29 to 56;
- comprising the steps of contacting a biological sample with a reagent which specifically interacts with the polynucleotide specified in (a) to (d) or the polypeptide specified in (e) and (f).
4. A diagnostic kit for conducting the method of anyone of claims 1 to 3.

5. A method of screening for agents which regulate the activity of a polypeptide encoded by a polynucleotide selected from the group consisting of:

- 5 a) a polynucleotide comprising at least one of the sequences of SEQ ID NO. 1 to 28;
- 10 b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;
- 15 c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;
- d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

comprising the steps of:

20

contacting a test compound with at least one polypeptide encoded by a polynucleotide specified in (a) to (d); and

25

detecting binding of the test compound to the polypeptide, wherein a test compound which binds to the polypeptide is identified as a potential therapeutic agent for modulating the activity of the polypeptide in order to prevent or treat chronic lung disease.

6. A method of screening for agents which regulate the activity of a polypeptide encoded by a polynucleotide selected from the group consisting of:

30

- a) a polynucleotide comprising at least one of the sequences of SEQ ID NO. 1 to 28;
- b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;
- c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;
- d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

comprising the steps of:

contacting a test compound with at least one polypeptide encoded by a polynucleotide specified in (a) to (d); and

detecting the activity of the polypeptide as specified for the respective sequence in Table 3, wherein a test compound which increases the activity is identified as a potential preventive or therapeutic agent for increasing the in chronic lung disease, and wherein a test compound which decreases the activity of the polypeptide is identified as a potential therapeutic agent for decreasing the activity in chronic lung disease.

7. A method of screening for agents which regulate the activity of a polynucleotide selected from group consisting of;

- 200 -

- 5
- a) a polynucleotide comprising at least one of the sequences of SEQ ID NO. 1 to 28;
- b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;
- 10 c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;
- d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);
- 15
- comprising the steps of:
- contacting a test compound with at least one polynucleotide specified in (a) to (d), and
- 20
- detecting binding of the test compound to the polynucleotide, wherein a test compound which binds to the polynucleotide is identified as a potential preventive or therapeutic agent for regulating the activity of the polynucleotide in chronic lung disease.

25

## 8. Use of

- a) a polynucleotide comprising at least one of the sequences of SEQ ID NO. 1 to 28;

30

- 5
- b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) and encodes a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;
- c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;
- 10 d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);
- e) an antisense molecule targeting one of the polynucleotide sequences specified in (a) to (d);
- 15 f) a purified polypeptide encoded by a polynucleotide sequence specified in (a) to (d)
- g) a purified polypeptide comprising at least one of the sequences of SEQ ID NO. 29 to 56;
- 20 h) an antibody capable of binding to one of the polynucleotide specified in (a) to (d) or a polypeptide specified in (f) and (g)
- 25 i) a reagent identified by any of the methods of claim 5 to 7 that modulates the amount or activity of a polynucleotide sequence specified in (a) to (d) or a polypeptide specified in (f) and (g)
- 30 for the prevention, prediction, diagnosis, prognosis and treatment of chronic lung disease.

9. Use of claim 8 wherein the disease is COPD.

10. A reagent that regulates the activity of a polynucleotide selected from the group consisting of:

5

a) a polynucleotide comprising at least one of the sequences of SEQ ID NO. 1 to 28;

10

b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;

15

c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;

20

d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

or a polypeptide encoded by at least one of the polynucleotides specified in (a) to (d). wherein said reagent is identified by the method of any of the claims 5 to 7.

25

11. A pharmaceutical composition, comprising:

an expression vector containing at least one polynucleotide selected from the group consisting of:

30

a) a polynucleotide comprising at least one of the sequences of SEQ ID NO. 1 to 28;

5           b)     a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 3;

          c)     a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the degeneracy of the genetic code;

10

          d)     a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

or the reagent of claim 10 and a pharmaceutically acceptable carrier.

15

12.     A method of treating or preventing chronic lung disease by administering to a mammal in need thereof an agent identified by the method of claim 5, 6 or 7.



Table 1: Genes, up-regulated in COPD patients

No	Accession- No	Fold Change		Description
		Arrays	TaqMan	
1	L 13463	25,2	21,0	Cluster Incl L13463:Human helix-loop-helix basic phosphoprotein (GOS8) mRNA, complete cds
2	X 14787	15,5	16,4	X14787 //DEFINITION=HSTS Human mRNA for thrombospondin
3	K 00650	15,1	5,4	K00650 //DEFINITION=HUMFOS Human fos proto-oncogene (c-fos), complete cds
4	M 29039	6,2	4,3	M29039 //DEFINITION=HUMJUNCAA Human transactivator (jun-B) gene, complete cds
5	S 59049	5,9	12,6	Cluster Incl S59049:BL34=B cell activation gene [human, mRNA, 1398 nt]
6	X 68277	4,4	5,2	X68277//HSCCL100 H.sapiens CL 100 mRNA for protein tyrosine phosphatase
7	M 36820	4,3	5,5	Cluster Incl M36820:Human cytokine (GRO-beta) mRNA, complete cds
8	M 36821	3,8	5,3	Cluster Incl M36821:Human cytokine (GRO-gamma) mRNA, complete cds
9	X02530	3,5	7,8	X02530 /Human mRNA for gamma-interferon inducible early response gene (with homology to platelet proteins)
10	L 20971	3,3	4,1	Cluster Incl L20971:Human phosphodiesterase mRNA, complete cds
11	L 22214	3,3	3,4	Cluster Incl L22214:Human adenosine A1 receptor (ADORA1) mRNA exons 1-6, complete cds
12	U 67784	3,1	4,4	Cluster Incl U67784:Human orphan G protein-coupled receptor (RDC1) mRNA, partial cds
13	V 01512	3,0	5,9	V01512 //DEFINITION=HSCFOS Human cellular oncogene c-fos (complete sequence)
14	AL 021154	2,6	5,1	Cluster Incl AL021154:(Inhibitor of DNA binding 3 (dominant negative helix-loop-helix protein, 1R21, HEIR-1))
15	M 20681	2,4	2,1	Cluster Incl M20681:Human glucose transporter-like protein-III (GLUT3), complete cds
16	L 06797	2,4	2,9	L06797/HUMGPCR Human (clone L5) orphan G protein-coupled receptor mRNA, complete cds
17	M 60278	2,1	2,2	Cluster Incl M60278:Human heparin-binding EGF-like growth factor mRNA, complete cds
18	M 96233	2,0	1,4	M96233 /Human glutathione transferase class mu number 4 (GSTM4) gene, complete cds

Table 2: Genes, down-regulated in COPD patients

No	Accession- No	Fold Change		Description
		Arrays	TaqMan	
19	U03688	-8.7	-8.11	Cluster Incl U03688:Human dioxin-inducible cytochrome P450 (CYP1B1) mRNA, complete cds
20	AF 052124	-5.9	-5,2	Cluster Incl AF052124:Homo sapiens osteopontin mRNA, complete cds
21	M 28225	-3.8	-6,9	Cluster Incl M28225:Human JE gene encoding a monocyte secretory protein
22	L 23808	-3.2	-4,4	L23808 /DEFINITION=HUMHME Human metalloproteinase (HME) mRNA, complete cds
23	AA 883870	-2.4	-1,8	Cluster Incl AA883870, similar to gb:M29844 APOLIPOPROTEIN C-II PRECURSOR (HUMAN)
24	AF 012024	-2.1	-1.0	AF012024/ integrin cytoplasmic domain associated protein (Icap-1b) mRNA, complete cds
25	AB 13382	-2.0	-2,0	Cluster Incl AB013382:Homo sapiens mRNA for DUSP6, complete cds
26	D 26443	-1.98	-2,0	Cluster Incl D26443:Human mRNA for glutamate transporter, complete cds
27	X16302	-1.51	-1.24	Cluster Incl X16302:Human mRNA for insulin-like growth factor binding protein (IGFBP-2)
28	M26683	-1.51	-10.13	M26683 / Human interferon gamma treatment inducible mRNA

Table 3: Biochemical function of the „COPD GENE“

No	mRNA Accession-No	Protein Accession-No	Biochemical Function of the „COPD GENE“
1	L 13463	AAC37587	GTPase activating protein, Regulator of G-protein signaling 2, one of a family of proteins that negatively regulates G protein-coupled receptor signaling by binding to G-protein alpha subunits and stimulating their intrinsic GTPase activity
2	X 14787	CAA32889	Inhibitor or repressor; Adhesin/agglutinin, Thrombospondin-1, a member of a family of adhesive molecules, expressed in platelets and in extracellular matrix of developing and repairing tissues, has a role in blood clotting and in angiogenesis
3	K 00650	AAA52471	Transcription factor
4	M 29039	AAA59198	Transcription factor, Proto-oncoprotein with similarity to Jun that participates in AP-1 transcriptional activation and in malignant transformation, when coexpressed with Jun may act as an inhibitor of Jun-mediated activation and transformation
5	S 59049	AAB26289	GTPase activating protein, Regulator of G-protein signaling 1, one of a family of proteins that negatively regulates G protein-coupled receptor signaling by binding to G-protein alpha subunits and stimulating their intrinsic GTPase activity, expressed specifically in B lymphocytes
6	X 68277	CAA48338	Protein phosphatase, Non-receptor protein-tyrosine phosphatase that has similarity to dual specificity serine/tyrosine phosphatases, upregulated by oxidative stress and heat shock
7	M 36820	AAA63183	Ligand, Chemotactic agent for human polymorphonuclear leukocytes, has strong similarity to human GRO1, which is a mitogenic factor involved in inflammatory processes and that has growth-regulatory properties

Table 3: (continued)

No	mRNA Accession-No	Protein Accession-No	Biochemical Function of the „COPD GENE“
8	M 36821	AAA63184	Ligand, Paralog of human GRO1, which is a mitogenic factor involved in inflammatory processes and that has growth-regulatory properties
9	X02530	CAA26370	Protein kinase, IFN-gamma-inducible protein, a member of the C-X-C chemokine family with homology to platelet factor 4 and beta-thromboglobulin
10	L 20971	AAA03593	Hydrolase, cAMP-specific phosphodiesterase that is sensitive to the antidepressant rolipram
11	L 22214	AAA17544	Receptor (signalling), Adenosine A1 receptor, a G protein-coupled receptor that selectively binds adenosine, stimulates cell death of thymocytes and phagocytosis
12	U 67784	AAB16913	Receptor (signalling), Protein with strong similarity to murine Cmkor1, which is a member of the G protein-coupled receptor family and related to chemokine receptors of the CXC family
13	V 01512	CAA24756	Transcription factor, Transcription factor that forms a heterodimer with c-Jun and stimulates transcription of genes containing activator protein-1 (AP-1) regulatory elements,
14	AL 021154, X66924	CAA48862	Transcription factor; Inhibitor or repressor, Member of the Id helix-loop-helix family of proteins that negatively regulate cell differentiation by inhibiting helix-loop-helix transcription factors via a dominant-negative mechanism
15	M 20681	AAB61083	Major Facilitator Superfamily; Transporter, Facilitated glucose transporter
16	L 06797	AAA03209	Receptor (signalling), CXC chemokine receptor (fusin), a G protein-coupled receptor that binds CXC cytokines and mediates intracellular calcium flux
17	M 60278	AAA35956	Ligand, Heparin-binding EGF-like growth factor that binds to EGF receptors and TM4 superfamily members
18	M 96233	AAA57346	Transferase, Member of the mu class of glutathione S-transferases, a family of detoxification enzymes that catalyzes the conjugation of glutathione to electrophilic compounds

Table 3: (continued)

No	mRNA Accession-No	Protein Accession-No	Biochemical Function of the „COPD-GENE“
19	U03688	AAA19567	Oxidoreductase; Transporter, Cytochrome P450 1B1
20	AF 052124	AAC28619	Extracellular matrix protein
21	M 28225	AAA60308	Ligand, Human JE gene encoding a monocyte secretory protein
22	L 23808	AAA58658	Protease, Matrix metalloprotease that degrades elastin
23	AA 883870	XP_008846	Hydrolase, Apolipoprotein CII, cofactor for lipoprotein lipase which hydrolyzes triglyceride-rich lipoproteins
24	AF 012024	AAB88672	Adhesin/agglutinin, Protein that interacts with the cytoplasmic domain of beta1 integrin and has a role in cell signaling
25	AB 013382	BAA34369	Protein phosphatase, Dual specificity protein phosphatase, a constitutively expressed cytosolic protein that selectively dephosphorylates and inactivates mitogen-activated protein kinase
26	D 26443	BAA05462	Transporter, High affinity sodium-dependent glutamate/aspartate transporter, important for reuptake of glutamate in excitatory neurotransmission
27	X16302	CAA34373	Structural protein, Insulin-like growth factor binding protein that binds to and modulates insulin-like growth factor activity
28	M26683	AAA96708	Ligand, Cytokine A 2, a chemotactic factor for monocytes

Table 4: Primer and Probe Sequences for TaqMan Analyses

Genes, up-regulated in COPD patients

No	Accession-No	PRIMER AND PROBES SEQUENCES 5' > 3'
1	L 13463	Probe forward primer reverse primer ACGTTCAGACCACCTATTCCCTTCTTGCG ATTCCTGCTTTTCAGAGTGAGACA GAACAGCTTCCCTCACTGTGTACA
2	X 14787	Probe forward primer reverse primer TCGCCAGGCCCTGTGGTGG AGGGCGTTCGCGAGGT GGCCCTTCTGTGAAAGTTGTAA
3	K 00650	Probe forward primer reverse primer TTCCCAGCATCATCCAGGCCCCAG CGAGCCCTTTGATGACTTCCT GGCTCCCAGTCTGCTGCATA
4	M 29039	Probe forward primer reverse primer CGCGTCTCTCAAGCTCGCCTCTTC CAGTACTTTTCTGGTCAGGGC ACAAATCAGGCGTTCAGCTC
5	S 59049	Probe forward primer reverse primer ACAGTGGAGCTCTGTATTAGAAAGCCCTCA GAGGAAGATACTGTGGTACTGTCAATAAAA GAGTTACCTGGCCTTCCCAGTT
6	X 68277	Probe forward primer reverse primer CCAACTTCAGCTTCATGGGCCAGCT GCAGAGGCGAAGCATCATCT CAGCACCTGGGACTCAAACTG
7	M 36820	Probe forward primer reverse primer TGCCAGTGTGTCAGACCCCTGC CCTGGCCACTGAACCTGCG TGGATGTTCTTGAGGTGAATTCC
8	M 36821	Probe forward primer reverse primer TGAATTCCCTGCAGTGTCTGCAAGCA ACATTACACTTTGGATGTTCTTTGA GAGCGTCCGTGGTCACTGA

Table 4: (continued)

<u>No</u>	<u>Accession-No</u>	<u>PRIMER AND PROBES SEQUENCES 5' &gt; 3'</u>
9	X 02530	Probe forward primer reverse primer TTCTGACTCTAAGTGGCATTCAAGGAGTACCTCTC CGATTCTGATTTGCTGCTTATC GCAGGTACAGCGTACGGTTCT
10	L 20971	Probe forward primer reverse primer TGATGCGGTCTGTCCAATTGCCGATA TTGTCTCCCTGCTGGAAAAATT TGAGCAACCCCAACCAAGTC
11	L 22214	Probe forward primer reverse primer CCGCATCCAGAAGTTCCGCGTC GCAACTCGGCCATGAACC GGTCATTCCAAATCTTAAGGAAGG
12	U 67784	Probe forward primer reverse primer CCGGTCCCTTCTACCCCGAGCACA CTGCGTCCAACAATGAGACCTA CCGATCAGCCACTCCTTGAT
13	V 01512	Probe forward primer reverse primer AGGACCTTATCTGTGCGTGAAACACACCA CCTAGAGGGTTCTGTAGACCTAGG AGTCCTTGAGGCCACAGC
14	AL 021154	Probe forward primer reverse primer CGCCGCCCTTGGCATAAGTTTGA TCCAAGGAGACCAAGAACCA GTGTGGAAGGAGTGGCTGCT
15	M 20681	Probe forward primer reverse primer TAGATCTGGAAGGACGGCGTCATGGA GCAGGCACACGGTGCA GCAGGCTCGATGCTGTTTCAT
16	L 06797	Probe forward primer reverse primer CTCCAAAGGAAAGCGAGGTGGACATTTCAT GGGTCCAGCCTCAAGATCC ACTTGAAGACTCAGACTCAGTGGAAA
17	M 60278	Probe forward primer reverse primer CGTGTCCCTCTCCCTGCCAAGTCTCAG GTCAAAGTGTAACAGATATCAGTGTCTCC TACAGGCATGGAAGCCCAAC
18	M 96233	Probe forward primer reverse primer CCTACAATGATGCAGCACTTCTCACAGTTCC GAAGCCAGAATACTTGGAGGAACT ATACGGTGGAGGTCAAGGACAT

Table 5

Genes, down-regulated in COPD patients

No	Accession-No	PRIMER AND PROBES SEQUENCES 5' > 3'
19	U 03688	Probe forward primer reverse primer ATTCCTCATGCCACCACCTGCCAACA TCCAGCTTTGTGCCTGTCCAC GGGAATGTGGTAGCCCAAGA
20	AF 052124	Probe forward primer reverse primer TCACCTCACACATGGAAGCGAGGAGT CCAGTACCTGTGATGCTACAGACG TGGCCTTGATGCACCATTC
21	M 28225	Probe forward primer reverse primer ACCAAACCTCCGAAGACTTGAACACCCAC TGGACCACCTGGACAAGCA GCTGCAGATTCTTGGGTGTG
22	L 23808	Probe forward primer reverse primer CCTTATGGCCAAACCTTGCCATCTGG CTGAGAGACCAAGACCAAGTGTAAAT TTCTGGCTTCAATTTCATAAGCA
23	AA 883870	Probe forward primer reverse primer ACAGCCGCCAGAACCTGTACGAGA TCCAGTTACTGGGAGTCAGCAA TCTACAGCGGGCAGGTATGTC
24	AF 012024	Probe forward primer reverse primer TTCAGATCCAGCACTGTGGCCA AGCAAGTCTGTGGATTCTAGCCTT CCTGAGCTTTTGGTGGAAATCTG
25	AB 13382	Probe forward primer reverse primer ATGCTTGACTTTACCAATTCCTGATGACATCTTTACG GAAAATTGTGCTCTGTGTGTAATCCA GCCATAACAAGGTCTTAGTGATAATAGTG
26	D 26443	Probe forward primer reverse primer TGGACTGGTTCCCTGGATCGCCTCC CACTGACGACATCAGCTCAT CCCAGTACGTTGGTGGTGGT
27	X 16302	Probe forward primer reverse primer AACCTCAAACAGTGCAAGATGTCTCTGAACG CTGTGACAAGCATGGCCCTGTA GCACTCCCCACGCTGC
28	M 26683	Probe forward primer reverse primer CCATGGCAGCCCTTTGGTGCA ACCTGCGTTCTCTCCTCTAGCT TGGTCTAGAAAGTGCAGCCCATTT



Table 6

No	Accession No.	TaqMan at SC	Fold Change	Primer and Probes sequences 5' > 3'	Forward Concen- tration (nM)	Reverse Concen- tration (nM)	Probe Concen- tration (nM)
1	L13463	YES	1,86	Forward Sequence Reverse Sequence Probe Sequence GAACAGCTTCCCTCACTGTGTACA ATTCCTGCTTTTCAGAGTGAGACAC ACGTTACAGACCACCTATTCCCTTCTTGCG	900	900	200
2	X14787	YES	20,45	Forward Sequence Reverse Sequence Probe Sequence TCTTTGGGTACCACTCCAGCA AAAGGCCCGAGTATCCCTGA CAAGTCACCCAGTCTTACTGGGACACCAA	900	900	200
3	K00650	YES	8,79	Forward Sequence Reverse Sequence Probe Sequence TCTAGGACTTCTGCACGGACCT AGGTCCGGACTGGTCGAGAT CAGTGCCAACTTCATTCCCACGGTCAC	900	900	200
4	M29039	YES	2,5	Forward Sequence Reverse Sequence Probe Sequence CAGTACTTTTCTGGTCAGGGC ACAAATCAGGCGTTCAGCTC CGCGTCTCTCAAGCTCGCCTCTTC	900	900	200
5	S59049	YES	4,79	Forward Sequence Reverse Sequence Probe Sequence TGGCTGAAGGGAATTAACAGATAGT TTTCAATGTTAATCCATTCTGAGTCATTT TCACCCAGGGAGCCATATACTGGCACAT	900	900	200
6	X68277	YES	1,52	Forward Sequence Reverse Sequence Probe Sequence GCAGAGGCGAAGCATCATCT CAGCACCTGGGACTCAAACTG CCAACTTCAGCTTCATGGGCCAGCT	900	900	200
7	M36820	YES	2,95	Forward Sequence Reverse Sequence Probe Sequence TGCTGCTCTGCTCTCTGG AGGTGAATTCCTTGCAGGGT CACTGAACTGCGCTGCCAGTGCTTG	900	900	200
8	M36821	YES	2,01	Forward Sequence Reverse Sequence Probe Sequence GAGCGTCGTTGGTCACTGA ACATTACACTTTGGATGTTCTTGA TGAAATTCCTTGCACTGTCTGCAAGCA	900	900	200
10	L20971	YES	1,39	Forward Sequence Reverse Sequence Probe Sequence TGCCAGCTATGTGGTAGGGC GTCAACCGTGGGATTTTCCTG CCTGGCCTTTTCACTTACTTGAGTTGGAGTC	50	50	175

Table 6: (continued)

No	Accession No.	TaqMan at SC	Fold Change	Primer and Probes sequences 5' > 3'	Forward Concen- tration (nM)	Reverse Concen- tration (nM)	Probe Concen- tration (nM)
12	U67784	YES	11,54	Forward Sequence Reverse Sequence Probe Sequence CGTGCAAGTCACACACCTCAT GGTGTGGTGAAGTAGGTGATGG CTTCTCAGGTGCATGAGCGTGGAC	900	900	200
15	M20681	YES	26,96	Forward Sequence Reverse Sequence Probe Sequence CCTTTGAAGGGCAGGCACA GCAGGCTCGATGCTGTTTCAT TCCATGACGCCGTCCTTTCCAGATCTA	900	900	200
16	L06797	YES	2,24	Forward Sequence Reverse Sequence Probe Sequence CTGAGAACATGACGGACAAGTAC AGGAAAGCGTGATGACAAAGAG CTGCACCTGTCAGTGGCCGACCTC	900	900	200
17	M60278	YES	27,48	Forward Sequence Reverse Sequence Probe Sequence AGGAACACAGGGAACATTGGAGCT TTGGTGAGGTGGGTGGGAT AACTGATTACCTGCCAATTGCTACCGAGAAGGT	900	900	200
18	M96233	YES	7,7	Forward Sequence Reverse Sequence Probe Sequence AGAGGAGGTCCGAGTTCAGC CAGTACCCAGTGTTCATGGACA TCTGCAGAATCGACACCAACCAGCATC	900	900	200

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
5 December 2002 (05.12.2002)

PCT

(10) International Publication Number  
**WO 02/097127 A3**

(51) International Patent Classification<sup>7</sup>: **C12Q 1/68**,  
G01N 33/48, C07K 14/47, A61K 48/00

(21) International Application Number: PCT/EP02/05835

(22) International Filing Date: 28 May 2002 (28.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0113266.1 31 May 2001 (31.05.2001) GB

(71) Applicant (for all designated States except US): **BAYER AKTIENGESELLSCHAFT** [DE/DE]; 51368 Leverkusen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **OELLERS, Nadja** [DE/DE]; Steinfeldergasse 9, 50670 Köln (DE). **GEHRMANN, Mathias** [DE/DE]; Rosenstr. 68, 47918 Tönisvorst (DE). **KALLABIS, Harald** [DE/DE]; Wolfskaul 8, 51061 Köln (DE). **HALL, Roderick** [GB/GB]; 10 Acacia Close, Woodham, Surrey KT 15 3SJ (GB). **SCHULZE, Thomas** [DE/DE]; Kaiserwaldstr. 2c, 91083 Baiersdorf (DE). **KROEGEL, Claus** [DE/DE]; Falkenweg 18, 07751 Cospeda (DE).

(74) Common Representative: **BAYER AKTIENGESELLSCHAFT**; 51368 Leverkusen (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,

LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

**Published:**

— with international search report

(88) Date of publication of the international search report:  
21 August 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GENES AND PROTEINS FOR PREVENTION, PREDICTION, DIAGNOSIS, PROGNOSIS AND TREATMENT OF CHRONIC LUNG DISEASE

(57) Abstract: Genes that are differentially expressed in lung tissue of COPD patients versus lungs of normal people are disclosed. The genes provide novel methods for the prevention, prediction, diagnosis, prognosis and treatment of chronic lung disease.



WO 02/097127 A3

## INTERNATIONAL SEARCH REPORT

International Application No.

T/EP 02/05835

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12Q1/68 G01N33/48 C07K14/47 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

SEQUENCE SEARCH, EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, EMBASE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 55174 A (HUMAN GENOME SCIENCES INC ;ROSEN CRAIG A (US); RUBEN STEVEN M (US)) 21 September 2000 (2000-09-21) Seq Id Nos 567 and 1507 page 3, line 3 - line 22 page 354, line 27 - line 29 page 365, line 19 -page 366, line 10 page 380, line 13 - line 17 claims 17-20 ---	1-12
X	US 5 955 314 A (GOLI SURYA K ET AL) 21 September 1999 (1999-09-21) Seq Id No 4 column 7, line 45 -column 8, line 13 abstract --- -/--	1-12

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&amp;" document member of the same patent family

Date of the actual completion of the international search

23 January 2003

Date of mailing of the international search report

13. 05. 2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

CEDER O.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 02/05835

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 854 191 A (SMITHKLINE BEECHAM CORP) 22 July 1998 (1998-07-22) abstract ---	1-12
A	EP 1 043 406 A (TEIJIN LTD) 11 October 2000 (2000-10-11) abstract -----	1-12

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP 02/05835

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: -  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.: -  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-12 all partly

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.1

Although claim 12 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

-----

## Continuation of Box I.1

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

-----

## Continuation of Box I.2

Present claim 8 relates to the use of a compound defined by reference to a desirable characteristic or property, namely it being identifiable by any of the methods of claims 5-7.

The claims cover all compounds having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compound by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compounds identified in parts a) - h) of claim 8, and only as far as the search fees have been paid.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

Invention 1: Claims 1-12 all partly

A method for prevention, prediction, diagnosis and treatment of chronic lung disease utilizing the polynucleotide of Seq Id No 1 or the polypeptide of Seq Id No 29, a method for screening for agents which regulates the polypeptide/polynucleotide or a pharmaceutical composition containing them.

Inventions 2-28: Claims 1-12 all partly

Idem as for invention 1 but for polynucleotide 2-28 and polypeptide 30-56, respectively.



## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

/EP 02/05835

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0055174	A	21-09-2000	AU 3395900 A	04-10-2000
			AU 3617600 A	04-10-2000
			AU 3617700 A	04-10-2000
			AU 3619400 A	04-10-2000
			AU 3619500 A	04-10-2000
			AU 3869400 A	04-10-2000
			CA 2364567 A1	21-09-2000
			CA 2364590 A1	21-09-2000
			CA 2364629 A1	21-09-2000
			CA 2366130 A1	21-09-2000
			CA 2366174 A1	21-09-2000
			CA 2366195 A1	21-09-2000
			EP 1168917 A2	09-01-2002
			EP 1165588 A1	02-01-2002
			EP 1169469 A1	09-01-2002
			EP 1165589 A1	02-01-2002
			EP 1159420 A1	05-12-2001
			EP 1163358 A1	19-12-2001
			JP 2003512815 T	08-04-2003
			JP 2003512816 T	08-04-2003
			WO 0055173 A1	21-09-2000
			WO 0055350 A1	21-09-2000
			WO 0055351 A1	21-09-2000
			WO 0055180 A2	21-09-2000
			WO 0055174 A1	21-09-2000
			WO 0055320 A1	21-09-2000
			US 2003054421 A1	20-03-2003
			US 2002081659 A1	27-06-2002
			US 2002039764 A1	04-04-2002
			US 2002055627 A1	09-05-2002
			US 2002151681 A1	17-10-2002
			US 2002052308 A1	02-05-2002
			US 2002044941 A1	18-04-2002
US 5955314	A	21-09-1999	AU 5238398 A	29-05-1998
			EP 0958363 A1	24-11-1999
			JP 2001527522 T	25-12-2001
			WO 9820128 A1	14-05-1998
			US 2002034777 A1	21-03-2002
EP 0854191	A	22-07-1998	US 6008017 A	28-12-1999
			EP 0854191 A2	22-07-1998
			JP 11042092 A	16-02-1999
			JP 2002191375 A	09-07-2002
			US 6365715 B1	02-04-2002
EP 1043406	A	11-10-2000	AU 747991 B2	30-05-2002
			AU 1682399 A	05-07-1999
			CA 2315218 A1	24-06-1999
			EP 1043406 A1	11-10-2000
			US 6346385 B1	12-02-2002
			CN 1285002 T	21-02-2001
			WO 9931271 A1	24-06-1999